



Investigation of ET changes in Synoptical stations of

h. sharifan, M. Mahmodian, and H. Ansari

University Gorgan, irrigation, Gorgan, Islamic Republic Of Iran (h_sharifan47@yahoo.com)

Investigation of ET changes in Synoptical stations of
Golestan region(Iran)

H. Sharifan(1) , M.Mahmodian(2) and H.Ansari(3)

(1) Department of water Eng., Gorgan University of Agricultural Sciences and
Natural Resources, Iran, (h_sharifan47@yahoo.com / Fax +98 171 4420981).

(2) meisam_mahmodian@yahoo.com

(3) Ferdoswi university of Mashhad/Iran, ansari_hos@yahoo.com

Abstract Evapotranspiration (ET) is one of the most important factors in irrigation systems designing. The different methods estimate ET which FAO-Penman-Monteith (F-P-M) method is one of the validest of methods which it estimates ET. Thus aim of this research was investigation of ET changes in synoptical stations on Golestan region/Iran (stations: Gorgan, Gonbad and Maraveh- Tappeh) in duration(1981-2005). Results shown that for yearly scale: ET values in Gorgan and Gonbad were increasing procedure exsept 1992-2000 duration, and in Maraveh- Tappeh station ET was decreasing procedure until 2000 year. But the later years were increasing procedure. In this manner for seasonly scale: ET maximum for autumn, winter and spring seasons, was happen in 1985. The maximum value of ET in Gorgan, Gonbad and Maraveh-Tappeh station were 1000, 1260 and 2000 mm/yr respectively and 510, 530 and 1000 mm/season respectively.

Keyword : Evapotranspiration, synoptical station, Golestan.