

Does Hydro and Osmo-Priming Improve Fennel (*Foeniculum vulgare*) Seeds Germination and Seedlings Growth?

Ehsan NEAMATOLLAHI¹, Mohammad BANNAYAN², Ali GHANBARI¹

Mohammad HAYDARI¹, Ali AHMADIAN¹

¹ Zabol University, Faculty of Agriculture, Department of Agronomy and Plant Breeding, Zabol, Iran; aghanbari@cic.aut.ac.ir

² Ferdowsi University of Mashhad, Faculty of Agriculture, Mashhad, Iran; bannayan@ferdowsi.um.ac.ir (corresponding author)

Abstract

This experiment was conducted to investigate the effects of hydropriming and osmopriming on germination rate, percentage of root–shoot length and root–shoot weight of fennel (*Foeniculum vulgare*) seeds. Priming was done by: hydropriming with distilled water, osmopriming with NaCl at four levels (-0.3, -0.6, -0.9, -1.2 MPa), osmopriming with K₂SO₄ in four levels (-0.3, -0.6, -0.9, -1.2 MPa), priming with PEG6000 in four levels (-0.3, -0.6, -0.9, -1.2MPa) and 5. seeds with unprime control at treat. In this study, RCBD experimental design was used for the analysis of experimental factors. The results showed that priming significantly effected at all treatment methods. Maximum and minimum germination percentage were obtained with PEG (-0.9 MPa) applied, and in control. Maximum and minimum germination rates were obtained when K₂SO₄ (-0.3 MPa), NaCl (-0.3 MPa) were used. Maximum and minimum root length were obtained when NaCl [(-1.2 MPa), PEG (-0.3 MPa)], NaCl (-0.6 MPa) were used. Maximum and minimum shoot length were obtained when PEG (-0.3 MPa), NaCl (-0.6 MPa) were used. Maximum and minimum root weight, root/shoot length were obtained when NaCl (-1.2 MPa), NaCl (-0.6 MPa) were used. Maximum and minimum shoot weight were obtained when NaCl (-1.2 MPa), NaCl (-0.3 MPa) were used. Maximum and minimum root/shoot weight were obtained when [PEG (-0.3 MPa), K₂SO₄ (-0.3, -0.6, -0.9 MPa), NaCl (-0.6, -0.9 MPa), hydropriming] and [PEG (-0.6, -1.2 MPa), NaCl (-1.2 MPa)] were used.

Keywords: fennel, *Foeniculum vulgare*, NaCl, K₂SO₄, PEG, hydropriming, osmopriming, germination

Introduction

Fennel (*Foeniculum vulgare*) is an aromatic biennial plant with soft, feathery, almost hair-like foliage. Native to coastal areas in the Mediterranean region, and widely naturalized in Europe and North America (Christman., 2004). Similar to some other medicinal plants (Bannayan *et al.*, 2008; Khazaie *et al.*, 2008) supplemental irrigation would improve its production during dry periods. Fennel belongs to the Umbelliferae (Apiaceae) family, a medicinal plant used as anti-spasmodic, appetite stimulant, stomachic, diuretic, anti-inflammatory, anti-diarrheic, against colic and as a lactation promoter (Marotti *et al.*, 1993; Piccaglia and Marotti, 1993; Cavaleiro *et al.*, 1993). Several components of the essential oil of this plant show important applications, including, fenchone as counterirritant; limonene as solvent, resins, wetting and dispersing agent; trans-anethole, flavoring agent in perfumery, cosmetics, soap; methylchavicol or estragole is used in perfumeries and as flavor in foods and liquors; α -pinene, used in manufacture of camphor, insecticides, solvents, perfume bases (Marotti *et al.*, 1993; Piccaglia and Marotti, 1993; Cavaleiro *et al.*, 1993). Seed germination is mostly an issue in medicinal plant seeds emergence (Nadjafi *et al.*, 2006). Good seedling establishment is an important constraint to such crop production

(Harris *et al.*, 1999). Poor seedbed, low quality seed, environmental stresses such as high and low temperature and salinity constrains to good establishment include (Weaich *et al.*, 1992; Towned *et al.*, 1996). A robust seedling establishment enhances competitiveness against weeds, improves tolerance to environmental stresses and maximizes biological and grain yields (Ghiyasi *et al.*, 2008).

Several approaches including, hardening, seed priming, seed soaking and seed coating have been employed to precondition seeds to improve germination and seedling growth of various crops (Basra *et al.*, 2003). Seed priming treatments such as osmopriming, hydropriming, matricopriming, hormonal-priming have been employed to accelerate the germination, seedling growth and yield in most of the crops under normal and stress conditions (Basra *et al.*, 2003). Osmopriming is most common type of seed priming in which seeds are soaked in aerated low water potential solution (Farooq *et al.*, 2005). Although, the mechanism of seed priming treatments is not fully understood, it has been observed that physiological and biochemical changes take place during the seed treatments (Basra *et al.*, 2005; Ghiyasi *et al.*, 2008), which could allow seeds to begin the germination sequences before sowing.

Rapid germination and emergence is an important determinant of successful establishment (Harris *et al.*, 1991).

Germination is one of the most salt-sensitive plant growth stages and severely inhibited with increasing salinity both in glycophytes and halophytes (Sosa *et al.*, 2005). Seed priming accelerates seed germination and seedling establishment under both normal and stressful environments (Ashraf and Foolad., 2005). Although priming is one of the physiological methods, which improves seed performance and provides faster and synchronised germination (Sivritepe and Dourado., 1995) it has been shown that NaCl priming could be used as an adaptation method to improve salt tolerance of seeds (Wiebe and Muhyaddin., 1987; Cano *et al.*, 1991; Cayuela *et al.*, 1996). Successful results have been obtained for tomatoes, (Pill *et al.*, 1991) and asparagus (Passam and Kakouriotis., 1994) under saline conditions.

Hydropriming is the simplest approach to hydrate seeds and minimize the use of chemicals. However, if the seeds are not accurately hydrated, hydration rate cannot be exactly controlled. It was observed that hydropriming practically ensured rapid and uniform germination accompanied with low abnormal seedling percentage (Singh, 1995; Shivankar *et al.*, 2003). They underline that hydropriming has high potential in improving field emergence and ensures early flowering and harvest under stress conditions especially in dry areas. Hydrated seeds with higher germination percentage under salt stress or micronutrient application increased tolerance of seeds to salt stress. In addition, reported protocol is simple, cheap and does not require expensive chemicals and sophisticated equipment. The protocol has practical importance and could be recommended to farmers to achieve higher germination and uniform emergence under field conditions.

The objective of this study was to explore the effects of two different priming (hydropriming and osmopriming) treatments on germination of fennel seeds (*Foeniculum vulgare*).

Materials and methods

In order to determine the impact of different primes on germination of Fennel seeds, an experiment was conducted at Zabol University in 2007. Seeds were primed with various materials, including hydroprime as seeds were primed with distilled water, osmopriming seeds were treated with three different chemical as NaCl, K₂SO₄, polyethylene glycol 6000. All three different osmopriming provided

osmotic potential of -0.3, -0.6, -0.9 and -1.2 MPa. Simulating various osmotic potential using PEG was according to Michel and Kaufmann (1973) and Money (1989).

Initially seeds were disinfected by sodium hypochlorite (NaOCl). Seeds were kept in sodium hypochlorite (1.5%) for one minute and then were washed with distilled water. After disinfecting, seeds were put in disinfected Petry dish. Each Petry dish contained 25 seeds. After 24 hours of priming seeds were washed with distilled water and then dried and kept in laboratory room at temperature of 25° C for two hours. Afterwards dried seeds were located in Petry dishes and treated with distilled water at temperature of 25°C for seven days. Statistical experimental design was randomized completely block, with three replications. The differences between the means were compared using Duncan test (P < 0.01).

Germination tests

Three replicates of 25 seeds were put between double layered rolled. The papers were replaced every 2 days to prevent accumulation of salts. The rolled paper with seeds was put into sealed plastic bags to avoid moisture loss. Seeds were allowed to germinate at 25 ± 1°C for 7 days. Germination was considered when the radicles were 2 mm long. Germination percentage was recorded every 24 h for 7 days. Root length and shoot length were measured after the 7th day. The germination rate was calculated as follows (according to Bannayan *et al.*, 2006):

$$\text{Germination rate} = \sum_{n=1}^{45}$$

number germinating since n-1 / n where, n is the days of incubation.

Results and discussion

For NaCl primed seeds the germination rate increased at osmotic pressure of -0.6 and -0.9 MPa, but germination rate decreased with increasing the osmotic pressure to -1.2 MPa (Tab. 1). The germination percentage of NaCl primed seeds decreased by increasing the osmotic pressure to -0.6 MPa, but germination percentage increased by increasing the osmotic pressure to -0.9 and -1.2 MPa. However, the best results of germination percentage in NaCl primed were obtained with -0.3 MPa pressure (Tab. 1). The root length of NaCl primed seeds decreased with increasing the osmotic pressure to -0.6 MPa, but root length increased


Osmotic level (Mpa)	Control		Hydropriming		NaCl		K ₂ SO ₄		PEG	
	G.Rate	G.Per	G.Rate	G.Per	G.Rate	G.Per	G.Rate	G.Per	G.Rate	G.Per
0	9.6 cd	4.3 g	9.3 cd	17.6bc						
-0.3					0.6 f	18.0b	12.0 a	17.6bc	9.7 cd	17.3b
-0.6					10.2cd	15.0f	9.1cde	17.6bc	11.9ab	16.6bcd
-0.9					9.5cd	16.0def	10.3bcd	17.3bcd	10.8abc	20.0a
-1.2					7.5e	17.3bcd	8.8de	15.6ef	9.3cd	16.3cdef

Tab. 2. Comparison of root length and shoot length of fennel seedling treated with NaCl, K₂SO₄ and PEG.

Osmotic level (Mpa)	Control		Hydropriming		NaCl		K ₂ SO ₄		PEG	
	R.length	S.length	R.length	S.length	R.length	S.length	R.length	S.length	R.length	S.length
0	5.7bc	2.9cd	6.3bc	3.4abcd						
-0.3					6.1bc	3.2abcd	6.2bc	3.9ab	8.2a	4.1a
-0.6					0.5d	0.3e	6.2bc	3.2abcd	5.0c	3.0bcd
-0.9					5.8bc	2.7d	5.2bc	2.7d	5.5bc	3.1bcd
-1.2					9.2a	3.9ab	6.5b	3.5abcd	6.2bc	3.6abc


at the osmotic pressure of -0.3, -0.9 and -1.2 MPa. Maximum root length was obtained at NaCl level providing -1.2 MPa osmotic pressure. This might be due to the fact that seedlings send their photosynthesis products to roots when they were exposed to salt stress in order to tolerate the imposed stress, and minimum root length was obtained at -0.6 MPa (Tab. 2). The shoot length at NaCl primed seeds decreased with increasing the osmotic pressure to -0.6 and -0.9 MPa, but shoot length increased as the osmotic pressure increased to -1.2 MPa (Tab. 2). Root weight and shoot weight of NaCl primed seeds increased with increasing the osmotic pressure, and maximum root weight and shoot weight was obtained at -1.2 MPa pressure (Tab. 3). Maximum and minimum root/shoot (R/S) length in NaCl primed seeds were obtained at -1.2 MPa and -0.6 MPa pressure, respectively. Maximum and minimum R/S weight in NaCl primed seeds were obtained at -0.9 MPa and -0.3 MPa pressure, respectively (Fig. 1, 2).

The maximum germination rate of K₂SO₄ primed seeds was obtained at -0.3 MPa pressure and beyond that pressure the germination rate decreased (Tab. 1). The germinated percentage of K₂SO₄ primed seeds decreased with increasing the osmotic pressure (Tab. 1). The root length of K₂SO₄ primed seeds increased with increasing the osmotic pressure (Tab. 2). Both shoot length and root weight showed significant effects of priming with K₂SO₄ at all potentials, except -0.9 MPa (Tab. 2, 3). Shoot weight of K₂SO₄ primed seeds decreased with increasing the osmotic pressure (Tab. 3). Maximum R/S length and R/S weight of K₂SO₄ primed seeds were obtained at -0.9 MPa pressure. Minimum R/S length and R/S weight in K₂SO₄ seeds primed were obtained at -0.3 MPa and -1.2 MPa pressures, respectively (Fig. 1, 2).

Fig. 1. Root/Shoot length (cm) of fennel seedling primed by hydropriming, NaCl, K₂SO₄, PEG and control

Using PEG priming, the germination rate increased at -0.6 and -0.9 MPa pressures (Tab. 1). The germination percentage of PEG primed seeds almost decreased by increasing solution concentration, except at -0.9 MPa pressure. Maximum germination percentage of PEG primed seeds was obtained at -0.9 MPa pressure and all seeds in Petry dishes were germinated (Tab. 1). The root length and shoot length of PEG primed seeds decreased with increasing osmotic pressure, and maximum root and shoot length were obtained at -0.3 MPa (Tab. 2). The root weight in PEG primed seeds generally decreased, but in PEG -0.6 MPa increased (Tab. 3). The shoot weight of PEG primed seeds decreased, except -0.9 MPa pressure, when increased again (Tab. 3). Maximum R/S weight and R/S length were obtained at -0.3 MPa pressure, and minimum R/S weight and R/S length were obtained at -0.9 MPa pressure (Fig. 1, 2). Several authors described positive effects of seed priming with water alone (Harris *et al.*, 1999, 2002, 2004; Rashid *et al.*, 2002). Seedlings from seeds primed with water alone are known to emerge more quickly and grow more vigorously than those from non-primed seeds (Rashid *et al.*, 2002; Arif *et al.*, 2005; Miraj, 2005). Our data confirmed that simple seed priming with water is an effective way to increase all characteristics compared to control treatment (Tab. 1, 2, 3 and Fig. 1, 2).

It is evident from the results that NaCl salinity caused growth inhibition in fennel seeds due to a decrease in total germination. These effects of salinity on fennel seeds supported by the previous findings (Franco *et al.*, 1993, 1997; Botia *et al.*, 1998; Carvajal *et al.*, 1998) studying on melon cultivars. Accumulation of Na ion changes ion balances such as Na:Ca and K:Na in plant cells under saline conditions. While the change in Na:Ca balance results in

Fig. 2. Root/Shoot weight (mg plant⁻¹) of fennel seedling primed by hydroprime, NaCl, K₂SO₄, PEG and control

Tab. 3. Root weight and shoot weight of fennel seedling treated with NaCl, K₂SO₄ and PEG

Osmotic level (Mpa)	Control		Hydropriming		NaCl		K ₂ SO ₄		PEG	
	R.weight	S.weight	R.weight	S.weight	R.weight	S.weight	R.weight	S.weight	R.weight	S.weight
0	0.6c	1.2abcd	0.8bc	1.0bcd						
-0.3					0.7bc	0.1e	0.7bc	1.1abcd	0.7bc	1.3abc
-0.6					0.1d	1.0cd	0.7bc	1.0cd	0.9ab	0.9d
-0.9					0.7bc	1.5ab	0.6c	1.0cd	0.5c	1.1abcd
-1.2					1.1a	1.5a	0.7bc	1.0bcd	0.7c	1.1bcd

increased cell permeability, the change in K:Na balance cause decreasing use of metabolic energy (Levitt, 1980).

Priming may improve germination by accelerating imbibition, which in turn would facilitate the emergence phase and the multiplication of radicle cells (McDonald, 1999). This process is important because allows the subsequent development of the embryo, especially in seeds characterised by a morphological dormancy (immature embryo), like *Chamaecyparis nootkatensis* seeds (Schimtz et al., 2001). In tomato, priming improved the germination capacity by increasing endosperm volume (Dahal et al., 1990).

The technique of seed priming is becoming familiar to farmers in several parts of the world, and has now been promoted there on a range of crops, for example wheat (Harris et al., 2001), maize (Harris et al., 2002), and mung bean (Rashid et al., 2004), where similar responses to those reported here have been found. Equally encouraging results have been found for these crops in other countries, and for other crops, such as chickpea in India and Bangladesh (Harris et al., 1999; Musa et al., 2001), upland rice in India (Harris et al., 1999, 2002), and finger millet in India (Kumar et al., 2002).

In many coated seeds, germination and subsequent seedling growth can be inhibited by mechanical restriction exerted by the seed coat (Sung and Chiu., 1995). Priming may be helpful in reducing the risk of poor stand establishment under drought and salt stress and permit more uniform growth under conditions of irregular rainfall and drought on saline soils.

References

- Arif, M., S. Ali, A. Shah, N. Javid, and A. Rashid (2005). Seed priming maize for improving emergence and seedling growth. *Sarhad J. Agric.* 21(4):539-543.
- Ashraf, M. and M. R. Foolad (2005). Pre-sowing seed treatment- a shotgun approach to improve germination, plant growth and crop yield under saline and non-saline conditions. *Adv. Agron.* 88:223-271.
- Bannayan, M., F. Nadjafi, M. Azizi, L. Tabrizi and M. Rastgoo (2008). Yield and seed quality of *Plantago ovata* and *Nigella sativa* under different irrigation treatments. *J. Ind. Crops and Prod.* 27:11-16.
- Basra, S. M. A., N. Zia, T. Mahmood, A. Afzal and A. Khaliq (2003). Comparison of different in vigorization techniques in wheat (*Triticum aestivum* L.) seeds. *Pak. J. Arid. Agric.* 5:11-16.
- Botia, P., M. Carvejal, A. Cerda and V. Martinez (1998). Response of eight *Cucumis melo* cultivars to salinity during germination and early vegetative growth. *Agronomic.* 18:503-513.
- Burget, K. L. and O. C. Burnside (1972). Optimum temperature for germination and seedling development of black night. *North Cent. Weed Control Conf. Residue. Rev.* 29:56-57.
- Cano, E. A., M. C. Bolarin, F. Perez-Alfocea and M. Caro (1991). Effect of NaCl priming on increased salt tolerance in tomato. *J. Hort. Sci.* 66:621-628.
- Cavaleiro, C. M. F., O. Roque, L. Proencada and A. Cunha (1993). Contribution for the characterization of Portuguese fennel chemotypes. *J. Essential Oil Res.* 5:223-225.
- Cayuela, E., F. Perez-Alfocea, M. Caro, and M. C. Bolarin (1996). Priming of seeds with NaCl induces physiological changes in tomato plants grown under salt stress. *Physiol. Plant.* 96:231-236.
- Carvajal, M., del Amor, F. M., Fernandez-Ballester, G., Martinez, V. and A. Cerda (1998) Time course of solute accumulation and water relations in muskmelon plants exposed to salt during different growth stages. *Plant Sci.* 138:103-112.
- Dahal, P., K. J. Bradford and R. A. Jones (1990). Effects of priming and endosperm integrity on seed germination rates of tomato genotypes germination at suboptimal temperatures. *J. Exp. Bot.* 41:1431-1439.
- Farooq, M., S. M. A Basra, K Hafez and N. Ahmad (2005). Thermal hardening: A new seed vigor enhancement tool in rice. *Acta Botanica Sinica.* 47:187-193.
- Franco, J. A., C. Esteban and C. Rodriguez (1993). Effects of salinity on various growth stages of muskmelon cv. Revigal. *J. Hort. Sci.* 68:899-904.
- Franco, J. A., J. A. Fernandez, S. Banon and A. Gonzalez (1997). Relationship between the effects of salinity on seedling leaf area and fruit yield of six muskmelon cultivars. *Hort. Sci.* 34:642-644.
- Ghiyasi, M., A. S. Abbasi, M. Tajbakhsh, R. Amirnia and H. Salehzade (2008). Effect of osmopriming with poly ethylene glycol 8000 (PEG8000) on germination and seedling growth of wheat (*Triticum aestivum* L.) seeds under salt stress. *Res. J. Biol. Sci.* 3(10):1249-1251.
- Harris, D., A. Joshi, P. A. Khan, P. Gothkar and P. S. Sodhi

- (1999). On-farm seed priming in semi-arid agriculture: development and evaluation in maize, rice and chickpea in India using participatory methods. *Exp. Agric.* 35:15-29.
- Harris, D., B. S. Raghuvanshi, J. S. Gangwar, S. C. Singh, K. D. Joshi, A. Rashid and P. A. Hollington (2001). Participatory evaluation by farmers of 'on-farm' seed priming in wheat in India, Nepal and Pakistan. *Exp. Agric.* 37:403-415.
- Harris, D., A. Rashid, P. A. Hollington, L. Jasi, and C. Riches (2002). Prospects of improving maize yields with 'on-farm' seed priming, p. 180-185. In: N. P. Rajbhandari, J. K. Ransom, K. Adikhari, R. A. F. E. Palme (Eds.). *Sustainable Maize Production Systems for Nepal: Proceedings of a Maize Symposium held, Kathmandu, Nepal, December 3-5, 2001.* Narc and Cimmyt.
- Harris, D., A. Rashid, S. Ali, and P. A. Hollington (2004). 'On-farm' seed priming with maize in Pakistan, p. 316-324. In: G. Srinivasan, P. H. Zaidi, B. M. Prasanna, F. Gonzalez, K. Lesnick (Eds.). *Proceedings of the 8th Asian Regional Maize Workshop: New Technologies for the New Millennium held Bangkok, Thailand, Cimmyt, Mexico, D.F., August 5-8, 2002.*
- Khazaie, H. R., F. Nadjafi and M. Bannayan (2008). Effect of irrigation frequency and planting density on herbage biomass and oil production of Thyme (*Thymus vulgaris*) and Hyssop (*Hyssopus officinalis*). *Journal of Industrial Crops and Products.* 27:315-321.
- Kumar, A., J. S. Gangwar, S. C. Prasad and D. Harris (2002). 'On-farm' seed priming increases yield of direct-sown finger millet (*Eleusine coracana*) in India. *Int. Sorghum Millets Newslett.* 43:90-92.
- Levit, J. (1980). *Response of plants to environmental stresses, Vol. II.* Academic press, New York.
- Marotti, M., V. Dellacecca, R. Piccaglia, E. Giovanelli, D. Palevitch and J. E. Simon (1993). Agronomic and chemical evaluation of three varieties of *Foeniculum vulgare* Mill. *Acta Hort.* 331:63-69.
- McDonald, M. B. (1999). Seed deterioration: physiology, repair and assessment. *Seed Sci. Technol.* 27:177-237.
- Michel, B. and E. Kaufmann (1973). The osmotic potential of polyethylene glycol 6000. *Plant Physiol.* 51:914-916.
- Miraj, G. (2005). Effect of phosphorus and zinc priming on germination, seedling growth and yield of maize. M.Sc. (Hons.) Thesis. Dept. Agric. Chem., NWFP Agricultural University, Peshawar, Pakistan.
- Money, N. P. (1989). Osmotic pressure of aqueous polyethylene glycols, relationship between molecular weight and vapor pressure deficit. *Plant physiology.* 91:766-769.
- Musa, A. M., D. Harris, C. Johansen and J. Kumar (2001). Short duration chickpea to replace fallow after aman rice: the role of on-farm seed priming in the High Barind Tract of Bangladesh. *Exp. Agric.* 37:509-521.
- Nadjafi, F., M. Bannayan, L. Tabrizi and M. Rastgoo (2006). Seed germination and dormancy breaking techniques for *Ferula* gummosa and *Teucrium polium*. *J. Arid Environments.* 64:542-547.
- Parera, C. A. and D. J. Cantliffe (1994). Presowing seed priming. *Hort. Rev.* 16: 109-141.
- Passam, H. C. and D. Kakouriotis (1994). The effects of osmoconditioning on the germination, emergence and early plant growth of cucumber under saline conditions. *Sci. Hort.* 57:233-240.
- Piccaglia, R. and M. Marrotti (1993). Characterization of several aromatic plants grown in northern Italy. *Flavor Fragrance Journal.* 8:112-115.
- Pill, W. G., J. J. Frett and D. C. Morneau (1991). Germination and seedling emergence of primed tomato and asparagus seeds under adverse conditions. *HortSci.* 26:1160-1162.
- Rashid, A., D. Harris, P. A. Hollington and R. A. Khattak (2002). On-farm seed priming: a key technology for improving the livelihoods of resource-poor farmers on saline lands, p. 423-431. In: R. Ahmad, K. A. Malik (Eds.). *Prospects for Saline Agriculture.* Kluwer Academic Publishers, The Netherlands.
- Rashid, A., D. Harris, P. A. Hollington, and M. Rafiq (2004). Improving the yield of mungbean (*Vigna radiata*) in the North West Frontier Province of Pakistan using on-farm seed priming. *Exp. Agric.* 40:233-244.
- Schintz, N., J. H. Xia and A. R. Kermod (2001). Dormancy of yellow cedar seeds is terminated by gibberellic acid in combination with fluridone or with osmotic priming and moist chilling. *Seed Sci. Technol.* 29:331-346.
- Shivankar, R. S., D. B. Deore and N. G. Zode (2003). Effect of pre-sowing seed treatment on establishment and seed yield of sunflower. *J. Oilseeds Res.* 20:299-300.
- Singh, B. G. (1995). Effect of hydration-dehydration seed treatments on vigour and yield of sunflower. *Indian. J. Plant physiol.* 38:66-68.
- Sivritepe, H. O. and A. M. Dourado (1995). The effect of priming treatments on the viability and accumulation of chromosomal damage in aged pea seeds. *Ann. Bot.* 75:165-171.
- Sosa L., A. Llanes, H. Reinoso, M. Reginato and V. Luna (2005). Osmotic and specific ion effect on the germination of *Prosopis strombulifera*. *Ann. Bot.* 96:261-267.
- Sung, J. M. and K. Y. Chiu (1995). Hydration effects on seedling emergence strength of watermelon seed differing in ploidy. *Plant Sci.* 110:21-26.
- Towned, J., P. W. Mtakwa, C. E. Mullins and L. P. Simmonds (1996). Soil physical factors limiting establishment of sorghum and cowpea in two contrasting soil types in the semi-arid tropics. *Soil Till. Res.* 40:89-106.
- Weaich, K., K. L. Bristow and A. Cass (1992). Preemergent shoot growth of maize under different drying conditions. *Soil Sci. Society Am. J.* 56:1272-1278.
- Wiebe, H. J. and T. Muhyaddin (1987). Improvement of emergence by osmotic seed treatments in soil of high salinity. *Acta Hort.* 198:91-100.