

## 2-Aminopyrimidinium hydrogen oxalate monohydrate

Hossein Eshtiagh-Hosseini,\* Zakieh Yousefi and Masoud Mirzaei

Department of Chemistry, School of Sciences, Ferdowsi University of Mashhad, Mashhad 917791436, Iran

Correspondence e-mail: heshtiagh@ferdowsi.um.ac.ir


Received 3 September 2009; accepted 30 September 2009

Key indicators: single-crystal X-ray study;  $T = 120$  K; mean  $\sigma(\text{C}-\text{C}) = 0.004$  Å;  $R$  factor = 0.055;  $wR$  factor = 0.148; data-to-parameter ratio = 12.2.

In the title hydrated salt,  $\text{C}_4\text{H}_6\text{N}_3^+\cdot\text{C}_2\text{HO}_4^-\cdot\text{H}_2\text{O}$ , intermolecular  $\text{N}-\text{H}\cdots\text{O}$  and  $\text{O}-\text{H}\cdots\text{O}$  hydrogen bonding helps to stabilize the crystal structure.

### Related literature

For the biological properties of pyrimidines, see: Rabie *et al.* (2007). For the applications of aminopyrimidines, see: Rospenk & Koll (2007). For aminopyrimidine salts, see: Childs *et al.* (2007).



### Experimental

#### Crystal data

$\text{C}_4\text{H}_6\text{N}_3^+\cdot\text{C}_2\text{HO}_4^-\cdot\text{H}_2\text{O}$   
 $M_r = 203.16$   
 Triclinic,  $P\bar{1}$

$a = 6.295$  (2) Å  
 $b = 6.339$  (2) Å  
 $c = 11.111$  (4) Å

$\alpha = 75.045$  (6)°  
 $\beta = 84.302$  (6)°  
 $\gamma = 86.026$  (7)°  
 $V = 425.8$  (2) Å<sup>3</sup>  
 $Z = 2$

Mo  $K\alpha$  radiation  
 $\mu = 0.14$  mm<sup>-1</sup>  
 $T = 120$  K  
 $0.35 \times 0.15 \times 0.07$  mm

#### Data collection

Bruker SMART 1000 CCD area-detector diffractometer  
 Absorption correction: none  
 3983 measured reflections

1835 independent reflections  
 1177 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.030$

#### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.055$   
 $wR(F^2) = 0.148$   
 $S = 1.02$   
 1835 reflections  
 151 parameters

H atoms treated by a mixture of independent and constrained refinement  
 $\Delta\rho_{\text{max}} = 0.44$  e Å<sup>-3</sup>  
 $\Delta\rho_{\text{min}} = -0.29$  e Å<sup>-3</sup>

**Table 1**  
 Hydrogen-bond geometry (Å, °).

| $D-H\cdots A$                                 | $D-H$    | $H\cdots A$ | $D\cdots A$ | $D-H\cdots A$ |
|---|----------|-------------|-------------|---------------|
| $\text{N1}-\text{H1}\cdots\text{O2}$          | 0.93 (3) | 1.75 (3)    | 2.671 (3)   | 173 (3)       |
| $\text{N2}-\text{H2A}\cdots\text{O1}$         | 0.96 (3) | 1.87 (3)    | 2.827 (3)   | 170 (3)       |
| $\text{N2}-\text{H2B}\cdots\text{O2}^i$       | 0.91 (4) | 1.99 (4)    | 2.885 (3)   | 171 (3)       |
| $\text{O4}-\text{H4O}\cdots\text{O1W}^{ii}$   | 0.89 (3) | 1.69 (4)    | 2.584 (3)   | 176 (4)       |
| $\text{O1W}-\text{H1WA}\cdots\text{O3}^{iii}$ | 0.97 (5) | 1.91 (5)    | 2.827 (3)   | 158 (4)       |
| $\text{O1W}-\text{H1WB}\cdots\text{O1}$       | 0.82 (5) | 2.14 (4)    | 2.812 (3)   | 139 (4)       |
| $\text{O1W}-\text{H1WB}\cdots\text{O3}$       | 0.82 (5) | 2.31 (5)    | 3.002 (3)   | 144 (4)       |

Symmetry codes: (i)  $x, y + 1, z$ ; (ii)  $x, y - 1, z$ ; (iii)  $-x + 1, -y + 1, -z$ .

Data collection: *SMART* (Bruker, 1998); cell refinement: *SAINT-Plus* (Bruker, 1998); data reduction: *SAINT-Plus*; program(s) used to solve structure: *SHELXTL* (Sheldrick, 2008); program(s) used to refine structure: *SHELXTL*; molecular graphics: *SHELXTL*; software used to prepare material for publication: *SHELXTL*

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: XU2606).

### References

- Bruker (1998). *SMART* and *SAINT-Plus*. Bruker AXS Inc., Madison, Wisconsin, USA.  
 Childs, S. L., Stahly, G. P. & Park, A. (2007). *Mol. Pharm.* **4**, 323–338.  
 Rabie, U. M., Abou-El-Wafa, M. H. & Mohamed, R. A. (2007). *J. Mol. Struct.* **871**, 6–13.  
 Rospenk, M. & Koll, A. (2007). *J. Mol. Struct.* **844–845**, 232–241.  
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.

**supplementary materials**

*Acta Cryst.* (2009). E65, o2816 [ doi:10.1107/S1600536809039907 ]

## 2-Aminopyrimidinium hydrogen oxalate monohydrate

H. Eshtiagh-Hosseini, Z. Yousefi and M. Mirzaei

### Comment

Pyrimidines have been attracted special attention because of unique biological properties, such as fungicides, vermicides, insecticides and medicines (Rabie *et al.*, 2007). Among them, aminopyrimidines are as interesting matters for chemists and pharmacist. They are a part of nucleic bases, cytosine, adenine and guanine which are responsible for molecular recognition and replication of DNA, through the formation and breakage of N—H···N hydrogen bonds (Rospenk & Koll, 2007). 2-Aminopyrimidine (2-apym), with amino group as two H-bond donor atoms and two N atoms as two H-bond acceptors atoms is particularly attractive as a very simple self-complementary prototype for chain formation with other organic molecules. Until now, a lot of co-crystals and proton transfer compounds were synthesized using 2-apym and a variety of carboxylic acid derivatives (Childs *et al.*, 2007). In this report, we choose oxalic acid (OxH<sub>2</sub>) in according to their difference in *pK<sub>a</sub>*.

In the title compound, the oxalic acid is mono-deprotonated while 2-apym is protonated (Fig. 1). The cation is hydrogen bonded to the anion with a cyclic  $R_2^2(8)$  pattern (Table 1). The anionic and cationic moieties link with the water molecule into layers by hydrogen bondings, resulting in beautiful picture as Flag-like structure. The O1W—H1WA···O3<sup>iii</sup> hydrogen bond [symmetry code: (iii) 1 - x, 1 - y, -z] between water and carboxyl group produces the three dimensional supra-molecular structure.

### Experimental

The title compound was synthesized *via* the reaction of OxH<sub>2</sub> (0.047 g, 0.375 mmol) with 2-apym (0.050 g, 0.5 mmol) in a water solution (5 ml). The solution was stirred for 3 h in 323 K. Colorless crystals were obtained after a week.

### Refinement

Nitrogen- and oxygen-bound H atoms were located in a difference Fourier map and refined isotropically. Carbon-bound H atoms were placed in calculated positions and were refined in riding model with  $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ .

### Figures




Fig. 1. The molecular structure of the title compound, showing the atom-numbering scheme. Displacement ellipsoids are drawn at the 50% probability level. Dashed lines indicate hydrogen bonding.

## 2-Aminopyrimidinium hydrogen oxalate monohydrate

### Crystal data

|  |   |
|--|---|
| $C_4H_6N_3^+ \cdot C_2HO_4^- \cdot H_2O$ | $Z = 2$   |
| $M_r = 203.16$                           | $F_{000} = 212$   |
| Triclinic, $P\bar{1}$                    | $D_x = 1.585 \text{ Mg m}^{-3}$                         |
| Hall symbol: -P 1                        | Melting point: 300 K                                    |
| $a = 6.295 (2) \text{ \AA}$              | Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$ |
| $b = 6.339 (2) \text{ \AA}$              | Cell parameters from 851 reflections                    |
| $c = 11.111 (4) \text{ \AA}$             | $\theta = 3.3\text{--}27.7^\circ$                       |
| $\alpha = 75.045 (6)^\circ$              | $\mu = 0.14 \text{ mm}^{-1}$                            |
| $\beta = 84.302 (6)^\circ$               | $T = 120 \text{ K}$                                     |
| $\gamma = 86.026 (7)^\circ$              | Prism, colorless  |
| $V = 425.8 (2) \text{ \AA}^3$            | $0.35 \times 0.15 \times 0.07 \text{ mm}$               |

### Data collection

|  |  |
|--|--|
| Bruker SMART 1000 CCD area-detector diffractometer | 1177 reflections with $I > 2\sigma(I)$ |
| Radiation source: normal-focus sealed tube         | $R_{\text{int}} = 0.030$               |
| Monochromator: graphite                            | $\theta_{\text{max}} = 27.0^\circ$     |
| $T = 120 \text{ K}$                                | $\theta_{\text{min}} = 1.9^\circ$      |
| $\varphi$ and $\omega$ scans                       | $h = -8 \rightarrow 8$                 |
| Absorption correction: none                        | $k = -8 \rightarrow 8$                 |
| 3983 measured reflections                          | $l = -14 \rightarrow 14$               |
| 1835 independent reflections                       |  |

### Refinement

|  |  |
|--|--|
| Refinement on $F^2$  | Secondary atom site location: difference Fourier map                   |
| Least-squares matrix: full                                     | Hydrogen site location: mixed  |
| $R[F^2 > 2\sigma(F^2)] = 0.055$                                | H atoms treated by a mixture of independent and constrained refinement |
| $wR(F^2) = 0.148$  | $w = 1/[\sigma^2(F_o^2) + (0.047P)^2 + 0.72P]$                         |
| $S = 1.02$   | where $P = (F_o^2 + 2F_c^2)/3$   |
| 1835 reflections   | $(\Delta/\sigma)_{\text{max}} < 0.001$                                 |
| 151 parameters   | $\Delta\rho_{\text{max}} = 0.44 \text{ e \AA}^{-3}$                    |
| Primary atom site location: structure-invariant direct methods | $\Delta\rho_{\text{min}} = -0.29 \text{ e \AA}^{-3}$                   |
|  | Extinction correction: none  |

### Special details

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations

between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

|      | $x$        | $y$         | $z$          | $U_{\text{iso}}^*/U_{\text{eq}}$ |
|------|------------|-------------|--------------|----------------------------------|
| N1   | 0.2235 (4) | 0.3419 (4)  | 0.6301 (2)   | 0.0225 (5)                       |
| H1   | 0.235 (5)  | 0.290 (5)   | 0.559 (3)    | 0.037 (9)*                       |
| N2   | 0.2566 (4) | 0.6890 (4)  | 0.5036 (2)   | 0.0263 (6)                       |
| H2A  | 0.270 (5)  | 0.637 (5)   | 0.429 (3)    | 0.027 (8)*                       |
| H2B  | 0.258 (5)  | 0.836 (6)   | 0.492 (3)    | 0.040 (9)*                       |
| N3   | 0.2128 (4) | 0.6436 (4)  | 0.7189 (2)   | 0.0249 (5)                       |
| C2   | 0.2305 (4) | 0.5594 (4)  | 0.6174 (2)   | 0.0232 (6)                       |
| C4   | 0.1869 (4) | 0.5030 (5)  | 0.8289 (3)   | 0.0264 (6)                       |
| H4A  | 0.1759     | 0.5588      | 0.9010       | 0.032*                           |
| C5   | 0.1746 (5) | 0.2770 (5)  | 0.8472 (3)   | 0.0270 (6)                       |
| H5A  | 0.1530     | 0.1818      | 0.9284       | 0.032*                           |
| C6   | 0.1952 (4) | 0.2013 (5)  | 0.7428 (3)   | 0.0262 (6)                       |
| H6A  | 0.1895     | 0.0492      | 0.7495       | 0.031*                           |
| O1   | 0.2756 (3) | 0.4892 (3)  | 0.30262 (17) | 0.0276 (5)                       |
| O2   | 0.2664 (3) | 0.1595 (3)  | 0.43700 (17) | 0.0268 (5)                       |
| O3   | 0.3456 (3) | 0.2958 (3)  | 0.11122 (17) | 0.0316 (5)                       |
| O4   | 0.3088 (4) | -0.0288 (3) | 0.24753 (19) | 0.0323 (5)                       |
| H4O  | 0.314 (5)  | -0.099 (6)  | 0.187 (3)    | 0.039 (9)*                       |
| C7   | 0.2830 (4) | 0.2859 (4)  | 0.3300 (2)   | 0.0229 (6)                       |
| C8   | 0.3161 (4) | 0.1829 (4)  | 0.2173 (2)   | 0.0230 (6)                       |
| O1W  | 0.3244 (4) | 0.7844 (3)  | 0.0656 (2)   | 0.0298 (5)                       |
| H1WA | 0.460 (8)  | 0.772 (7)   | 0.018 (4)    | 0.078 (15)*                      |
| H1WB | 0.306 (7)  | 0.661 (8)   | 0.109 (4)    | 0.070 (14)*                      |

*Atomic displacement parameters ( $\text{\AA}^2$ )*

|    | $U^{11}$    | $U^{22}$    | $U^{33}$    | $U^{12}$     | $U^{13}$     | $U^{23}$     |
|----|-------------|-------------|-------------|--------------|--------------|--------------|
| N1 | 0.0308 (13) | 0.0209 (12) | 0.0172 (12) | -0.0009 (9)  | -0.0015 (9)  | -0.0078 (10) |
| N2 | 0.0432 (15) | 0.0175 (12) | 0.0176 (12) | -0.0027 (10) | -0.0019 (10) | -0.0035 (10) |
| N3 | 0.0325 (13) | 0.0240 (12) | 0.0193 (12) | -0.0007 (10) | -0.0031 (9)  | -0.0071 (10) |
| C2 | 0.0284 (15) | 0.0221 (14) | 0.0205 (13) | -0.0005 (11) | -0.0036 (11) | -0.0078 (11) |
| C4 | 0.0299 (15) | 0.0321 (16) | 0.0192 (14) | -0.0013 (12) | -0.0026 (11) | -0.0100 (12) |
| C5 | 0.0320 (16) | 0.0273 (15) | 0.0191 (14) | -0.0006 (12) | -0.0038 (11) | -0.0008 (11) |
| C6 | 0.0316 (16) | 0.0211 (13) | 0.0244 (14) | -0.0023 (11) | -0.0026 (11) | -0.0028 (11) |
| O1 | 0.0423 (12) | 0.0178 (10) | 0.0224 (10) | -0.0008 (8)  | -0.0025 (8)  | -0.0050 (8)  |
| O2 | 0.0458 (12) | 0.0174 (9)  | 0.0166 (10) | -0.0010 (8)  | -0.0031 (8)  | -0.0029 (8)  |
| O3 | 0.0544 (14) | 0.0221 (10) | 0.0175 (10) | -0.0041 (9)  | -0.0013 (9)  | -0.0038 (8)  |

## supplementary materials

|     |             |             |             |              |              |              |
|-----|-------------|-------------|-------------|--------------|--------------|--------------|
| O4  | 0.0605 (15) | 0.0183 (10) | 0.0196 (10) | -0.0015 (9)  | -0.0015 (9)  | -0.0080 (8)  |
| C7  | 0.0268 (15) | 0.0228 (14) | 0.0195 (13) | 0.0003 (11)  | -0.0016 (11) | -0.0068 (11) |
| C8  | 0.0296 (15) | 0.0195 (13) | 0.0208 (14) | -0.0010 (11) | -0.0022 (11) | -0.0065 (11) |
| O1W | 0.0455 (14) | 0.0198 (11) | 0.0240 (11) | -0.0019 (9)  | -0.0029 (9)  | -0.0054 (9)  |

### Geometric parameters ( $\text{\AA}$ , $^\circ$ )

|             |            |               |            |
|-------------|------------|---------------|------------|
| N1—C6       | 1.340 (3)  | C5—H5A        | 0.9500     |
| N1—C2       | 1.353 (3)  | C6—H6A        | 0.9500     |
| N1—H1       | 0.93 (4)   | O1—C7         | 1.244 (3)  |
| N2—C2       | 1.320 (3)  | O2—C7         | 1.250 (3)  |
| N2—H2A      | 0.96 (3)   | O3—C8         | 1.215 (3)  |
| N2—H2B      | 0.91 (4)   | O4—C8         | 1.299 (3)  |
| N3—C4       | 1.316 (4)  | O4—H4O        | 0.90 (4)   |
| N3—C2       | 1.359 (3)  | C7—C8         | 1.545 (4)  |
| C4—C5       | 1.400 (4)  | O1W—H1WA      | 0.97 (5)   |
| C4—H4A      | 0.9500     | O1W—H1WB      | 0.82 (5)   |
| C5—C6       | 1.358 (4)  |               |            |
| C6—N1—C2    | 121.5 (2)  | C6—C5—H5A     | 121.8      |
| C6—N1—H1    | 119 (2)    | C4—C5—H5A     | 121.8      |
| C2—N1—H1    | 119 (2)    | N1—C6—C5      | 119.7 (3)  |
| C2—N2—H2A   | 123.4 (18) | N1—C6—H6A     | 120.1      |
| C2—N2—H2B   | 120 (2)    | C5—C6—H6A     | 120.1      |
| H2A—N2—H2B  | 116 (3)    | C8—O4—H4O     | 119 (2)    |
| C4—N3—C2    | 116.5 (2)  | O1—C7—O2      | 127.2 (3)  |
| N2—C2—N1    | 118.4 (2)  | O1—C7—C8      | 115.1 (2)  |
| N2—C2—N3    | 120.5 (2)  | O2—C7—C8      | 117.7 (2)  |
| N1—C2—N3    | 121.2 (2)  | O3—C8—O4      | 124.8 (3)  |
| N3—C4—C5    | 124.7 (3)  | O3—C8—C7      | 121.1 (2)  |
| N3—C4—H4A   | 117.7      | O4—C8—C7      | 114.1 (2)  |
| C5—C4—H4A   | 117.7      | H1WA—O1W—H1WB | 104 (4)    |
| C6—C5—C4    | 116.4 (3)  |               |            |
| C6—N1—C2—N2 | -179.3 (3) | C2—N1—C6—C5   | -0.5 (4)   |
| C6—N1—C2—N3 | 1.2 (4)    | C4—C5—C6—N1   | -0.6 (4)   |
| C4—N3—C2—N2 | 179.9 (3)  | O1—C7—C8—O3   | 4.3 (4)    |
| C4—N3—C2—N1 | -0.6 (4)   | O2—C7—C8—O3   | -175.8 (3) |
| C2—N3—C4—C5 | -0.7 (4)   | O1—C7—C8—O4   | -175.5 (2) |
| N3—C4—C5—C6 | 1.3 (4)    | O2—C7—C8—O4   | 4.4 (4)    |

### Hydrogen-bond geometry ( $\text{\AA}$ , $^\circ$ )

| $D-H\cdots A$                       | $D-H$    | $H\cdots A$ | $D\cdots A$ | $D-H\cdots A$ |
|-------------------------------------|----------|-------------|-------------|---------------|
| N1—H1 $\cdots$ O2                   | 0.93 (3) | 1.75 (3)    | 2.671 (3)   | 173 (3)       |
| N2—H2A $\cdots$ O1                  | 0.96 (3) | 1.87 (3)    | 2.827 (3)   | 170 (3)       |
| N2—H2B $\cdots$ O2 <sup>i</sup>     | 0.91 (4) | 1.99 (4)    | 2.885 (3)   | 171 (3)       |
| O4—H4O $\cdots$ O1W <sup>ii</sup>   | 0.89 (3) | 1.69 (4)    | 2.584 (3)   | 176 (4)       |
| O1W—H1WA $\cdots$ O3 <sup>iii</sup> | 0.97 (5) | 1.91 (5)    | 2.827 (3)   | 158 (4)       |
| O1W—H1WB $\cdots$ O1                | 0.82 (5) | 2.14 (4)    | 2.812 (3)   | 139 (4)       |

O1W—H1WB...O3

0.82 (5)

2.31 (5)

3.002 (3)

144 (4)

Symmetry codes: (i)  $x, y+1, z$ ; (ii)  $x, y-1, z$ ; (iii)  $-x+1, -y+1, -z$ .

Fig. 1

