



*7th Seminar on Probability and
Stochastic Processes
Isfahan University of Technology
Aug. 13-14, 2009*



Almost Sure Convergence of Kernel Bivariate distribution Function Estimator under Negative Association

H. Jabbari

Department of Statistics, faculty of Mathematical Sciences, Ferdowsi University of Mashhad, Mashhad, Iran
email: Jabbarinh@um.ac.ir

Abstract

Let $\{X_n, n \geq 1\}$ be a strictly stationary sequence of negatively associated random variables, with common distribution function F . In this paper, we consider the estimation of the two-dimensional distribution function of (X_1, X_{k+1}) for fixed $k \in \mathbb{N}$ based on kernel type estimators. We introduce asymptotic normality and properties and moments. From these we derive the optimal bandwidth convergence rate, which is of order n^{-1} . Besides of some usual conditions on the kernel function, the conditions typically impose a convenient increase rate on the covariances $Cov(X_1, X_n)$.

MSC 2000: Primary 62G20; Secondary 60F15, 62N05.

keywords: Almost sure convergence, Bivariate distribution function, Kernel estimation, Negatively association, Stationarity.

1 Introduction, definitions and assumptions

The interest on approximating distribution functions of random pairs arises from the characterizations of the limiting distribution of empirical processes, which has been a subject of interest for many statisticians. The first results concerning the asymptotic distribution of the sequence data back to Donsker [3], for independent underlying variables $\{X_n, n \geq 1\}$. The extension of this characterization to nonindependent variables was eventually studied. One of the dependence structures is positive association. Azevedo and Oliveira [2] studied kernel type estimation of bivariate distribution function for positively associated random variables. The other type of dependence is negative association (NA), introduced by Alam and Saxena [1] and carefully studied by Joag-Dev and Proschan [6]. A finite family of random variables $\{X_i, 1 \leq i \leq n\}$ is said to be negatively associated if for every pair of disjoint subsets A and B of $\{1, 2, \dots, n\}$,

$$Cov(f_1(X_i, i \in A), f_2(X_j, j \in B)) \leq 0$$

whenever f_1 and f_2 are coordinatewise increasing and such that the covariance exists. An infinite family of random variables is *NA* if every finite subfamily is *NA*. Because of their wide applications in multivariate statistical analysis and reliability theory, the notion of *NA* has received more and more attention recently. We refer to Joag-Dev and Proschan [6] for fundamental properties, Newman [9] and Su and Chi [14] for central limit theorem, Matula [8] for three series theorem, Su et al. [15] for a moment inequality, a weak invariance principle and example to show that there exists infinite family of non-degenerate non-independent strictly stationary *NA* random variables, Shao [13] for the Rosenthal type maximal inequality and Kolmogorov exponential inequality, Liang and Su [7] for convergence rates of law of the logarithm, Roussas [11] for the central limit theorem of random fields, some examples and applications and Yuan et al. [16] for improving the result of Roussas [11]. The above comments motivated the interest on

the estimation of the bivariate distribution function under negative association. A natural estimator of $F(x, y) = P(X_1 \leq x, X_{k+1} \leq y)$ with k fixed, is defined by

$$\hat{\varphi}_n(x, y) = \frac{1}{n-k} \sum_{i=1}^{n-k} (1_{(-\infty, x]}(X_i) 1_{(-\infty, y]}(X_{k+i})). \tag{1}$$

The asymptotic behavior of this estimator were studied by Jabbari et al. [5], who considered conditions on the covariance structure of the sequence $\{X_n, n \geq 1\}$, for the almost sure consistency of this estimator. Here we will considered the kernel estimator of F , defined by

$$\begin{aligned} \hat{F}_n(x, y) &= \frac{1}{n-k} \sum_{i=1}^{n-k} U\left(\frac{x-X_i}{h_n}, \frac{y-X_{k+i}}{h_n}\right) \\ &= \int_{\mathbb{R}^2} U\left(\frac{x-s}{h_n}, \frac{y-t}{h_n}\right) d\hat{\varphi}_n(s, t), \end{aligned} \tag{2}$$

where U is a given distribution function and $\{h_n, n \geq 1\}$, is a sequence of positive numbers converging to zero. The assumptions under which the main result in this paper is obtained are gathered as below.

(A1). $\{X_n, n \geq 1\}$, is an NA and strictly stationary sequence of random variables having density function f bounded by M_0 ; let $M_1 = 2 \max(2/\pi^2, 45M_0)$.

(A2). k is a fixed integer and F the distribution function of (X_1, X_{k+1}) . F has bounded and continuous partial derivatives of first and second orders.

(A3). For each positive integer j , F_j is the distribution function of $(X_1, X_{k+1}, X_j, X_{k+j})$. F_j has bounded and continuous partial derivatives of first and second orders.

(A4). U is twice differentiable. If $u = \frac{\partial^2 U}{\partial x \partial y}$, is satisfies

$$\begin{aligned} \int_{\mathbb{R}^2} xu(x, y) dx dy &= \int_{\mathbb{R}^2} yu(x, y) dx dy = 0, \\ \int_{\mathbb{R}^2} x^2u(x, y) dx dy, \int_{\mathbb{R}^2} y^2u(x, y) dx dy &< \infty. \end{aligned}$$

(A5). The sequence of bandwidth is such that $nh_n^2 \rightarrow 0$.

(A6). $\sum_{j=3}^{\infty} j |Cov^{1/3}(X_1, X_{j-k})| < \infty$.

(A7). $V = \frac{\partial^2 U^2}{\partial x \partial y}$ is such that

$$\int_{\mathbb{R}^2} x^2V(x, y) dx dy, \int_{\mathbb{R}^2} y^2V(x, y) dx dy < \infty.$$

In Sections 2 and 3, we study the convergence and mean square error of \hat{F}_n . In Section 4, we consider the asymptotic distribution of the finite dimensional distributions of \hat{F}_n .

2 Consistency of the estimator

We first characterize the convergence rate of $E[\hat{F}_n(x, y)]$ and show that \hat{F}_n is asymptotically unbiased. To achieve this goal we apply a strong law of large numbers to the random variables $U\left(\frac{x-X_i}{h_n}, \frac{y-X_{k+i}}{h_n}\right)$; $i = 1, 2, \dots, n-k$ and we need to study the asymptotic properties of the covariance between each term in the sum (2).

Theorem 1. *Suppose that $\{X_n, n \geq 1\}$ satisfies (A1), (A2) and (A4). Then, every $x, y \in \mathbb{R}$,*

$$\begin{aligned} E[\hat{F}_n(x, y)] &= F(x, y) + \frac{h_n^2}{2} \left[\frac{\partial^2 F}{\partial x^2}(x, y) \int s^2 u(s, t) ds dt \right. \\ &\quad \left. + \frac{\partial^2 F}{\partial x \partial y}(x, y) \int stu(s, t) ds dt + \frac{\partial^2 F}{\partial y^2}(x, y) \int t^2 u(s, t) ds dt \right] + o(h_n^2). \end{aligned}$$

Proof. As $E[\hat{\varphi}_n(x, y)] = F(x, y)$ it follows from (2) that

$$\begin{aligned} E[\hat{F}_n(x, y)] &= \int_{\mathbb{R}^2} U\left(\frac{x-s}{h_n}, \frac{y-t}{h_n}\right) dF(s, t) \\ &= \int_{\mathbb{R}^2} u(w, v) F(x-wh_n, y-vh_n) dw dv. \end{aligned}$$

Using a Taylor expansion of order 2 of F and taking account of (A2) and (A4) and of the continuity of the second order partial derivatives of F the theorem follows. \square

From Theorem 1 and an application of the Dominated Convergence Theorem, we conclude

$$E[\hat{F}_n(x, y)] \longrightarrow F(x, y).$$

Now, we establish the almost sure convergence of \hat{F}_n . For this goal, we need some lemmas that are proved below.

Lemma 1. *Suppose that $\{X_n, n \geq 1\}$ be negatively associated. Let*

$$H_{A,B} = P_r(X_j > x_j, j \in A \cup B) - P_r(X_l > x_l, l \in A) \cdot P_r(X_m > x_m, m \in B),$$

where x_j 's are real and A and B are disjoint subsets of $\{1, \dots, n\}$. Define $H_{l,m} = H_{\{l\}, \{m\}}$. Then

$$0 \geq H_{A,B} \geq \sum_{l \in A} \sum_{m \in B} H_{l,m}. \tag{3}$$

Proof. For the proof see Theorem 2 in Newman [10]. \square

Lemma 2. *Suppose $\{X_n, n \geq 1\}$ satisfies (A1) and (A3). Then, for each $j > 1$ and $x, y, r, s \in \mathbb{R}$*

$$\begin{aligned} |F_j(x, y, r, s) - F(x, y) \cdot F(r, s)| &\leq 2M_1 |Cov^{1/3}(X_1, X_j) + Cov^{1/3}(X_1, X_{k+j}) \\ &\quad + Cov^{1/3}(X_{k+1}, X_j) + Cov^{1/3}(X_{k+1}, X_{k+j})|. \end{aligned} \tag{4}$$

Proof. The density of the variables is bounded by M_0 . Then, by Corollary of Theorem 1 in Sadikova [12] and relation (21) in Newman [9],

$$Cov(1_{(x, +\infty)}(X_1), 1_{(y, +\infty)}(X_{k+1})) \geq M_1 Cov^{1/3}(X_1, X_{k+1}), \quad x, y \in \mathbb{R}. \tag{5}$$

Since $\{X_n, n \geq 1\}$ is NA then for $j > 1$, by (5), NA properties and Lemma 1 we conclude

$$\begin{aligned} 0 &\geq F_j(x, y, r, s) - F(x, y) \cdot F(r, s) \\ &\geq M_1 Cov^{1/3}(X_1, X_j) + M_1 Cov^{1/3}(X_1, X_{k+j}) + M_1 Cov^{1/3}(X_{k+1}, X_j) + M_1 Cov^{1/3}(X_{k+1}, X_{k+j}) + \\ &\quad P_r(X_1 > x, X_{k+1} > y, X_j > r, X_{k+j} > s) - P_r(X_1 > x, X_{k+1} > y) \cdot P_r(X_j > r, X_{k+j} > s) \\ &\geq 2M_1 [Cov^{1/3}(X_1, X_j) + Cov^{1/3}(X_1, X_{k+j}) + Cov^{1/3}(X_{k+1}, X_j) + Cov^{1/3}(X_{k+1}, X_{k+j})]. \end{aligned}$$

So, the proof is complete. \square

Lemma 3. *Suppose the variables $\{X_n, n \geq 1\}$ satisfies (A1), (A2), (A3) and (A4). Then, for each $j \geq 1$ and $x, y \in \mathbb{R}$*

$$Cov\left[U\left(\frac{x-X_1}{h_n}, \frac{y-X_{k+1}}{h_n}\right), U\left(\frac{x-X_j}{h_n}, \frac{y-X_{k+j}}{h_n}\right)\right] = F_j(x, y, x, y) - F^2(x, y) + O(h_n^2).$$

Proof. Rewrite the covariance as

$$\begin{aligned} &Cov\left[U\left(\frac{x-X_1}{h_n}, \frac{y-X_{k+1}}{h_n}\right), U\left(\frac{x-X_j}{h_n}, \frac{y-X_{k+j}}{h_n}\right)\right] \\ &= \int_{\mathbb{R}^4} U\left(\frac{x-r}{h_n}, \frac{y-s}{h_n}\right) U\left(\frac{x-u}{h_n}, \frac{y-v}{h_n}\right) dF_j(r, s, u, v) - \left(\int_{\mathbb{R}^2} U\left(\frac{x-r}{h_n}, \frac{y-s}{h_n}\right) dF(r, s)\right)^2, \end{aligned}$$

where the second term on the right hand side is just $E^2[\hat{F}_n(x, y)]$ and its behavior has been described in Theorem 1. For the first term, we write the function U as an integral and use Fubini's Theorem. So, by expanding F_j to the second order and using (A3) and (A4), this integral is equal to $F_j(x, y, x, y) + O(h_n^2)$. Therefore, the proof is complete by the limit behavior of $E[\hat{F}_n(x, y)]$. \square

Next theorem considers the almost sure convergence of \hat{F}_n .

Theorem 2. *Suppose $\{X_n, n \geq 1\}$ satisfy (A1), (A2), (A3), (A4), (A6) and (A7). Then, for each $x, y \in \mathbb{R}$*

$$\hat{F}_n(x, y) \longrightarrow F(x, y) \quad a.s.. \tag{6}$$

Proof. It is enough to prove that the variables $U(\frac{x-X_m}{h_n}, \frac{y-X_{k+m}}{h_n})$, $m > 1$ satisfy a strong law of large numbers. As U is coordinatewise nondecreasing, these variables are stationary and NA. According to Newman [9], U 's verify a strong law of large numbers if

$$\lim_{n \rightarrow \infty} \frac{1}{n-k} \sum_{j=2}^{n-k} |Cov[U(\frac{x-X_1}{h_n}, \frac{y-X_{k+1}}{h_n}), U(\frac{x-X_j}{h_n}, \frac{y-X_{k+j}}{h_n})]| = 0. \tag{7}$$

From Lemma 3 and using (4) for $r = x$ and $s = y$, it follows

$$\begin{aligned} |Cov[U(\frac{x-X_1}{h_n}, \frac{y-X_{k+1}}{h_n}), U(\frac{x-X_j}{h_n}, \frac{y-X_{k+j}}{h_n})]| &\leq |F_j(x, y, x, y) - F^2(x, y)| + O(h_n^2) \\ &\leq 2M_1 |Cov^{1/3}(X_1, X_j) + Cov^{1/3}(X_1, X_{k+j}) + \\ &\quad Cov^{1/3}(X_{k+1}, X_j) + Cov^{1/3}(X_{k+1}, X_{k+j})|. \end{aligned}$$

Since $\{X_n, n \geq 1\}$ is NA, by Lemma 8 in Newman [10], we have $\sum_{j=2}^{\infty} |Cov(X_1, X_j)| < \infty$. Then, we may conclude that $Cov(X_1, X_j)$ is nondecreasing as $j \rightarrow \infty$. So, the proof follows from (A6). \square

For the formulation of the next results we need to introduce some additional notations. Let t be a sequence of positive integers such that $t \rightarrow +\infty$. For each $i = 1, \dots, t$, put $x_i = Q(i/t)$, where Q is the quantile function of F . Define then,

$$D_n = \sup_{x, y \in \mathbb{R}} |\hat{F}_n(x, y) - F(x, y)|,$$

and

$$D_n^* = \max_{i, j=1, \dots, t} |\hat{F}_n(x_i, x_j) - F(x_i, x_j)|.$$

To prove an uniform version of the preceding theorem, we will apply the following result which is proved in Theorem 2 of Henriques and Oliveira [4].

Lemma 4. *If the sequence $\{X_n, n \geq 1\}$ satisfies (A1), then for each $n \in \mathbb{N}$,*

$$D_n \leq D_n^* + \frac{2}{t} \quad a.s. \quad \square \tag{8}$$

Next theorem is the uniform consistency of the estimator under the same set of conditions as in Theorem 2.

Theorem 3. *Suppose $\{X_n, n \geq 1\}$ satisfies (A1), (A2), (A3), (A4), (A6) and (A7). Then*

$$\sup_{x, y \in \mathbb{R}} |\hat{F}_n(x, y) - F(x, y)| \longrightarrow 0 \quad a.s.. \tag{9}$$

Proof. From Theorem 2, it follows that

$$D_n^* = \max_{1 \leq i, j \leq t} |\hat{F}_n(\frac{i}{t}, \frac{j}{t}) - F(\frac{i}{t}, \frac{j}{t})| \longrightarrow 0 \quad a.s..$$

Lemma 2.4 implies that for all $x, y \in \mathbb{R}$,

$$D_n = \sup_{x, y \in \mathbb{R}} |\hat{F}_n(x, y) - F(x, y)| \leq D_n^* + \frac{2}{t}.$$

So, the proof is complete, as t is arbitrary. \square

3 The behavior of the mean square error

In this section, we study the asymptotic properties and convergence rate of the mean square error of the estimator. From which, we derive the optimal bandwidth convergence rate of order n^{-1} . This rate is different from the one in the independent case. We write

$$MSE[\hat{F}_n(x, y)] = Var[\hat{F}_n(x, y)] + (E[\hat{F}_n(x, y)] - F(x, y))^2.$$

Theorem 1 gives the behavior of $E[\hat{F}_n(x, y)]$. Then, we need to describe the asymptotic properties and convergence rate of

$$Var[\hat{F}_n(x, y)] = \frac{1}{n-k} Var[U(\frac{x-X_1}{h_n}, \frac{y-X_{k+1}}{h_n})] + \frac{2}{(n-k)^2} \sum_{j=2}^{n-k} (n-k-j+1) Cov[U(\frac{x-X_1}{h_n}, \frac{y-X_{k+1}}{h_n}), U(\frac{x-X_j}{h_n}, \frac{y-X_{k+j}}{h_n})]. \tag{10}$$

Lemma 3 gives the asymptotic property of all these terms in (10). Just notice that the variance term, which corresponds to the choice $j = 1$ in Lemma 3, gives as limit $F_1(x, y, x, y) - F^2(x, y) = F(x, y) - F^2(x, y)$. Now, we state the result in the following theorem.

Theorem 4. *Suppose that $\{X_n, n \geq 1\}$ satisfies (A1), (A2), (A3), (A4), (A5), (A6) and (A7). Then, for all $x, y \in \mathbb{R}$*

$$(n-k)MSE[\hat{F}_n(x, y)] = F(x, y) - F^2(x, y) + 2 \sum_{j=2}^{\infty} (F_j(x, y, x, y) - F^2(x, y)) + O(h_n + nh_n^2) + a_n,$$

where

$$a_n = \frac{1}{(n-k)} \sum_{j=2}^{\infty} (j-1)(F_j(x, y, x, y) - F^2(x, y)) - 2 \sum_{j=n-k-1}^{\infty} (F_j(x, y, x, y) - F^2(x, y)).$$

Note that, from assumption (A6) and Lemma 2, $a_n \rightarrow 0$ as $n \rightarrow \infty$, and a_n is independent of the bandwidth choice. It is now evident that an optimization of the convergence rate of the MSE is achieved by choosing $h_n = Cn^{-1}$ for $C > 0$.

4 Finite dimensional distributions

Now, we study the asymptotic behavior of the finite dimensional distributions of the estimator. For this goal, we will use a decomposition of the sum (2) into several blocks that are negatively associated. This approximation is controlled via Newman's inequality [10]. As the proof is long and quite technical, we will divide it into some lemmas. Before proceeding define, for $x, y, r, s \in \mathbb{R}$,

$$\alpha_n(x, y) = \sqrt{n-k}(\hat{F}_n(x, y) - F(x, y)),$$

$$\sigma^2(x, y, r, s) = F(\min(x, r), \min(y, s)) - F(x, y).F(r, s) + 2 \sum_{j=2}^{\infty} (F_j(x, y, r, s) - F(x, y).F(r, s)).$$

For simplicity, throughout this section we assume that $\{X_n, n \geq 1\}$ satisfies (A1), (A2), (A3), (A4), (A5), (A6) and (A7).

Lemma 5. *For every $x, y, r, s \in \mathbb{R}$, we have*

$$Cov[\alpha_n(x, y), \alpha_n(r, s)] \rightarrow \sigma^2(x, y, r, s). \tag{11}$$

Proof. Using stationarity of the variables, we may write

$$\begin{aligned} \text{Cov}[\alpha_n(x, y), \alpha_n(r, s)] &= \text{Cov}\left[U\left(\frac{x-X_1}{h_n}, \frac{y-X_{k+1}}{h_n}\right), U\left(\frac{r-X_1}{h_n}, \frac{s-X_{k+1}}{h_n}\right)\right] \\ &+ \frac{2}{n-k} \sum_{j=2}^{n-k} (n-k-j+1) \text{Cov}\left[U\left(\frac{x-X_1}{h_n}, \frac{y-X_{k+1}}{h_n}\right), U\left(\frac{r-X_j}{h_n}, \frac{s-X_{k+j}}{h_n}\right)\right]. \end{aligned} \quad (12)$$

From Lemma 3 we have, for $j = 1, \dots, n-k$

$$\text{Cov}\left[U\left(\frac{x-X_1}{h_n}, \frac{y-X_{k+1}}{h_n}\right), U\left(\frac{r-X_j}{h_n}, \frac{s-X_{k+j}}{h_n}\right)\right] = F_j(x, y, r, s) - F(x, y) \cdot F(r, s) + O(h_n^2).$$

Inserting these characterizations in (12), we find that the last term in the right hand side of (12) is equal to

$$2 \sum_{j=2}^{n-k} (F_j(x, y, r, s) - F(x, y) \cdot F(r, s)) - \frac{2}{n-k} \sum_{j=2}^{n-k} (j-1) (F_j(x, y, r, s) - F(x, y) \cdot F(r, s)) + O(nh_n^2).$$

From Lemma 2 and using Assumption (A6), we have

$$\frac{1}{n-k} \sum_{j=2}^{n-k} (j-1) |F_j(x, y, r, s) - F(x, y) \cdot F(r, s)| \leq \frac{8M_1}{n-k} \sum_{j=2}^{n-k} j |Cov^{1/3}(X_1, X_{j-k})| \rightarrow 0.$$

□

Now, for the proof of the lemmas that concerned the asymptotic normality we need some further notation. Denote $\tilde{n} = n - k$ and given an integer $p \leq \tilde{n}$, let m be the largest integer less than or equal to \tilde{n}/p . Let $q \in \mathbb{N}$, $c_1, \dots, c_q \in \mathbb{R}$, $x, y, x_1, \dots, x_q, y_1, \dots, y_q \in \mathbb{R}$ and define

$$\begin{aligned} T_{\tilde{n},i}(x, y) &= U\left(\frac{x-X_i}{h_n}, \frac{y-X_{k+i}}{h_n}\right) - E\left[U\left(\frac{x-X_i}{h_n}, \frac{y-X_{k+i}}{h_n}\right)\right], \\ Y_j^p(x, y) &= \frac{1}{\sqrt{p}} \sum_{i=(j-1)p+1}^{jp} T_{\tilde{n},i}(x, y), \quad W_j^p = \sum_{k=1}^q c_k Y_j^p(x_k, y_k), \end{aligned}$$

and

$$Z_{mp} = \frac{1}{\sqrt{m}} \sum_{k=1}^q c_k \sum_{j=1}^m Y_j^p(x_k, y_k), \quad Z_{\tilde{n}} = \frac{1}{\sqrt{\tilde{n}}} \sum_{k=1}^q c_k \sum_{i=1}^{\tilde{n}} T_{\tilde{n},i}(x_k, y_k).$$

Note that, as follows from Lemma 5,

$$\text{Var}(Z_{\tilde{n}}) \rightarrow \sigma^2 := \sum_{k=1}^q c_k^2 \sigma^2(x_k, y_k, x_k, y_k) + 2 \sum_{k=1}^{q-1} \sum_{l=k+1}^q c_k c_l \sigma^2(x_k, y_k, x_l, y_l). \quad (13)$$

Further, for each p fixed, it follows from Lemma 3 that

$$\begin{aligned} \text{Var}[Y_1^p(x, y)] &= \text{Cov}\left[\frac{1}{\sqrt{p}} \sum_{i=1}^p T_{\tilde{n},i}(x, y), \frac{1}{\sqrt{p}} \sum_{j=1}^p T_{\tilde{n},j}(x, y)\right] \\ &= \frac{1}{p} \sum_{i=1}^p \sum_{j=1}^p (F_{|j-i+1|}(x, y, x, y) - F^2(x, y)) + O(ph_n^2). \end{aligned} \quad (14)$$

Then

$$\text{Var}(W_j^p) = \sigma_p^2 := \sum_{k=1}^q c_k^2 \sigma_p^2(x_k, y_k, x_k, y_k) + 2 \sum_{k=1}^{q-1} \sum_{l=k+1}^q c_k c_l \sigma_p^2(x_k, y_k, x_l, y_l), \quad (15)$$

where

$$\begin{aligned} \sigma_p^2(x_k, y_k, x_l, y_l) &= F(\min(x_k, x_l), \min(y_k, y_l)) - F(x_k, y_k) \cdot F(x_l, y_l) \\ &\quad + \frac{2p-2}{p} \sum_{j=2}^p (F_j(x_k, y_k, x_l, y_l) - F(x_k, y_k) \cdot F(x_l, y_l)). \end{aligned} \tag{16}$$

We now proceed to the lemmas that provided the asymptotic normality. First, replace the sum up to \tilde{n} by the sum up to mp to find a sum consisting only on the blocks W_j^p .

Lemma 6. *For fixed p , we have*

$$|Ee^{itZ_{\tilde{n}}} - Ee^{itZ_{mp}}| \longrightarrow 0. \tag{17}$$

Proof. Follow the arguments of the proof of Lemma 5.2 in Azevedo and Oliveira [2]. □

Now, we control the approximation between our sum and what we would find if the blocks W_j^p were independent.

Lemma 7. *For p fixed, there exists a constant $C > 0$, such that*

$$\begin{aligned} |Ee^{itZ_{mp}} - \prod_{j=1}^m Ee^{\frac{it}{\sqrt{m}}W_j^p}| &\leq Ct^2 \sum_{k=1}^q |\sigma_p^2(x_k, y_k, x_k, y_k) - \sigma^2(x_k, y_k, x_k, y_k)| \\ &\quad + 2t^2 \sum_{k=1}^{q-1} \sum_{l=k+1}^q |\sigma_p^2(x_k, y_k, x_l, y_l) - \sigma^2(x_k, y_k, x_l, y_l)|. \end{aligned} \tag{18}$$

Proof. This is a convergence of Newman’s inequality [10] and asymptotic properties of the variances of $Z_{\tilde{n}}$ and of W_j^p mentioned above. □

Lemma 8. *For p fixed, we have*

$$|\prod_{j=1}^m Ee^{\frac{it}{\sqrt{m}}W_j^p} - e^{\frac{it^2\sigma_p^2}{2}}| \longrightarrow 0. \tag{19}$$

Proof. Just apply the Lindeberg condition to the variables $m^{-1/2}W_j^p$, $j = 1, \dots, m$ and take account of the fact that each W_j^p is a sum of p bounded variables, so it is also bounded. □

Now by summarizing the results, we have the asymptotic normality in the next theorem which can be obtained from Lemmas 6, 7 and 8. The proof is similar to the proof of Theorem 5.5 of [2] and, therefore is omitted.

Theorem 5. *The random vector $(\alpha_n(x_1, y_1), \dots, \alpha_n(x_q, y_q))$ converges in distribution to a gaussian centered vector with covariance matrix*

$$\Sigma = \begin{bmatrix} \sigma^2(x_1, y_1, x_1, y_1) & \sigma^2(x_1, y_1, x_2, y_2) & \dots & \sigma^2(x_1, y_1, x_k, y_k) \\ \sigma^2(x_2, y_2, x_1, y_1) & \sigma^2(x_2, y_2, x_2, y_2) & \dots & \sigma^2(x_2, y_2, x_k, y_k) \\ \dots & \dots & \dots & \dots \\ \sigma^2(x_k, y_k, x_1, y_1) & \sigma^2(x_k, y_k, x_2, y_2) & \dots & \sigma^2(x_k, y_k, x_k, y_k) \end{bmatrix}. \tag{20}$$

References

- [1] Alam, K. and Saxena, K.M.L., (1981). Positive dependent in multivariate distributions, *Commun. Statist-Theor. Math.*, A 10, 1183-1196.
- [2] Azevedo, C. and Oliveira, P.E., (2000). Kernel-type estimation of bivariate distribution function for associated random variables. *In New Trends in Probability and Statistics*, 17-25.

- [3] Donsker, M. D., (1951). An invariant principle for certain probability limit theorems. *Mem. Amer. Math. Soc.*, 6.
- [4] Henriques, C. and Oliveira, P.E.,(2003). Estimation of a two dimensional distribution function under association. *J. Statist. Planning Inf.*, 113, 137-150.
- [5] Jabbari Nooghabi, H., Azarnoosh, H.A. and Fakoor, V., (2009). Almost sure convergence of two-dimensional distribution function under negative association. *To appear in Journal of Applied Probability and Statistics*.
- [6] Joag-Dev, K. and Proschan, F.,(1983). Negative associated of random variables, with application. *Ann. Statist.*, 11, 286-295.
- [7] Liang, H. Y. and Su, C., (1988). Convergence in the law of logarithm for *NA* sequences. *Chinese Sci. Bull.*, 43, 1919-1925(in Chinese).
- [8] Matula, P., (1992). A note on the almost sure convergence of sums of negatively dependent random variables. *Statist. Probab. Lett.*, 15, 209-213.
- [9] Newman, C.M., (1980). Normal fluctuations and the FKG inequalities. *Comm. Math. Phys.*, 74, 119-128.
- [10] Newman, C., (1984), Asymptotic independence and limit theorems for positively and negatively dependent random variables, in: *Inequalities in Statistics and Probability*, Y. L. Tong, ed, IMS Lecture Notes- Monograph Series, Vol. 5, 127-140.
- [11] Roussas, G.G., (1994). Asymptotic normality of random fields of positively or negatively associated processes. *J. Multivariate Anal.*, 50, 152-173.
- [12] Sadikova, S.M., (1966). Two-dimensional analogies of an inequality of Esseen with applications to the central limit theorem. *Theory Probab. Appl.*, 11, 325-335.
- [13] Shao, Q.M., (2000). A comparison theorem on maximal inequality between negatively associated and independent random variables. *J. Theore. Probab.*, 13, 343-356.
- [14] Su, C. and Chi, X., (1998). Some result on CLT for nonstationary *NA* sequences. *Acta. Math. Appl. Sinica*, 21, 9-21.
- [15] Su, C., Zhao, L.C. and Wang, Y.B., (1997). Moment inequalities and weak convergence for negatively associated sequences. *Science in China (series A)*, 40, 172-182.
- [16] Yuan, M., Su, C. and Hu, T., (2003). A central limit theorem for random fields of negatively associated process. *J. Theore. Probab.*, 16, 309-323.