

# Proceedings

of the British Society of Animal Science  
and the Agricultural Research Forum

## 2010

### Advances in Animal Biosciences

This book is part of a series which is a companion to the journal ANIMAL



- 257 The effect of offering grass silage alone or in combination with legume:cereal wholecrop silage on methane emissions of Holstein steers  
P C Kennedy, L E Dawson, D J Kilpatrick
- 258 The effect of electromagnetic water treatment on *in vitro* methane production  
M O'Brien, P O'Kiely
- 259 Effect of legume and perennial ryegrass herbage on *in vitro* methane output using the total gas production technique  
A Navarro-Villa, M O'Brien, S Lopez, T M Boland, P O'Kiely
- 260 *In vitro* methane output of perennial ryegrass produced under four grazing management regimes and sampled throughout the growing season  
P Purcell, M O'Brien, T M Boland, M O'Donovan, P O'Kiely
- 261 The effect of sward maturity on the *in vitro* digestibility and methane production of sward components  
C J Quinlan, M B Lynch, M O'Brien, A Navarro, T M Boland
- 262 Effect of sward maturity on the dry matter intake, enteric methane emission and milk solids production of pasture grazed dairy cows  
M H Deighton, C M Wims, B M O'Loughlin, E Lewis, M O'Donovan

## FEED EVALUATIONS/TECHNIQUES

- 263 Effect of adding different levels of probiotic on *in vitro* gas production of noodle waste  
M Besharati, A Taghizadeh, A Ansari
- 264 Effect of peppermint (*Mentha piperita*) essential oil on *in vitro* gas production parameters of lucerne hay and cottonseed hulls  
E Jani, M Danesh Mesgaran, A R Vakili, A Soleimani, H Jahani-Azizabadi
- 265 *In vitro* gas production parameters of chickpea (*Cicer arietinum* L.) by-product  
E Abdi Ghezeljeh, M Danesh Mesgaran
- 266 Use of *in situ* technique to evaluate three weed forages  
M Kazemi, A M Tahmasbi, R Valizadeh, A R Vakili, M M Moheghi
- 267 Kinetic of *in vitro* gas production of high fat sunflower meal treated with sodium hydroxide and or formal dehyde by rumen bacteria+protozoa  
M Bojarpour, T Mohammadabadi, M Danesh Mesgaran, M Chaji
- 268 The kinetic of *in vitro* gas production of tannic acid treated sunflower meal with or without polyethylene glycol  
T Mohammadabadi, M Chaji, S Tabatabaei
- 269 Nitrogen fractionations, *in situ* ruminal degradation and post-ruminal crude protein disappearance of over heat and overheat-xylose processed guar meal  
H Jahani-Azizabadi, M Danesh Mesgaran, A R Vakili, M Vatandoost, M Mojtahedi, E Abdi Ghezeljeh, A Hojjat Panah, A Fanaie-Nokar
- 270 *In vitro* first order dry matter disappearance kinetics of chemically and physically treated cottonseed hulls  
A Famarzi Garmroodi, M Danesh Mesgaran, A R Vakili, A R Heravi Moussavi, A Tahmasbi, H Jahani-Azizabadi
- 271 Use of white rot fungi to improve the feed value of rice straw  
J W Cone, J J P Baars, A S M Sonnenberg
- 272 Cultivation of oyster mushrooms (*Pleurotus* species) to improve the *in vitro* dry matter digestibility of wheat straw for feeding to ruminants  
H Omed, A Avagyan, M Hale, J Gibbons
- 273 Evaluation of condensed tannin content of some native tanniniferous plants from semi-arid regions in Brazil  
R C Lucas, A L Abdalla, M E Q Vieira, J D F Gomes, M R R S Peçanha, M T Lima, R Moura, B Berenchtein, A S Morsy, Y A Soltan
- 274 Chemical composition and dry matter degradability coefficients of Fennel seed  
M Kazemi, A M Tahmasbi, R Valizadeh, M Danesh Mesgaran, A A Naserian

## Chemical composition and dry matter degradability coefficients of Fennel seed

M Kazemi, A M Tahmasbi, R Valizadeh, M Danesh Mesgaran, A A Naserian

Ferdowsi University of Mashhad, Mashhad, Khorasan Razavi, Islamic Republic of Iran

Email: miton62@yahoo.com

**Introduction** Fennel (*Foeniculum vulgare*), is a plant species in the genus *Foeniculum* (treated as the sole species in the genus by most botanists). It is a member of the family *Apiaceae* (formerly the *Umbelliferae*). It is a hardy, perennial, umbelliferous herb, with yellow flowers and feathery leaves. Fennel is widely cultivated, both in its native range and elsewhere, for its edible, strongly-flavoured leaves and seeds. Its aniseed flavor comes from anethole, an aromatic compound also found in anise and star anise, and its taste and aroma are similar to theirs, though usually not as strong. Seeds of the fennel plant are widely used in many of the culinary traditions of the world. There are historical anecdotes that fennel is a galactagogue improving the milk supply of a breastfeeding mother (Crellin *et al.*, 1989). This use, although not supported by direct evidence, is sometimes justified by the fact that fennel is a source of phytoestrogens, which promote growth of breast tissue (Mark, 2000). The effects in animals is unclear, thus determination of these effects will be important for animals especially ruminants. The objective of this experiment was determination of DM degradability and chemical composition of Fennel seeds.

**Materials and methods** Samples of Fennel seed were obtained from Golchekan-Zamani factory in the Razavi Khorasan district. The samples were analyzed for dry matter (DM), crude protein (CP), neutral detergent fiber (NDF), acid detergent fiber (ADF), and organic matter based on procedure described in AOAC methods (2000). Measurements of *in situ* DM were performed in 3 rumen fistulated bull cows (fed at maintenance) using the nylon bag technique (Krskov and McDonald, 1979). Nylon bags which were approximately (9×18 cm) containing 5 g (2mm screen) were incubated in the rumen of fistulated bull cows for 2, 4, 8, 16, 24, 48, 72, 96 and 120 h. There were 4 replications per treatment. The rate and extent of DM degradation were estimated according to the equation:  $P = a + b(1 - e^{-ct})$ . Effective degradability (ED) was calculated according to equation:  $ED = a + (b \times c)/(c + k)$ , assuming an outflow rate (k) of 0.02 h<sup>-1</sup>. The data were analyzed using the ANOVA procedure of SAS (SAS Institute, 2002).

**Results** Degradable coefficients and ruminal disappearance of DM are summarized in Table 1. The chemical composition of fennel seed is shown in Table 2.

**Table 1** *In situ* degradability coefficients disappearance (Mean with SEM) of DM of Fennel seed.

Fennel seed	
Degradation parameters	
a <sup>1</sup>	0.32±0.01
b <sup>2</sup>	0.46±0.01
c <sup>3</sup>	0.037±0.003
a+b <sup>4</sup>	0.58±0.02
Effective degradability	
k=0.02	0.62±0.03
k=0.03	0.58±0.03

<sup>1,2,3,4</sup> constants in the equation  $P=a+b(1-e^{-ct})$  where P = level of degradation at time t; a=readily soluble fraction; b=insoluble fraction but degradable in rumen; c=rate of degradation of b per hour and a+b = potentially degradable fraction.

**Table 2** Chemical composition(g/kg) of Fennel seed

Fennel seed	
Composition (%)	
DM	914±4
ADF	382±13
NDF	421±15
CP	182±11
ASH	117±2

**Conclusions** The crude protein and Ash contents of fennel seed, evaluated in the present study, were almost high. Rapidly and slowly degradable fractions of fennel seed for DM indicated that more than 0.58 of this feed could be digested in the rumen. High digestibility of fennel seed and high nutritional value of it shows that it can supply intermediate protein for animals.

**Acknowledgments** The authors thank the Ferdowsi University of Mashhad and Excellence Centre for Animal Science for financial and technical support.

## References

- AOAC, 2000. Official Methods of Analysis, 17<sup>th</sup> ed. Association of Official Analytical Chemists. Washington, DC.
- Crellin, J.K., Philpott, J. and Tommie Bass, A.L. 1989. Herbal Medicine Past and Present. Duke University Press, 207-208.
- Krskov, E.R. and McDonald, I. 1979. Journal of Agricultural Science 92, 499-504.
- Mark, A. P. 2000. Alpha Books. ISBN 0028639480, 142
- SAS Institute INC.2002. SAS User's Guide: Statistics. Statistical Analysis Systems Institute Inc. Cary NC.