

## Crude Protein Fractions and *in vitro* Gas Production of Alfalfa Silages Treated with Pistachio by-Products Extract

### Research Article

A. Mokhtarpour<sup>1\*</sup>, A.A. Naserian<sup>1</sup>, R. Valizadeh<sup>1</sup>, M. Danesh Mesgaran<sup>1</sup> and F. Pourmollae<sup>2</sup>

<sup>1</sup> Department of Animal Science, Faculty of Agriculture, Ferdowsi University of Mashhad, Mashhad, Iran

<sup>2</sup> Agricultural Jihad Organization of Khorasan Razavi, Mashhad, Iran

Received on: 7 Apr 2014

Revised on: 1 Jul 2014

Accepted on: 31 Jul 2014

Online Published on: Jun 2015

\*Correspondence E-mail: [am.mokhtarpour@stu-mail.um.ac.ir](mailto:am.mokhtarpour@stu-mail.um.ac.ir)

© 2010 Copyright by Islamic Azad University, Rasht Branch, Rasht, Iran

Online version is available on: [www.ijas.ir](http://www.ijas.ir)

### ABSTRACT

Two experiments were conducted to evaluate the crude protein fractionation, *in vitro* gas production and the biological effects of tannins in alfalfa silage treated with different levels of pistachio by-products (PB) extract. PB was soaked in water with a ratio of 1:5 (w/v) for 12 h. After filtering and concentrating, the crude extract was added to fresh alfalfa to a final concentration of 1%, 2% and 3% tannin dry matter (DM) and then ensiled for 60 days. Four treatments were as follows: alfalfa silage without addition of PB extract as a control (T0), PB tannin added at 1% (T1), 2% (T2) and 3% DM (T3). Treating alfalfa silage decreased the formation of non-protein nitrogen (NPN) compounds and increased crude protein fraction B2 during ensilage. Cumulative *in vitro* gas production after 96 h incubation, organic matter digestibility (OMD) and metabolizable energy (ME) decreased in T2 and T3 compared to the control. The gas production and short chain fatty acids (SCFA) of treated silages increased in the presence of polyethylene glycol (PEG) which indicates that the presence of phenolic compounds and tannins depressed the gas production. It can be concluded that treating alfalfa silage with PB extract can reduce the nitrogen losses during ensilage and degradability of organic matter (OM) by microbial inhibition.

**KEY WORDS** crude protein fractionation, *in vitro* gas production, pistachio by-products, polyethylene glycol, tannin bioassay.

### INTRODUCTION

It is well documented that the nutritive value of ensiled leguminous forage such as alfalfa is limited by rapid and excessive degradation of protein during ensiling (Albrecht and Muck, 1991) and rumen fermentation (Broderick, 1995). Improving forage quality by reducing proteolysis during ensiling will improve N utilization by ruminants (Givens and Rulquin, 2004). There has been increasing interest in reducing the rate and extent of protein degradation during ensiling and rumen fermentation by using natural products such as tannins. Tannins are secondary plant metabolites that react with proteins, carbohydrates and min-

erals thus retard their digestion (McSweeney *et al.* 2001). Pistachio (*Pistacia vera*) by-products (PB) contain phenolic compounds and tannins ranging from 9.06-14.57% and 4.97-8.67% DM, respectively, depending on the method of extraction including solvent, particle size and time of extraction (Mokhtarpour *et al.* 2014). The *in vitro* gas production method is widely used to evaluate the nutritive value of foodstuffs (Getachew *et al.* 2002) and is closely related to digestibility and energetic value (Menke and Steingass, 1988). Many papers have been published on the use of the gas production technique as a method to study the anti-nutritive factors (Makkar *et al.* 1995; Getachew *et al.* 2008; Kumara Mahipala *et al.* 2009) and is seemed to be more

efficient than other *in vitro* techniques in determining the nutritive value of feeds containing tannins (Getachew *et al.* 2002).

The potential of PB extract addition to alfalfa during silage making and its effect on gas production has not yet been exploited. Therefore, the aims of this study were to evaluate the effect of treating alfalfa silage with PB extract on *in vitro* gas production and assaying the biological effect of tannin by using PEG.

## MATERIALS AND METHODS

### Pistachio by-products extract preparation

Sun-dried pistachio by-products (PB) were obtained from a pistachio de-hulling factory in Feizabad (Khorasan Razavi Province, Iran) which is located on the north east part of Iran at 35 °01' N latitude and 58 °78' E longitude. Pistachio by-products were soaked in water by a ratio of 1:5 (w/v) at room temperature for 12 h. The contents were filtered through 4 layers of cheesecloth and then were concentrated by a rotary evaporator at 40 °C.

### Preparation of alfalfa silages

A second regrowth of alfalfa was harvested at early bloom. The alfalfa was chopped to a length of 3 cm with a cutter. Alfalfa herbage was ensiled in 12 plastic buckets with the capacity of 500 g. PB extract was added to fresh alfalfa at 0, 40, 80 and 120 mL/kg of fresh weight to achieve final concentrations of 0%, 1%, 2% and 3% tannin as tannic acid equivalent (24% DM), respectively. Three replicate treatments were set up and all were ensiled for 60 days at room temperature (25 °C±1) in a dark place.

### *In vitro* gas production

Approximately 200 mg dry weight of the samples were weighed in triplicate and placed in 120 mL gas-tight culture bottles following the gas pressure transducer technique of Theodorou *et al.* (1994). The culture bottles were filled with 30 mL of buffered rumen fluid. Rumen fluid was obtained from three fistulated bulls before the morning feeding. The donor animals were fed with corn silage and concentrate with a ratio of 70:30 twice daily. The rumen fluid collection and composition of buffer were performed according to the Menke and Steingass (1988) procedure. The culture bottles containing samples and mixture of rumen fluid and buffer were incubated in a water bath at 39 °C. Gas pressure in the head-space of each bottle was read from the display unit after 2, 4, 8, 12, 24, 48, 72 and 96 hours' incubation and the corresponding gas volume was determined by recording the volume of gas displaced into the syringe barrel. Bottles were shaken gently after each gas production reading without removing them from the water

bath. This procedure was performed twice. Total gas values were corrected for blank incubation. Cumulative gas production data were fitted to the model of Ørskov and McDonald (1979):

$$Y = b(1 - e^{-ct})$$

Where:

Y: gas produced at time.

b: gas production from the insoluble but fermentable fraction (mL).

c: gas production rate constant for b.

t: incubation time (h).

The OM digestibility, ME and NE<sub>L</sub> were calculated to the following equations according to Menke and Steingass (1988):

$$\text{OMD (\%)} = 14.88 + 0.8893 \text{ GP} + 0.448 \text{ CP} + 0.651 \text{ Ash}$$

$$\text{ME (MJ/kg DM)} = 2.20 + 0.1357 \text{ GP} + 0.057 \text{ CP}$$

$$\text{NEI (MJ/kg DM)} = 0.54 + 0.0959 \text{ GP} + 0.038 \text{ CP} + 0.001733 \text{ EE2}$$

Where:

GP: net production in 24 h in mL/200 mg of DM.

CP: % of DM, EE is % DM and ash is % of DM.

### *In vitro* tannin bioassay

*In vitro* gas production was carried out using PEG in order to identify the biological effects of tannins as described by Makkar *et al.* (1995). Briefly, 500 mg dry weight of silage samples were incubated without and with 1 g PEG (MW=6000) and filled with 40 mL rumen liquor and buffer mixture in triplicate. Culture bottles were placed in a water bath at 39 °C and gas production was recorded after 2, 4, 6, 8, 10, 12 and 24 hours' incubation. Short chain fatty acids were calculated by the equation of Getachew *et al.* (2002):

$$\text{SCFA (mmol/40 mL)} = -0.00425 + 0.0222 \times \text{gas (mL)}$$

### Chemical composition

Dry matter content of fresh alfalfa was determined by drying in an oven at 100 °C to a constant weight (AOAC 2005). Ash (method 942.05) ether extract (method 920.39) and CP (Kjeldahl N×6.25) content of silages was determined by the block digestion method using copper catalyst and steam distillation into boric acid (method 2001.11) on behr S 5 steam distillation unit (behr Labor Technik GmbH, Germany) as described in AOAC (2005). Crude protein fractions of alfalfa silages were determined by the method

of Licitra *et al.* (1996). NPN was determined by precipitation of true protein in tungstic acid solution. Soluble protein was determined using borate phosphate buffer. The amount of insoluble protein in acid detergent and neutral detergent were determined after measuring ADF and NDF (Van Soest *et al.* 1991). For total phenolics (TP) quantification, approximately 200 mg ( $\pm 0.001$ ) silage samples was extracted in 10 mL 70% aqueous acetone (v/v) in quadruplicate by using an ultrasonic bath for 20 min. After centrifugation (3000 $\times$ g, 4 °C, 10 min) the supernatant was collected and kept in refrigerator (4 °C). Non-tannin phenolics (NTP) were determined following absorption of tannins in total phenolic extract to insoluble polyvinylpyrrolidone (PVPP). Total phenolics and non-tannin phenolics were determined by folin-ciocalteu reagent using tannic acid (Merck GmbH, Darmstadt, Germany) as a standard, expressing results as tannic acid equivalent. Total tannins (TT) were calculated as the difference between TP and NTP (Makkar, 2000). Condensed tannins (CT) concentration was measured on phenolics supernatant using the butanol-HCl reagent (Makkar, 2000). The values of CT were expressed as leucocyanidin equivalent.

### Statistical analysis

Data from chemical composition and gas production were analyzed using the GLM procedure in SAS (2001) as a completely randomized design. Means were separated by the least squares means procedure (LSMEANS) when a significant ( $P < 0.05$ ) treatment effect was observed. For tannin bioassay, data were analyzed as a completely randomized design with a 4  $\times$  2 factorial arrangement treatment.

The model included the effects of PB tannin level, presence or absence of PEG and the interaction between tannin level and PEG.

## RESULTS AND DISCUSSION

The amount of total phenolics, tannins and CP fractions of silages are presented in Table 1. Total phenolics and tannins increased by increasing PB extract. Treating alfalfa silage with PB extract decreased fraction A and increased the concentrations of fractions B1 and B2. The reduction in fraction A (NPN compounds) of treated silages can be attributed to tannin binding to forage proteins and preventing already formed tannin-protein complexes from enzyme hydrolysis or may be due to the action of tannins to bind to plant proteolytic enzymes (Guo *et al.* 2007). Although changes in activity of proteolytic enzymes were not determined in our study, decreased in NPN concentration may not be associated with inhibition of proteolytic enzymes activities as the pH values for all treatments were above 4.5 (data not shown).

**Table 1** Phenolic compounds (% DM) and crude protein fractions (% CP) of alfalfa silages

Item	Treatments				SEM
	T0	T1	T2	T3	
<b>Phenolic compounds (% DM)</b>					
Total phenolics	1.42 <sup>d</sup>	2.78 <sup>c</sup>	4.56 <sup>b</sup>	7.01 <sup>a</sup>	0.791
Total tannins	0.61 <sup>d</sup>	1.11 <sup>c</sup>	2.21 <sup>b</sup>	3.17 <sup>a</sup>	0.375
Condensed tannins	ND	0.12 <sup>c</sup>	0.25 <sup>b</sup>	0.35 <sup>a</sup>	0.043
<b>CP fractions (% CP)</b>					
A	62.76 <sup>a</sup>	57.02 <sup>b</sup>	58.01 <sup>b</sup>	58.26 <sup>b</sup>	0.883
B1	0.45 <sup>b</sup>	0.97 <sup>b</sup>	1.96 <sup>a</sup>	2.52 <sup>a</sup>	0.310
B2	28.80 <sup>b</sup>	33.25 <sup>a</sup>	32.40 <sup>a</sup>	32.36 <sup>a</sup>	0.686
B3	5.08	5.07	4.66	3.72	0.248
C	2.93 <sup>b</sup>	3.69 <sup>a</sup>	2.99 <sup>b</sup>	3.15 <sup>ab</sup>	0.151

T0: control; T1: 1% of DM PB tannin; T2: 2% of DM PB tannin; T3: 3% of DM PB tannin.

SEM: standard error of the means; ND: not detected.

The means within the same row with at least one common letter, do not have significant difference ( $P > 0.05$ ).

Henderson (1993) stated that the optimal pH for plant peptidase activity is between 5-7 and it will be reduced at pH below 4. Our results are consistent with Tabacco *et al.* (2006) and Guo *et al.* (2007) who reported that addition of tannic acid to alfalfa decreased the NPN concentration of silages. An increase in fraction B1 and B2 was associated with a reduction in fraction A of silages treated at 2% and 3% tannin.

However, a trend ( $P = 0.06$ ) for more concentrated fraction B1 was observed in T1 than T0. The concentration of fraction B3 was not affected by treatments. Treating alfalfa silage at 1% tannin (T1) increased fraction C compared with T0 and T2. Fraction C are heat-damaged proteins, proteins associated with lignins and tannin-protein complexes which are not degradable in the rumen and are indigestible in the intestine (Krishnamoorthy *et al.* 1982; Licitra *et al.* 1996). *In vitro* gas production and fermentation characteristics of alfalfa silages are shown in Table 2.

**Table 2** cumulative *in vitro* gas productions (mL/200 mg DM) after 24 and 96 h of incubation and estimated parameters of alfalfa silages

Item	Treatments				SEM
	T0	T1	T2	T3	
<b>Cumulative gas production</b>					
24 h	48.7 <sup>a</sup>	44.8 <sup>ab</sup>	41.2 <sup>bc</sup>	35.2 <sup>c</sup>	1.75
96 h	78.3 <sup>a</sup>	76.4 <sup>a</sup>	67.6 <sup>b</sup>	54.5 <sup>c</sup>	2.93
<b>Estimated parameters</b>					
b	76.8 <sup>a</sup>	75.3 <sup>a</sup>	65.7 <sup>b</sup>	52.6 <sup>c</sup>	2.98
c	0.044 <sup>b</sup>	0.038 <sup>c</sup>	0.042 <sup>b</sup>	0.049 <sup>a</sup>	0.001
OMD (%)	73.3 <sup>a</sup>	70.8 <sup>ab</sup>	65.3 <sup>bc</sup>	60.8 <sup>c</sup>	1.73
ME (MJ/kg DM)	9.94 <sup>a</sup>	9.31 <sup>ab</sup>	8.44 <sup>bc</sup>	7.55 <sup>c</sup>	0.306
NEI (MJ/kg DM)	6.04 <sup>a</sup>	5.60 <sup>ab</sup>	4.98 <sup>bc</sup>	4.36 <sup>c</sup>	0.215

T0: control; T1: 1% of DM pistachio by-products tannin; T2: 2% of DM pistachio by-products tannin; T3: 3% of DM pistachio by-products tannin.

b: gas production from the insoluble but fermentable fraction (mL/200 mg DM); c: gas production rate constant (mL/h); OMD: organic matter digestibility; ME: metabolizable energy; NEI: net energy for lactation; SEM: standard error of the means.

The means within the same row with at least one common letter, do not have significant difference ( $P > 0.05$ ).

There were significant differences among the treatments. Addition of PB extract (tannin) lowered gas production with a decrease in gas production from the insoluble but fermentable fraction along with both OMD and nutritive value of silages. However, PB tannin at 1% DM had no effect on these parameters. These results are in accordance with those obtained by [Tabacco et al. \(2006\)](#) who reported that treating alfalfa silage with hydrolysable tannin (chestnut) at the rate of 2%, 4% and 6% DM decreased gas production and estimated parameters including OMD, ME and NE<sub>L</sub>. The rate constant of gas production (c) decreased in T1, but increased in T3. The lower gas production and OMD in treated silages could be attributed to the inhibitory effects of phenolic compounds, especially tannins to the rumen microorganisms' activity. It is suggested that high concentration of condensed tannin (more than 5.5% DM) has negative effects on digestibility *in vivo* ([Min et al. 2003](#)). In the present study, using culture bottles, low levels of tannins (2%, 3% DM) inhibited rumen fermentation and it seems that rumen microbial population could not counteract the effects of tannins.

This is because the *in vitro* gas production technique is a static system compared to a dynamic system (*in vivo*). As 200 mg alfalfa silage was incubated in 30 mL medium, the rumen microorganisms were able to utilize alfalfa silage as a sole energy source. However, in an *in vivo* experiment, a tannin-treated feed is in combination with other feeds and a large population of microorganisms ([Tabacco et al. 2006](#)). [Getachew et al. \(2002\)](#) reported that over 90% of the variation in gas production was explained by SCFA produced by incubation of browsed leaves. Hence, the amount of gas production reflects the extent of feed degradability. The higher gas production in control and silage treated with 1% tannin compared to other treatments indicated the higher extent of fermentation of OM. [Tabacco et al. \(2006\)](#) reported that the gas production decreased by 10.2%, 7.9% and 10.2% when chestnut tannin was added to alfalfa silage at the rate of 2%, 4% and 6% DM respectively. In our experiment, the gas production decreased by 8.0%, 15.4% and 27.7% in T1, T2 and T3 respectively.

**Table 4** *In vitro* gas production at 24 h of incubation (mL/0.5g DM) and estimated SCFA (mmol/40mL) without and with polyethylene glycol of alfalfa silages

Item	Tannin level								SEM
	T0		T1		T2		T3		
	-PEG	+PEG	-PEG	+PEG	-PEG	+PEG	-PEG	+PEG	
GV	84.4	96.0	82.0	99.1	78.2	97.3	71.7	99.2	2.18
SCFA	1.88	2.14	1.82	2.20	1.84	2.17	1.60	2.21	0.049
Item	Significance of effects								
	TL		PEG		TL × PEG				
	GV	0.09	< 0.001						0.01
SCFA	0.09	< 0.001						0.01	

T0: control; T1: 1% of DM pistachio by-products tannin; T2: 2% of DM pistachio by-products tannin and T3: 3% of DM pistachio by-products tannin.

TL: tannin level; GV: cumulative gas volume at 24 h of incubation (mL/0.5g DM); SCFA: short-chain fatty acids (mmol/40mL) and PEG: polyethylene glycol.

SEM: standard error of the means.

This difference may be due to higher inhibitory effect of PB tannin than chestnut tannin. Moreover, other factors such as concentration of phenolic compounds may affect rumen microorganisms' activity or interfere with their enzyme secretion ([Yanez Ruiz et al. 2004](#)). Correlation between phenolic compound and gas production and estimated parameters are shown in Table 3. A decreased in gas production, OMD and ME in treated silages suggests a strong negative relationship between gas production, OM and ME and total phenolic and total tannins.

**Table 3** Correlation coefficients (r) of phenolic compounds and *in vitro* gas production characteristics of alfalfa silages

Item	Characteristics				
	GV	b	c	OMD	ME
Total phenolics	-0.998**	-0.982*	0.636	-0.994**	-0.998**
Total tannins	-0.989**	-0.981*	0.642	-0.999**	-0.998**

GV: cumulative gas volume at 24 h of incubation; b: gas production from the insoluble but fermentable fraction; c: gas production rate constant; OMD: organic matter digestibility and ME: metabolizable energy.

\* (P<0.05) and \*\* (P<0.01).

Addition of PEG increased the gas production in treated silages, but had no effect on control silage (Table 4). At 24 h of incubation, an increase in gas production as a result of PEG inclusion was 20.9%, 24.4% and 28.7% for T1, T2 and T3 respectively. Increase in gas production by the addition of PEG revealed a negative influence of tannins on digestibility ([Makkar, 2005](#)). Polyethylene glycol is a polymer that binds to tannins irreversibly and thereby increase the availability of nutrients ([Makkar et al. 1995](#); [Provenza et al. 2000](#)) resulting in increased in microbial activity and gas production ([Makkar, 2005](#)). Our results are consistent with [Osuga et al. \(2006\)](#), [Kumara Mahipala et al. \(2009\)](#) and [Yusef Elahi et al. \(2012\)](#). Therefore, an increase in gas production by adding PEG is considered a biological effect of tannins ([Makkar et al. 1995](#)). Short-chain fatty acids were also increased by PEG inclusion. [Getachew et al. \(2002\)](#) found a significant correlation between percent changes in gas production and SCFA by PEG addition and phenolic compounds.

## CONCLUSION

The results of our study showed that even lower level of PB tannin (1% DM) inhibited proteolysis during ensiling resulting in lower concentration of NPN compared to untreated silage. At moderate levels of tannins (2-3% DM) the gas production at 24 h and 96 h of incubation, ME and OMD decreased. PEG inclusion increased cumulative gas production at 24 h of incubation in PB extract treated silage. Therefore, it can be concluded that phenolic compounds especially tannins in PB can interfere with rumen microorganisms' activity leading to decreased proteolysis and OM fermentation.

## ACKNOWLEDGEMENT

The authors would like to acknowledge from Department of Animal Science of Ferdowsi University of Mashhad for the cooperation.

## REFERENCES

- Albrecht K.A. and Muck R.E. (1991). Proteolysis in ensiled forage legumes that vary in tannin concentration. *Crop Sci.* **31**, 464-469.
- AOAC. (2005). Official Methods of Analysis. 18<sup>th</sup> Ed. Association of Official Analytical Chemists, Arlington, VA, USA.
- Broderick G.A. (1995). Performance of lactating dairy cows fed either alfalfa silage or alfalfa hay as the sole forage. *J. Dairy Sci.* **78**, 320-329.
- Getachew G., Makkar H.P.S. and Becker K. (2002). Tropical browses: contents of phenolic compounds, *in vitro* gas production and stoichiometric relationship between short chain fatty acid and *in vitro* gas production. *J. Agric. Sci.* **139**, 341-352.
- Getachew G., Pittroff W., Putnam D.H., Dandekar A., Goyal S., and De Peters E.J. (2008). The influence of addition of gallic acid, tannic acid, or quebracho tannins to alfalfa hay on *in vitro* rumen fermentation and microbial protein synthesis. *Anim. Feed Sci. Technol.* **140**, 444-461.
- Givens D.I. and Rulquin H. (2004). Utilisation by ruminants of nitrogen compounds in silage-based diets. *Anim. Feed Sci. Technol.* **114**, 1-18.
- Guo X., Zhou H., Yu Z. and Zhang Y. (2007). Changes in the distribution of nitrogen and plant enzymatic activity during ensilage of lucerne treated with different additives. *Grass Forage Sci.* **62**, 35-43.
- Henderson N. (1993). Silage additives. *Anim. Feed Sci. Technol.* **45**, 35-56.
- Krishnamoorthy U., Muscato T.V., Sniffen C.J. and Van Soest P.J. (1982). Nitrogen fractions in selected feedstuffs. *J. Dairy Sci.* **65**, 217-225.
- Kumara Mahipala M.B.P., Krebs G.L., McCafferty P. and Gunaratne L.H.P. (2009). Chemical composition, biological effects of tannin and *in vitro* nutritive value of selected browse species grown in the west Australian Mediterranean environment. *Anim. Feed Sci. Technol.* **153**, 203-215.
- Licitra G., Hernandez T.M. and Van Soest P.J. (1996). Standardization of procedures for nitrogen fractionation of ruminant feeds. *Anim. Feed Sci. Technol.* **57**, 347-358.
- Makkar H.P.S. (2000). Quantification of Tannins in Tree Foliage. A Laboratory Manual for the FAO/IAEA Co-ordinated Research Project on Use of Nuclear and Related Techniques to Develop Simple Tannin Assays for Predicting and Improving the safety and Efficiency of Feeding Ruminants on Tanniniferous Tree Foliage. Joint FAO/IAEA, FAO/IAEA of Nuclear Techniques in Food and Agriculture. Animal Production and Health Sub-program, FAO/IAEA Working Document. IAEA, Vienna, Austria.
- Makkar H.P.S., Blümmel M. and Becker K. (1995). *In vitro* effects and interactions of tannins and saponins and fate of tannins in rumen. *J. Sci. Food Agric.* **69**, 481-493.
- Makkar H.P. (2005). *In vitro* gas methods for evaluation of feeds containing phytochemicals. *Anim. Feed Sci. Technol.* **123**, 291-302.
- Mcsweeney C.S., Palmer B., McNeill D.M. and Krause D.O. (2001). Microbial interaction with tannin: nutritional consequences for ruminants. *Anim. Feed Sci. Technol.* **91**, 83-93.
- Menke K.H. and Steingass H. (1988). Estimation of the energetic feed value obtained from chemical analysis and gas production using rumen fluid. *Anim. Res. Dev.* **28**, 7-55.
- Min B.R., Barry T.N., Attwood G.T. and McNabb W.C. (2003). The effect of condensed tannins on the nutrition and health of ruminants fed fresh temperate forages: a review. *Anim. Feed Sci. Technol.* **106**, 3-19.
- Mokhtarpour A., Naserian A.A., Valizadeh R., Danesh Mesgaran M. and Pourmollae F. (2014). Extraction of phenolic compounds and tannins from pistachio by-products. *Annu. Rev. Res. Biol.* **4**, 1330-1338.
- Ørskov E.R. and McDonald I. (1979). The estimation of protein degradability in the rumen from incubation measurements weighted according to rate of passage. *J. Agric. Sci.* **92**, 499-503.
- Osuga I.M., Abdulrazak S.A., Ichinohe T., Ondiek J.O. and Fujihara T. (2006). Degradation characteristics and tannin bioassay of some browse forage from Kenya harvested during the dry season. *Anim. Sci. J.* **77**, 414-421.
- Provenza F.D., Burritt E.A., Perevolotsky A. and Silanikove N. (2000). Self-regulation of intake of polyethylene glycol by sheep fed diets varying in tannin concentrations. *J. Anim. Sci.* **78**, 1206-1212.
- SAS Institute. (2001). SAS<sup>®</sup>/STAT Software, Release 9.1. SAS Institute, Inc., Cary, NC, USA.
- Tabacco E., Borreani G., Crovetto G.M., Galassi G., Colombo D. and Cavallarin L. (2006). Effect of chestnut tannin on fermentation quality, proteolysis and protein rumen degradability of alfalfa silage. *J. Dairy Sci.* **89**, 4736-4746.
- Theodorou M.K., Williams B.A., Dhanoa M.S., McAllan A. and France J. (1994). A simple gas production method using a pressure transducer to determine the fermentation kinetics of ruminant feeds. *Anim. Feed Sci. Technol.* **48**, 185-197.
- Van Soest P.J., Robertson J.B. and Lewis B.A. (1991). Methods for dietary fiber, neutral detergent fiber and non starch polysaccharides in relation to animal nutrition. *J. Dairy Sci.* **74**, 3583-3597.

Yanez Ruiz D.R., Moumen A., Martin Garcia A.I. and Molina Alcaide E. (2004). Ruminal fermentation and degradation patterns, protozoa population, and urinary purine derivatives excretion in goats and wethers fed diets based on two-stage olive cake: effect of PEG supply. *J. Anim. Sci.* **82**, 2023-2032.

Yousef Elahi M., Rouzbehan Y. and Rezaee A. (2012). Effects of phenolic compounds in three oak species on *in vitro* gas production using inoculums of two breeds of indigenous Iranian goats. *Anim. Feed Sci. Technol.* **176**, 26-31.

---