

Please ensure that your abstract fits into one column on one page and complies with the *Instructions to Authors* available from the Abstract Submission web page.

## **The vulnerability and quality protection zoning maps of Bojnourd aquifer (N Iran) using Fuzzy-AHP-DRASTIC techniques**

HOSSEIN MOHAMMADZADEH AND FAEZEH DAVODI

Groundwater Research Center (GRC), Ferdowsi University of Mashhad, Mashhad, Iran, P.O.Box: 91775-1436

Groundwater is the main resources of water supply for agricultural, drinking, and industrial purposes in Iran and, therefore, it is essential to protect aquifers from pollution. In this paper, the vulnerability and quality protection zoning maps of Bojnourd aquifer, the capital of N-Khorasan province - North of Iran, was prepared using Fuzzy-AHP- DRASTIC techniques in GIS environment. The groundwater risk map ( $C_{Risk}$ ) was prepared by integrating the aquifer intrinsic vulnerability map (DRASTIC- $D_i$ ) with the land-use map. To increase the accuracy of  $D_i$  and  $C_{Risk}$  maps, they were calibrated using aquifer nitrate concentrations map. After correcting the weights and ratings of the model parameters, the aquifer specific vulnerability (modified  $D_i$ - $D_{i-m}$ ) and special risk-taking ( $C_{Risk-m}$ ) maps were prepared. Finally, by improving the weights and rates of required parameters using analytical hierarchy process (AHP) and fuzzy logic techniques, the risk-taking maps of the aquifer were prepared with AHP-DRASTIC, modified AHP-DRASTIC, Fuzzy-AHP-DRASTIC and modified Fuzzy-AHP-DRASTIC models. The correlation coefficient between the vulnerability/Q-protection zoning maps with aquifer nitrate map indicate that the  $C_{Risk-m}$  model, is the best model for preparing the Bojnourd aquifer vulnerability zoning map, and about 26, 23 and 51 percent of the aquifer have high, very high and very susceptible potential for contamination.

**Keywords:** Aquifer vulnerability, Quality protection zoning map, DRASTIC model, AHP, Fuzzy Logic, Bojnourd plain.