



Priming impacts on seed quality of tomato (*Lycopersicon esculentum*) under salinity condition

Mohsen Poursoltan Hojagan¹, Hossein Arouiee^{*2}, Seyyed Jalal Tabatabaei,
Seyyed Hossein Neamati²

¹Horticultural Science (Olericulture), Ferdowsi University of Mashhad, Iran

²Department of Gardening, Faculty of Agriculture, Ferdowsi University of Mashhad, Iran

³Department of Physiology of Plant Nutrition, Shahed University, Iran

Article published on September 30, 2017

Key words: Germination rate, Priming, Salinity, Seedling, Tomato

Abstract

This study aimed to investigate the seed priming effects on the quality of tomato seeds under salinity and non-salinity conditions. The factorial design was carried out in the form of completely randomized design (CRD) with four replications. The treatments included priming (Control, tryptophan, proline and arginine with five mM concentration) and levels of salinity (0, 30, 60mM NaCl). As salinity increased, means of germination percentage, germination rate, length of radicle, plumule and seedling, and seedling dry weight decreased. Seed priming with proline compensated radicle length reductions caused by salinity. Seed priming had an improving effect on seedling dry weight and increased germination percentage, germination rate and length of radicle. Proline-primed seeds improved seed quality better than tryptophan and arginine-primed seeds.

*Corresponding Author: Hossein Arouiee ✉ arouiee@um.ac.ir

Introduction

Seed priming is a simple method for improving seedling vigor and establishment and also crop performance in the field (McDonald, 2000). The useful effects of priming have been proved for several crops such as barley (Abdulrahmani *et al.*, 2007), maize (Parera and Cantliffe, 1994), lentil (Ghassemi-Golezani *et al.*, 2008), chickpea and pinto bean (Ghassemi-Golezani *et al.*, 2010). These priming effects are linked with nucleic acid repairing and building up, increased protein synthesis, and the membrane repairing (McDonald, 2000). In addition, priming improves anti-oxidative enzymes activities of the seeds treated (Hsu *et al.*, 2003). Seed germination rate and uniformity and seedling emergence may be increased with the early improvements (Ghassemi-Golezani *et al.*, 2010), particularly in stress conditions, especially under stressful conditions (Ghassemi-Golezani *et al.*, 2008).

Two critical factors to crop production under salt-stress are rapid seed germination and stand establishment (Ashraf and Foolad, 2005). Seed germination may be affected by soil salinity through osmotic potential external to the seed preventing water uptake, or Na⁺ and Cl⁻ ions (Khajeh-Hosseini *et al.*, 2003). As previous studies showed, seed germination, seedling emergence and growth can be improved with priming under saline conditions (Sivirtepe *et al.*, 2003). The results of the studies on the effect of salt stress on growth indicated a relation between plant length decrease and sodium chloride concentration increase (Memon *et al.*, 2010).

The over salinity of the soil is one of the main factors that limits the spread of plants in their natural habitats. It is an ever-increasing problem in arid and semi-arid regions (Shanon, 1986). Fisher and Turner (1978) estimate that arid and semi-arid lands represent around 40% of the earth's area. The property of salinity tolerance is not a simple attribute, but it is an outcome of various features that depend on different physiological interactions, which are difficult to determine. Jamil *et al.* (2005) showed that concentration increase resulted in the adverse effects of salinity on seed germination rate and percentage,

and leaf number (Jamil *et al.*, 2005; Gama *et al.*, 2007; Ha *et al.*, 2008). Several studies show that changes in salinity concentration, type of salt present, or type of plant species affect the fresh and dry weights of the seedling either negatively or positively, (Saffan, 2008; Rui *et al.*, 2009; Taffouo *et al.*, 2010; Memon *et al.*, 2010). Satisfactory stand establishment and higher yields can be ensured with high and rapid (Ghassemi-Golezani *et al.*, 2010).

The morphological appearance presented by the plant in response to salinity, may not be enough to determine its effect, so it is important to recognize other physiological and biochemical factors, including toxic ions, osmotic potential, lack of elements and other physiological and chemical disorders, as well as the interactions between these various stresses (Hasegawa *et al.*, 2000). Although tomato response to salt stress has previously been evaluated, but there is not so much information on the effects of priming on tomato performance under salt stress. So, this study aims mainly at investigating the effects of priming on tomato germination under saline and non-saline conditions.

Materials and methods

An experimental study was designed to evaluate tomato seed quality parameters under saline and non-saline conditions. The proposed study was conducted in the Vegetable Seed Laboratory, Research Center of Agricultural and Natural Resources of Eastern Azarbaijan Province, Iran, to determine Priming impacts on seed quality of tomato (*Lycopersicon esculentum*) under salinity condition.

Priming Treatment

The treatments included priming (Control, tryptophan, proline and arginine with 5 mM concentration) and salinity levels (0, 30 and 60mM). Tomato seeds (*Lycopersicon esculentum*) were grouped into four subsamples, and one of them was left as control (unprimed) and the other three subsamples were prepared for priming Proline and arginine were Other sub-samples which were soaked in tryptophan, at 15C° for four hours Following priming, seeds were completely washed with distilled

water for one minute and then dried at 20-23 C° to its primary moisture in the laboratory.

Emergence Test

Laboratory tests were carried out as factorial, based on CRD design at the Seed Technology Laboratory of the University of Mashhad, Iran. Four replicates of 25 seeds of each sub-samples (Control, tryptophan, proline and arginine) were placed between moist filter papers and germinated in an incubator adjusted at 20°C for 10 days. Salinity levels were created by supplementing the Hoagland solution with NaCl; control (only Hoagland’s solution), 0, 30 and 60mM NaCl.

Data Collection

Germination (protrusion of radicle by 2mm) was recorded in daily intervals. At the end, percentage of normal seedlings and seedling dry weight were determined. Mean germination rate was calculated based on the following equation of Ellis and Roberts, 1980:

$$\bar{R} = \frac{\sum n}{\sum D.n}$$

Where n is the number of seeds that germinated on day D, D is the number of days since the start of the test and \bar{R} is the mean germination rate.

Statistical Analysis

MSTATC software was used to analyze the data variance. To compare the means of each trait, Duncan test was applied at $p \leq 0.05$. Excel software was used to draw Fig.s.

Results and discussion

The analysis of variance of the laboratory data showed significant effects of priming on germination percentage, germination rate, length of radicle and seedling dry weight. However, priming had no significant effect on length of plumule and length of seedling (Table 1). The highest germination percentage, germination rate, length of radicle and seedling dry weight were achieved by P₃. Seeds of P₂ and P₄ and P₁ germinated later than those of P₃ (Table 2). Germination percentage, germination rate, length of radicle, length of plumule and length of seedling and seedling dry weight significantly affected by salinity (Table 1). Germination percentage and length of radicle for S₁ was significantly.

Higher than those for S₂ and S₃. The highest germination rate, length of plumule, length of seedling and seedling dry weight were also recorded for S₁ followed by those for S₂ (Table 2).

Table 1. Analysis of variance of the effects of seed priming and salinity on seed quality parameters.

Source	d.f	MS					
		Germination percentage	Germination rate	Length of plumule	Length of radicle	Length of seedling	Seedling dry weight
Replication	3	29.688	0.0001	1.815	0.103	0.095	0.00001
Priming	3	78.299*	0.021**	1.326	0.248**	1.807	0.0001*
Salinity	2	228.646**	0.012**	15.319**	44.445**	42.173**	0.0002*
Priming × Salinity	6	43.924	0.0001	2.077	0.311**	4.610	0.00001
Error	33	28.551	0.001	1.470	0.028	4.172	0.00002
C.V (%)	-	6.14	8.00	22.04	4.39	26.26	14.99

*, **: significant at $p \leq 0.05$ and $p \leq 0.01$.

Table 2. Means of seed quality parameters affected by priming and salinity.

Treatments	Germination percentage (%)	Germination rate (per day)	Length of plumule (cm)	Length of radicle (cm)	Length of seedling (cm)	Seedling dry weight (mg)
Priming						
P ₁	88.75b	0.250c	5.34a	3.601b	7.47a	0.0337c
P ₂	93.33ab	0.300b	5.51a	3.802a	7.76a	0.0370b
P ₃	94.58a	0.350a	5.77a	3.946a	7.87a	0.0396a
P ₄	91.25ab	0.286b	5.54a	3.832a	7.56a	0.0351bc
Salinity						
S ₁	96.25a	0.322a	6.234a	5.602a	9.234a	0.0394a
S ₂	90.63b	0.300a	5.877a	3.470b	8.072a	0.0371a
S ₃	89.06b	0.267b	4.389b	2.318c	6.028b	0.0325b

Different letters in each column for each treatment indicate significant difference at $p \leq 0.05$.

P₁, P₂, P₃ and P₄: unprimed seeds and priming for tryptophan, proline and arginine respectively.

The interaction of priming × salinity was significant for length of radicle (Table 1). Length of radicle of tomato seeds with different levels of priming decreased with increasing salinity, although this reduction in primed seeds (P₂, P₃ and P₄) was lower than that in unprimed seed lot (Fig. 1). However, the highest improvement in length of radicle due to priming was also observed in P₃ seed lot (priming with proline) (Fig. 1).

Germination percentage, germination rate, length of radicle, plumule and seedling length showed positive significant correlations with the seedling dry weight.

Germination percentage and plumule length had the highest positive correlation with seedling dry weight. Germination percentage and plumule length with radicle and seedling length also showed significant and positive correlation (Table 3).

Table 3. Correlation coefficients between seed quality parameters of tomato.

Traits	1	2	3	4	5	6
1. Germination percentage (%)	1					
2. Germination rate (per day)	0.782**	1				
3. Length of plumule (cm)	0.788**	0.631*	1			
4. Length of radicle (cm)	0.831**	0.565	0.720**	1		
5. Length of seedling (cm)	0.854**	0.554	0.715**	0.934**	1	
6. Seedling dry weight (mg)	0.888**	0.784**	0.82**	0.643*	0.696*	1

*, **: Significant at p≤0.05 and p≤0.01, respectively




Fig. 1. Means of length of radicle affected by priming and salinity.

P₁, P₂, P₃ and P₄: unprimed seeds and priming for tryptophan, proline and arginine respectively.

S₁, S₂ and S₃: Control and salinity with 30 and 60 mM NaCl, respectively.

Seed priming improved germination percentage, germination rate, length of radicle and seedling dry weight (Table 2). It might be due to early synthesis of nucleic acids. Ultimately, DNA, RNA and proteins resulted in the improvement of seed germination energy (Bray *et al.*, 1989). Rapid seed germination can result in larger seedling production (Abdulrahmani *et al.*, 2007; Ghassemi-Golezani *et al.*, 2010a, b). Osmo-priming had also positive effects on

barley seed germination and seedling growth (Abdulrahmani *et al.*, 2007), cucumber (Ghassemi-Golezani and Esmailpour, 2008), fennel (Neamatollahi *et al.*, 2009) and winter rapeseed (Ghassemi-Golezani *et al.*, 2010b).

Mean number of germination percentage, germination rate, length of radicle, length of plumule and length of seedling, which is resulted in decreasing seedling dry weight particularly under severe salinity (Table 2). Similar results have been reported by Bagheri and Sadeghipour (2009) for barley. Reduction in seedling dry weight due to salinity was also reported for maize and sunflower (Katerji *et al.*, 1996) and wheat (Yildirim and Bilge, 2010).

Reduction in germination and seedling dry weight due to salinity stress was also reported for cotton and wheat (Cullu, 2003), rice (Mahmood *et al.*, 2009). Salinity may delay the onset, reduce the rate, and increase the dispersion of germination events, leading to reductions in plant growth and final crop yield (Ashraf and Foolad, 2005). Significant and positive correlations among these traits (Table 3) indicate that improving each of the former traits can enhance tomato seedling dry weight. The highest positive and significant correlations were shown between germination percentage and plumule length and

seedling dry weight (Table 3) Proline-primed seeds improved grain yield per plant, better than arginine- and tryptophan-primed seeds. Thus, priming the seeds with amino acid can promote tomato seed germination rate and seedling dry weight, and can improve seed quality.

References

Abdulrahmani B, Ghassemi-Golezani K, Valizadeh M, Feizi-Asl V. 2007. Seed priming and seedling establishment of barley (*Hordeum vulgare* L.). Journal of Food, Agriculture and Environment **5**, 179-184.

Ashraf M, Foolad MR. 2005. Pre-sowing seed treatment: A shotgun approach to improve germination, plant growth and crop yield under saline and non-saline conditions. Advances in Agronomy **88**, 223-271.

Bagheri A, Sadeghipour O. 2009. Effects of salt stress on yield, yield components and carbohydrates content in four hullless barley (*Hordeum vulgare* L.) cultivars. Journal of Biological Sciences **9**, 909-912.

Bray CM, Davison PA, Ashraf M, Taylor RM. 1989. Biochemical changes during priming of leek seeds. Annals of Botany **63**, 185-193.

Çullu MA. 2003. Estimation of the effect of soil salinity on crop yield using remote sensing and geographic information system. Turkish Journal of Agriculture and Forestry **27**, 23-28.

Ellis RH, Roberts EH. 1980. Towards a rational basis for testing seed quality. In: Hebblethwaite PD, Ed. Seed production. London, UK: Butterworth's 605-635.

Fisher RA, Turner NC. 1978. Plant productivity, in arid and semi-arid zones. Annual Review of Plant Physiology **29**, 897-912.

Gama PBS, Inanaga S, Tanaka K, Nakazawa R. 2007. Physiological response of common bean (*Phaseolus vulgaris* L.) seedlings to salinity stress. African Journal of Biotechnology **6**, 79-88.

Ghassemi-Golezani K, Aliloo AA, Valizadeh M, Moghaddam M. 2008. Effects of different priming techniques on seed invigoration and seedling establishment of lentil (*Lens culinaris*). Journal of Food Agriculture and Environment **6**, 222-226.

Ghassemi-Golezani K, Chadordooz-Jeddi A, Nasrollahzadeh S, Moghaddam M. 2010a. Effects of hydro-priming duration on seedling vigor and grain yield of pinto bean (*Phaseolus Vulgaris* L.) cultivars. Notulae Botanicae Horti Agrobotanici Cluj-Napoca Journal **38**, 109-113.

Ghassemi-Golezani K, Esmailpour B. 2008. The effect of salt priming on the performance of differentially matured cucumber (*Cucumis sativus*) seeds. Notulae Botanicae Horti Agrobotanici Cluj-Napoca Journal **36**, 67-70.

Ghassemi-Golezani K, Jabbarpour S, Zehtab-Salmasi S, Mohammadi A. 2010b. Response of winter rapeseed (*Brassica napus* L.) cultivars to salt priming of seeds. African journal of agricultural research **5**, 1089-1094.

Ha E, Ikhajiagba B, Bamidele JF, Ogic-odia E. 2008. Salinity effects on young healthy seedling of *Kyllingia peruviana* collected from escravos, Delta state. Global Journal of Environmental Research **2**, 74-88.

Hasegawa PM, Bressan RA, Zhu JK, Bohnert HJ. 2000. Plant cellular and molecular responses to high salinity. Annual review of plant physiology and plant molecular biology **51**, 463-499.

Hsu SY, Hsu YT, Kao CH. 2003. The effect of polyethylene glycol on proline accumulation in rice leaves. Biologia Plantarum **46**, 73-78.

Jamil M, Lee CC, Rehman SU, Lee DB, Ashraf M, Rha ES. 2005. Salinity (NaCl) tolerance of brassica species at germination and early seedling growth. Electronic Journal of Environmental, Agricultural and Food Chemistry **4**, 970-976.

- Katerji N, Van Hoorn JW, Hamdy A, Karam F, Mastrorilli A.** 1996. Effect of salinity on water stress, growth and yield of maize and sunflower. *Agricultural Water Management* **30**, 237-249.
- Khajeh-Hosseini M, Powell AA, Bingham IJ.** 2003. The interaction between salinity stress and seed vigour during germination of soybean seeds. *Seed Science and Technology* **31**, 715-725.
- Mahmood A, Latif T, Khan MA.** 2009. Effect of salinity on growth, yield and yield components in basmati rice germplasm. *Pakistan Journal of Botany* **41**, 3035-3045.
- McDonald MB.** 2000. Seed priming. Seed technology and its biological basis.
- Memon SA, Hou X, Wang LJ.** 2010. Morphological analysis of salt stress response of pak Choi. *Electronic journal of environmental, agricultural and food chemistry* **9**, 248-254.
- Neamatollahi E, Bannayan M, Ghanbari A, Haydari M, Ahmadian A.** 2009. Does hydro and osmo-priming improve fennel (*Foeniculum vulgare*) seeds germination and seedlings growth? *Notulae Botanicae Horti Agrobotanici Cluj-Napoca* **37**, 190-194.
- Parera CA, Cantliffe DJ.** 1994. Pre-sowing seed priming. *Horticultural Reviews* **16**, 109-141.
- Rui L, Wei S, Mu-xiang C, Cheng-jun J, Min W, Bo-ping Y.** 2009. Leaf anatomical changes of *Burquiera gymnorhiza* seedlings under salt stress. *Journal of Tropical and Subtropical Botany* **17**, 169-175.
- Saffan SE.** 2008. Effect of salinity and osmotic stresses on some economic plants. *Research journal of agriculture and biological sciences* **4**, 159-166.
- Shanon MC.** 1986. New insights in plant breeding efforts for improved salt tolerance. *Horticultural Science & Technology* **6**, 96-99.
- Sivirtepe N, Sivirtepe HO, Eris A.** 2003. The effects of NaCl priming on salt tolerance in melon seedling grown under saline conditions. *Scientia Horticulturae* **97**, 229-237.
- Taffouo VD, Wamba OF, Yombi E, Nono GV, Akoe A.** 2010. Growth, yield, water status and ionic distribution response of three bambara groundnut (*Vigna subterranean* (L.) verdc.) landraces grown under saline conditions. *International Journal of Botany* **6**, 53-58.
- Yildirim M, Bilge B.** 2010. Responses of some wheat genotypes and their F2 progenies to salinity and heat stress. *Scientific Research and Essays* **5**, 1734-1741.