



An eight-coordinate zinc complex containing the highly pre-organized ligand 1,10-phenanthroline-2,9-dicarboxylic acid: Solvothermal synthesis, supramolecular structure and CSD studies

Masoud Mirzaei ^{a,*}, Azam Hassanpoor ^a, Hanie Alizadeh ^a, Mahnaz Gohari ^a, Alexander J. Blake ^b

^a Department of Chemistry, Faculty of Science, Ferdowsi University of Mashhad, Mashhad, Iran

^b School of Chemistry, University of Nottingham, Nottingham, NG7 2RD, UK

ARTICLE INFO

Article history:

Received 9 April 2018

Received in revised form

12 June 2018

Accepted 14 June 2018

Available online 15 June 2018

Keywords:

Pre-organized ligand

1,10-Phenanthroline-2,9-dicarboxylic acid

Zinc complex

π - π stacking interaction

Hydrogen bonding

ABSTRACT

A new anionic coordination complex of Zn(II) containing the highly pre-organized ligand 1,10-phenanthroline-2,9-dicarboxylate (PDA), $(\text{H}_3\text{O})_2[\text{Zn}(\text{PDA})_2] \cdot 2\text{CH}_3\text{NO}_2 \cdot 2\text{H}_2\text{O}$ (**1**), has been synthesized by solvothermal methods and characterized by elemental analysis, infrared spectroscopy and single-crystal X-ray diffraction. The last of these techniques revealed a coordination sphere around Zn(II) consisting of four N atoms and four O donors from two 1,10-phenanthroline-2,9-dicarboxylate ligands which features a distorted dodecahedral geometry. The ionic coordination motifs in **1** are held together by intermolecular O–H \cdots O, N–H \cdots O and C–H \cdots O hydrogen bonding, C–H \cdots π and π - π stacking interactions, to produce an extended 2D architecture.

© 2018 Elsevier B.V. All rights reserved.

1. Introduction

Currently, a key goal for chemists lies in the design and synthesis of new materials with significantly improved properties and performance. In this arena, supramolecular chemistry offers powerful synthetic tools such as the ability to vary the metal centre [1–4], its counter ions [5–7], the steric, conformational and coordinative features of the ligand [8,9] and the reaction conditions [10–13]. This flexibility allows us to exert control over the self-assembly process and construct desirable target architectures for potential real-world applications. Among these factors, the selection of suitable organic ligands plays a significant role [14]. Pre-organized ligands have proven to be important in inorganic, biomedical, environmental, and industrial chemistry [15]. Such ligands possess rigid backbones, resulting in minimal motion of the donor groups due to the conformation of the free ligand being closely similar to that in the metal complex [16], allowing the pre-organized ligands to coordinate selectively to metal ions. The H₂PDA ligand, with its planar, rigid phenanthroline skeleton and two carboxylate groups

at the 2 and 9 positions, is thereby one of the most prominent such ligands and is used to prepare coordination compounds with selected metal ions [17] and luminescent [18] or magnetic properties [19]. A search of the Cambridge Structural Database [CSD version 5.38 (version of October 2017)/Conquest version 1.21] [20] revealed 49 coordination compounds of the PDA ligand which is mostly frequently found coordinated to metal ions with ionic radii of ca. 1 Å. With various functional groups, this symmetrically chelating ligand provides non-covalent interactions such as hydrogen bonding and aromatic stacking for controlling the inter-growth of supramolecular networks. In previous work, we focused on design and synthesis of lanthanoid coordination frameworks containing H₂PDA [21]. As a continuation of our efforts to develop novel systems containing dicarboxylic acids [22–25] and in order to allow comparisons to be drawn with the previous work, we synthesized a zinc(II) coordination complex, $(\text{H}_3\text{O})_2[\text{Zn}(\text{PDA})_2] \cdot 2\text{CH}_3\text{NO}_2 \cdot 2\text{H}_2\text{O}$, (**1**), using the H₂PDA ligand. Complex **1** has been characterized by elemental analysis and infrared spectroscopy. Its solid-state structure was determined by single crystal X-ray diffraction analysis which showed that π - π stacking and hydrogen bonding interactions are crucial to the formation and generation of higher-dimensional (D) network.

* Corresponding author.

E-mail address: mirzaesh@um.ac.ir (M. Mirzaei).

2. Experimental

2.1. Materials and instrumentation

All commercially available chemicals were of reagent grade and were used without further purification, except for PDA which was synthesized according to literature methods [26]. Infrared spectra were recorded as KBr disks in the range (4000–600 cm^{-1}) on a Buck 500 Scientific spectrometer. Elemental analysis was carried out with a Thermo Finnigan Flash-1112 EA microanalyzer and Perkin-Elmer 2004(II) apparatus. Melting points were determined on a Electrothermal Thermo Scientific IA 9000 apparatus.

2.2. Synthesis of $(\text{H}_3\text{O})_2[\text{Zn}(\text{PDA})_2]\cdot 2\text{CH}_3\text{NO}_2\cdot 2\text{H}_2\text{O}\cdot (1)$

A mixture of H_2PDA (54 mg, 0.2 mmol) and $\text{ZnCl}_2\cdot 5\text{H}_2\text{O}$ (22 mg, 0.1 mmol) in deionized water/acetonitrile (15 ml; 1:1) was stirred for about 30 min. The mixture was then transferred into a 23 ml Teflon-lined autoclave and kept at 130 °C for 3 days. After slow cooling (5 °C \cdot h $^{-1}$) to the room temperature, crystals of the complex **1** could be isolated as translucent light-yellow tablet crystals (see Scheme 1). Yield: 53% (based on Zn). Anal. Calcd. For $\text{C}_{30}\text{H}_{30}\text{ZnN}_6\text{O}_{16}$: C, 45.27; H, 3.80; N, 10.56. Found: C, 46.85; H, 3.41; N, 11.25%. IR (KBr pellet, cm^{-1}): 3377(s), 1637(s), 1570 (s), 1466(m), 1361(s), 1313(m), 1247(w), 867(m), 828(w), 780(m). m.p. > 400 °C.

2.3. X-ray crystallography

Single-crystal data collection for **1** was performed on a Bruker SMART APEX II CCD area detector diffractometer, equipped with an Oxford Cryosystems open-flow nitrogen cryostat operating at 120 K, using ω scans and graphite-monochromated Cu $K\alpha$ radiation ($\lambda = 1.54184$ Å). The crystal was mounted on a thin glass fibre using paraffin oil. All data sets were corrected for Lorentz, polarization and absorption effects as specified in Table 1. The structure was solved by direct methods using SHELXS97 [27], completed with difference Fourier synthesis, and refined on F^2 by full-matrix least-squares using SHELXL97 [28].

3. Results and discussion

3.1. Vibrational spectrum

In addition to elemental analysis, complex **1** was initially characterized by its IR spectrum. The medium intense broad absorption in the 3377 cm^{-1} region indicates the presence of water molecules.


The observed bands at 1570 and 1361 cm^{-1} are attributed to the antisymmetric $\nu_{\text{as}}(\text{COO}^-)$ and symmetric $\nu_{\text{s}}(\text{COO}^-)$ stretching vibrations of carboxylate groups. The resulting value of $\Delta\nu$ ($\Delta\nu = \nu_{\text{as}} - \nu_{\text{s}} = 209$ cm^{-1}) is characteristic of the monodentate bonding mode of the carboxylate group. The band at 867 cm^{-1} is assigned as ring breathing mode of the phen ligand [29]. These assignments are supported by the results of the single-crystal X-ray diffraction analysis.

3.2. Description of crystal structure

Crystal data for compound **1** are shown in Table 1 while selected bond lengths and bond angles are given in Table 2 and hydrogen bond parameters are shown in Table 3. As shown in Fig. 1, molecular structure of **1** contains discrete 1,10-phenanthroline-2,9-dicarboxylate zinc(II) anions, hydronium cations, water and nitromethane molecules. In the anionic complex, each Zn^{II} ion is eight-coordinated and surrounded by four O and four N donors from two PDA ligands in a distorted dodecahedral geometry, with Zn–O and Zn–N distances of 2.579(3) and 2.141(3) Å, respectively. Each $(\text{PDA})^{2-}$ acts as a tetradentate chelating ligand and the ligands are almost perpendicular to each other in a *cis* arrangement. As seen

Table 1
Selected crystal data and structure refinement details for **1**.

Empirical formula	$\text{C}_{28}\text{H}_{12}\text{ZnN}_4\text{O}_8\cdot 2(\text{CH}_3\text{NO}_2)\cdot 2(\text{H}_3\text{O})\cdot 2(\text{H}_2\text{O})$
Formula weight, g mol^{-1}	795.97
Temperature, K	120(2)
Wavelength λ , Å	1.54184
Crystal system	Tetragonal
Space group	$I4_1/a$
$a/\text{Å}$	10.4342(2)
$b/\text{Å}$	10.4342(2)
$c/\text{Å}$	30.6744(6)
$V/\text{Å}^3$	3339.60(14)
Z	4
$D_c/\text{Mg m}^{-3}$	1.583
F_{000}	1640
μ/mm^{-1}	1.79
Crystal size/ mm^3	$0.26 \times 0.17 \times 0.03$
Space range, °	4.5 to 67.9
Reflections collected	7862
Reflection independent	1520
$R_1, wR_2 [I > 2\sigma(I)]$	0.0678, 0.1967
R_1, wR_2 (all data)	0.0697, 0.1987
Goodness-of-fit on F^2	1.05
Largest differences peak and hole, $e \text{ Å}^{-3}$	0.73 and -0.49



Scheme 1. The synthetic procedure of **1**.

Table 2
Selected bond lengths (Å) and angles (°) for **1**.

Zn1–N1	2.141(3)	Zn1–O2	2.579(3)
O2–Zn1–N1	67.16(9)	O2–Zn1–O2	149.10(9)
O2–Zn1–N1	143.72(9)	O2–Zn1–O2	94.07(9)
O2–Zn1–N1	76.55(9)	O2–Zn1–N1	79.31(9)
N1–Zn1–N1	76.64(8)	N1–Zn1–N1	127.99(8)

Table 3
Selected hydrogen bonding interactions present in **1**.

D–H···A	d(D–H)/Å	D(H–A)/Å	d(D–A)/Å	D–H–A/°
O2S–H2SA···O1	0.96	1.78	2.696(5)	158
O2S–H2SB···O1	0.93	1.83	2.719(5)	152
C3–H4···O2	0.95	2.44	3.325(11)	154




Fig. 1. Ellipsoid plot of **1** showing the atom-numbering scheme. Displacement ellipsoids are drawn at the 50% probability level.

from Table 4, the bond distances in **1** are longer than the values reported in the literature. Connectivity within the coordination compound is supplemented by O–H···O [O···O distances are 2.696(5) and 2.719(5) Å] and C–H···O [C···O distance is 3.325(11) Å] hydrogen bonds between (PDA)²⁻ carboxylate groups and hydronium cations. Hydronium cations connect discrete anionic complexes into 2D layers with the graph-set notation $R_8^4(16)$ along the *ac* plane (Fig. 2). Additionally, C–H···O interactions involving the coordinated O of carboxylate and an H atoms from an aromatic ring create $R_2^2(14)$ graph-sets along the crystallographic *c* axis (Fig. 3). It

Table 4
Comparison of the bond distances (Å) within some PDA complexes.

Chemical formula	M–O	Ref
(H ₃ O) ₂ [Zn(PDA) ₂]·2CH ₃ NO ₂ ·2H ₂ O·(1)	2.579	This work
Cu ₂ (PDA) ₂ (H ₂ O) ₂ ·2H ₂ O	1.904, 2.073	[30]
(Hpyda) ₂ [Cu(PDA) ₂]·10H ₂ O	2.165, 2.232	[31]
[Ni(PDA)(H ₂ O) ₃]·H ₂ O	2.116	[32]
[Co(PDA)(H ₂ O) ₃]·2H ₂ O	2.336	[33]
[Fe(HPDA)(H ₂ O) ₃] (ClO ₄) ₂ ·3H ₂ O·1/2H ₂ O ₂	2.120	[34]
[Mn(PDA)(H ₂ O) ₃]	2.350	[35]
(Hpyda)[Cr(PDA) ₂]·5H ₂ O	1.976, 1.953	[36]

Abbreviationpyda = pyridine-2,6-diamine.




Fig. 2. (Top) Hydrogen bonds present in **1** generate a 2D layer, here viewed along *ac* plane; (bottom) schematic representation of graph-sets.




Fig. 3. C–H···O hydrogen bonds generate $R_2^2(14)$ graph-sets along the *c* axis.

is worth mentioning that the three aromatic rings of the (PDA)²⁻ ligand play an important role in self-assembly via π – π [centroid-centroid distances 3.993 Å (Cg1···Cg2) and dihedral angle 18.73°] stacking interactions along the crystallographic *a* axis (Fig. 4).

Database studies. Upon examination of the crystal structure of **1** and the crystal structures identified in the CSD search the following interesting issues arise. First, in the 49 complexes containing PDA as ligand, the highest and lowest abundances are related to lanthanoid and main group metals, respectively. The reason is that the highly pre-organized PDA ligand with its rigid backbone skeleton tends to form complex with metals whose ionic radii are close to 1 Å. As a result, PDA complexes with first-row transition metals and main group metals are less stable than those with lanthanoids. Also, as Fig. 5 clearly shows, no complexes have previously been reported between PDA and Zn, making complex **1** the first. Scheme 2 represents coordination modes of PDA in transition metal complexes. In most cases, the PDA ligand




Fig. 4. Partial view of the crystal packing of **1** with indication of the π - π stacking interactions along the a axis.





Fig. 5. Abundance of coordination compounds containing PDA as a ligand as a function of metal centre.

does not use its entire donor capacity for link to metal ion, but in **1** two $(\text{PDA})^{2-}$ act as tetradentate chelating ligands and are almost perpendicular to each other in a *cis* arrangement. This observation can be related to the electron configuration of Zn^{2+} (d^{10}) which makes it similar to the main group metals.

4. Conclusions

We report herein the synthesis and structural characterisation of the first zinc complex involving $(\text{PDA})^{2-}$ ligand, $(\text{H}_3\text{O})_2[\text{Zn}(\text{PDA})_2] \cdot 2\text{CH}_3\text{NO}_2 \cdot 2\text{H}_2\text{O}$ (**1**). The presence of an organic ligand with three aromatic rings in these systems provides an extended net of noncovalent interactions, namely hydrogen bonds and π - π stacking interactions, which play a prominent role in the assembly of low-dimensional entities into a high-dimensional supramolecular network. Additionally, in order to investigate the



Scheme 2. Coordination modes of PDA in the first-row transition metal complexes.

influence of the metal ion on the geometry of PDA complexes, we undertook a statistical study based on the results of database searching. The results clearly show that the coordination mode of the pre-organized ligand is strongly dependent on ionic radius and electron configuration of metal centre.

Acknowledgements

M.M. wishes to thank to the Ferdowsi University of Mashhad Research Council for their financial support of this project (Grant No. 1/30354). M.M. gratefully acknowledged the Cambridge Crystallographic Data Centre (CCDC) for access to the Cambridge Structural Database.

Appendix A. Supplementary material

CCDC 1835767 contains the supplementary crystallographic data for this paper. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre via www.ccdc.cam.ac.uk/data_request/cif. Supplementary data associated with this article can be found, in the online version, at <https://doi.org/10.1016/j.ica.2014.10.032>.

References

- [1] M. Mirzaei, H. Eshtiagh-Hosseini, Z. Bolouri, Z. Rahmati, A. Esmaeilzadeh, A. Hassanpoor, A. Bauza, P. Ballester, M. Barceló-Oliver, J.T. Mague, B. Notash, A. Frontera, Rationalization of noncovalent interactions within six new MII/8-aminoquinoline supramolecular complexes (MII = Mn, Cu, and Cd): a combined experimental and theoretical DFT study, *Cryst. Growth Des.* 15 (2015) 1351–1361.

- [2] H. Eshtiagh-Hosseini, M. Mirzaei, S. Zarghami, A. Bauza, A. Frontera, J.T. Mague, M. Habibi, M. Shamsipur, Crystal engineering with coordination compounds of Ni^{II}, Co^{II}, Cr^{III}, and Cd^{II} bearing 2,6-dicarboxy-4-hydroxypyridine and 9-aminoacridine fragments driven by different natures of the face-to-face $\pi \dots \pi$ stacking: X-ray structures, DFT calculations, and solution studies, *CrystEngComm* 16 (2014) 1359–1377.
- [3] M.C. Hong, Y.J. Zhao, W.P. Su, R. Cao, M. Fujita, Z.Y. Zhou, A.S.C. Chan, A nanometer-sized metallosupramolecular cube with O_h symmetry, *J. Am. Chem. Soc.* 122 (2000) 4819–4820.
- [4] M. Mishra, K. Tiwari, A.K. Singh, V.P. Singh, Versatile coordination behavior of a multi-dentate Schiff base with manganese(II), copper(II) and zinc(II) ions and their corrosion inhibition study, *Inorg. Chim. Acta.* 425 (2015) 36–45.
- [5] C.B. Aakeroy, N.R. Champness, C. Janiak, Recent advances in crystal engineering, *CrystEngComm* 12 (2010) 22–43.
- [6] L. Carlucci, G. Ciani, D.M. Proserpio, S. Rizzato, Complex interwoven polymeric frames from the self-assembly of silver(I) cations and sebacitrile, *Chem. Eur. J.* 5 (1999) 237–243.
- [7] M. Mirzaei, H. Eshtiagh-Hosseini, M. Mohammadi Abadeh, M. Chahkandi, A. Frontera, A. Hassanpoor, Influence of accompanying anions on supramolecular assembly and coordination geometry in Hg^{II} complexes with 8-aminoquinoline: experimental and theoretical studies, *CrystEngComm* 15 (2013) 1404–1413.
- [8] G.G. Sezer, O.Z. Yeşilel, O. Şahinc, A.D. Burrows, Zinc(II) and cadmium(II) coordination polymers containing phenylenediacetate and bis(imidazole-1-ylmethyl)benzene linkers: the effect of ligand isomers on the solid state structures, *J. Solid State Chem.* 252 (2017) 8–21.
- [9] P. Manna, B.K. Tripuramallu, S.K. Das, Influential role of geometrical disparity of linker and metal ionic radii in elucidating the structural diversity of coordination polymers based on angular dicarboxylate and bis-pyridyl ligands, *Cryst. Growth Des.* 14 (2014) 278–289.
- [10] L.K. Das, A.M. Kirillov, A. Ghosh, Discrete 0D and polymeric 2D and 3D derivatives assembled from [(CuL)₂Zn]²⁺ and dicyanamide blocks (H₂L = salen type Schiff bases): genuine supramolecular isomers with distinct topologies, *CrystEngComm* 16 (2014) 3029–3039.
- [11] H. Eshtiagh-Hosseini, M. Mirzaei, M. Biabani, V. Lippolis, M. Chahkandi, C. Bazzicaluppi, Insight into the connecting roles of interaction synthons and water clusters within different transition metal coordination compounds of pyridine-2,5-dicarboxylic acid: experimental and theoretical studies, *CrystEngComm* 15 (2013) 6752–6768.
- [12] H.R. Khavasi, B. Mir Mohammad Sadegh, Temperature-dependent supramolecular motif in coordination compounds, *Inorg. Chem.* 49 (2010) 5356–5358.
- [13] M.A. Withersby, A.J. Blake, N.R. Champness, P.A. Cooke, P. Hubberstey, W.-S. Li, M. Schroder, Solvent control in the synthesis of 3,6-bis(pyridin-3-yl)-1,2,4,5-tetrazine-bridged cadmium(II) and zinc(II) coordination polymers, *Inorg. Chem.* 38 (1999) 2259–2266.
- [14] D. Sertphon, D.J. Harding, P. Harding, H. Adams, Effect of the β -diketonate ligand on the spin states of [Ni(β -dkt)(NH₂-quin)] complexes, *Polyhedron* 30 (2011) 2740–2745.
- [15] D.L. Melton, D.G. VanDerveer, R.D. Hancock, Complexes of greatly enhanced thermodynamic stability and metal ion size-based selectivity, formed by the highly preorganized non-macrocyclic ligand 1,10-phenanthroline-2,9-dicarboxylic acid. A Thermodynamic and crystallographic study, *Inorg. Chem.* 45 (2006) 9306–9314.
- [16] D.J. Cram, J.M. Cram, Design of complexes between synthetic hosts and organic guests, *Acc. Chem. Res.* 11 (1978) 8–14.
- [17] M.A. Lashley, A.S. Ivanov, V.S. Bryantsev, S. Dai, R.D. Hancock, Highly preorganized ligand 1,10-phenanthroline-2,9-dicarboxylic acid for the selective recovery of uranium from seawater in the presence of competing vanadium species, *Inorg. Chem.* 55 (2016) 10818–10829.
- [18] B.V. Harbuzaru, A. Corma, F. Rey, J.L. Jordá, D. Ananias, L.D. Carlos, J. Rocha, A miniaturized linear pH sensor based on a highly photoluminescent self-assembled europium(III) metal–organic framework, *Angew. Chem. Int. Ed.* 48 (2009) 6476–6479.
- [19] Q. Chen, M.H. Zeng, L.Q. Wei, M. Kurmoo, A multifaceted cage cluster, [Co^{II}₆O₁₂ \rightarrow X]⁻ (X = Cl⁻ or F⁻): halide template effect and frustrated magnetism, *Chem. Mater.* 22 (2010) 4328–4334.
- [20] C.R. Groom, I.J. Bruno, M.P. Lightfoot, S.C. Ward, The Cambridge structural database, *Acta Crystallogr. B* 72 (2016) 171–179.
- [21] M. Alipour, O. Akintola, A. Buchholz, M. Mirzaei, H. Eshtiagh-Hosseini, H. Goris, W. Plass, Size-dependent self-assembly of lanthanide-based coordination frameworks with phenanthroline-2,9-dicarboxylic acid as pre-organized ligand in hybrid materials, *Eur. J. Inorg. Chem.* (2016) 5356–5365.
- [22] A. Najafi, M. Mirzaei, A. Bauzá, J.T. Mague, A. Frontera, The roles of H-bonding, π -stacking, and antiparallel CO \dots CO interactions in the formation of a new Gd(III) coordination polymer based on pyridine-2,6-dicarboxylic acid, *Inorg. Chem. Commun.* 83 (2017) 24–26.
- [23] M. Shahbazi, F. Mehrzad, M. Mirzaei, H. Eshtiagh Hosseini, J.T. Mague, M. Ardalani, M. Shamsipur, Synthesis, single crystal X-ray characterization, and solution studies of Zn(II), Cu(II)-, Ag(I)- and Ni(II)-pyridine-2,6-dipicolinate N-oxide complexes with different topologies and coordination modes, *Inorg. Chim. Acta.* 458 (2017) 84–96.
- [24] M. Mirzaei, H. Eshtiagh Hosseini, M. Alipour, A. Bauzá, J.T. Mague, M. Korabik, A. Frontera, Hydrothermal synthesis, X-ray structure and DFT and magnetic studies of a (H₂SiW₁₂O₄₀)²⁻ based one-dimensional linear coordination polymer, *Dalton Trans.* 44 (2015) 8824–8832.
- [25] M. Mirzaei, H. Eshtiagh Hosseini, Z. Karrabi, B. Notash, catena-Poly[[di- μ -2-aqua-hexaquaabis(μ -3-4-oxidopyridine-2,6-dicarboxylato)tri manganese(II)] trihydrate]: a new one-dimensional coordination polymer based on a trinuclear Mn^{II} complex of chelidamic acid, *Acta Crystallogr. C* 69 (2013) 1140–1143.
- [26] A. Angeloff, J.C. Daran, J. Bernadou, B. Meunier, The ligand 1,10-phenanthroline-2,9-dicarbaldehyde dioxime can act both as a tridentate and as a tetradentate ligand—synthesis, characterization and crystal structures of its transition metal complexes, *Eur. J. Inorg. Chem.* (2000) 1985–1996.
- [27] G.M. Sheldrick, Phase annealing in SHELX-90: direct methods for larger structures, *Acta Crystallogr. A* 46 (1990) 467.
- [28] G.M. Sheldrick, A short history of SHELX, *Acta Crystallogr. A* 64 (2008) 112–122.
- [29] K. Nakamoto, *Infrared, Raman Spectra, Of Inorganic and Coordination Compounds*, Part B, fifth ed., Wiley, New York, 1997.
- [30] L.L. Fan, C.J. Li, Z.S. Meng, M.L. Tong, Metal-mediated one-step in situ oxidation of 2,9-dimethyl-1,10-phenanthroline and formation of transition-metal and lanthanoid complexes, *Eur. J. Inorg. Chem.* (2008) 3905–3909.
- [31] A. Moghimi, R. Alizadeh, H. Aghabozorg, A. Shokravi, M.C. Aragoni, F. Demartin, F. Isaia, V. Lippolis, A. Harrison, A. Shokrollahi, M. Shamsipur, Ion pairing, H-bonding, and π - π interactions in copper(II) complex-organonetworks derived from a proton-transfer compound of the 1,10-phenanthroline-2,9-dicarboxylic acid, *J. Mol. Struct.* 750 (2005) 166–173.
- [32] Y.B. Xie, J.R. Li, X.H. Bu, The synthesis and structure of the NiII complex of 1,10-phenanthroline-2,9-dicarboxylate: a three-dimensional network via hydrogen bonding interactions, *J. Mol. Struct.* 741 (2005) 249–253.
- [33] Z.F. Shi, Z.Q. Gao, J.Z. Gu, Triaqua(1,10-phenanthroline-2,9-dicarboxylato)cobalt(II) dihydrate, *Acta Crystallogr. E* 66 (2010) m372.
- [34] N.J. Williams, N.E. Dean, D.G. VanDerveer, R.C. Luckay, R.D. Hancock, Strong metal ion size based selectivity of the highly preorganized ligand PDA (1,10-phenanthroline-2,9-dicarboxylic acid) with trivalent metal ions. A crystallographic, fluorometric, and thermodynamic study, *Inorg. Chem.* 48 (2009) 7853–7863.
- [35] O. Tamer, D. Avci, Y. Atalay, B. Cosut, Y. Zorlu, M. Erkovan, Y. Yerli, Synthesis, X-ray structure, spectroscopic characterization and nonlinear optical properties of triaqua(1,10-phenanthroline-2,9-dicarboxylato)manganese(II) dihydrate: a combined experimental and theoretical study, *J. Mol. Struct.* 1100 (2015) 605–613.
- [36] A. Moghimi, R. Alizadeh, M.C. Aragoni, V. Lippolis, H. Aghabozorg, P. Norouzi, F. Isaia, S. Sheshmani, Synthesis, characterization and X-ray crystal structure of a chromium(III) complex obtained from a proton-transfer compound containing 1,10-phenanthroline-2,9-dicarboxylic acid and 2,6-pyridinediamine, *Z. Anorg. Allg. Chem.* 631 (2005) 1941–1946.