

Niche modelling of the potential distribution of the Egyptian Vulture *Neophron percnopterus* during summer and winter in Iran, to identify gaps in protected area coverage

AZITA FARASHI and MOHAMMAD ALIZADEH-NOUGHANI

Summary

An analysis of the extent of overlap between habitats of the Egyptian Vulture *Neophron percnopterus* during summer and winter in Iran, and Iranian protected areas, discovered slight overlap between the two. Our study aimed to survey suitable habitats of the Egyptian Vulture during summer and winter in Iran using MaxEnt, a species distribution modelling method, and compare the results with the locations and span of protected areas. Model fitness was assessed using the area under the ROC-plot and True Skill Statistics. We found that habitat suitability in Iran varies for the Egyptian Vulture during summer and winter. Summer visitors are scattered in the north and west of Iran, concentrated along the Zagros and Alborz mountain ranges. Winter visitors and residents are scattered in coastal areas in the south. Habitat protection for winter visitors was the greatest, with 10% of suitable habitat covered, while less than 10% of suitable habitat for residents and summer visitors fell within protected areas. The Egyptian Vulture is categorised as 'Endangered' on the IUCN Red List. Different environmental factors influence the suitability of habitat for the species. While some factors such as NDVI and anthropogenic disturbance influence all visitors in a similar way, the effects of others such as elevation and distance from protected areas vary for summer and winter visitors. Since designation of protected areas in Iran is mainly determined by the distribution of mammal species, protection of bird habitats is often overlooked. Suitable habitat for the species should be considered when selecting protected areas in future.

Introduction

Knowledge of a species' habitat can help to understand its ecology and guide management and conservation efforts. Gaps in our understanding of population biology and habitats of birds of prey hinder accurate designation of conservation status as well as formulation and implementation of successful conservation strategies. Evidence-based methods, which take into account biological factors, species' distribution, and population density (Naveda-Rodriguez, 2013, 2015), are required and predictive models, such as species distribution models (SDMs), have become an essential tool for researchers in predicting and assessing the distribution of species by quantifying species-environment relationships (Loiselle *et al.* 2003, Guisan and Thuiller 2005, Elith and Leathwick 2009, Franklin 2010, Morán-Ordóñez *et al.* 2017, Flessner *et al.* 2017, Ramesh *et al.* 2017). SDMs may be used to identify areas with high conservation value, demarcate future protected areas, initiate conservation measures, and forecast how climate change will affect species' distribution (Stralberg *et al.* 2009, Austin and Van Niel 2011, Virkkala *et al.* 2013, Rempel and Hornseth 2017). They have been used to show that the combined effects of global environmental change (e.g. through changes in land use) are likely to cause unparalleled rates of environmental change

and biodiversity loss (Parmesan 2006, Riordan and Rundel 2014). Much attention has been paid to threats such as direct habitat loss, and range shifts due to climate change in the 21st century (Thuiller *et al.* 2005), but habitat conversion and degradation, resulting from overexploitation and agricultural activities, constitute potentially more imminent threat to species persistence (Sala *et al.* 2000, Maxwell *et al.* 2016). Minimising harm to ecosystems from present activities will help face challenges posed by climate change in the future (Maxwell *et al.* 2016).

There is also a long history of using SDMs to assess the effectiveness of protected areas (Farashi and Shariati 2017) by applying gap analyses, based on species distribution. Historically, protected areas have been designated in areas with low economic value at the expense of regions with ecological and conservation importance which may be insufficiently protected (Rodrigues *et al.* 2004). Venter *et al.* (2018) showed that neither existing nor new protected areas cover locations with high concentrations of threatened vertebrate species.

The Egyptian Vulture *Neophron percnopterus* has been listed on the IUCN Red List as 'Endangered' since 2007 (IUCN 2017). Multiple factors have contributed to population declines of the species (Nikolaus 1984, 2006, Cuthbert *et al.* 2006, Ogada *et al.* 2016) including electrocution by power lines (Nikolaus 1984, 2006, Angelov *et al.* 2013), wind farms (Ferrer and Penteriani 2008, Penteriani *et al.* 2011), direct persecution and changes in food availability (Velevski *et al.* 2015), and poisoning by drugs (e.g. diclofenac) and lead (from gun-shot) (Thiollay 2006a,b, Hernández and Margalida, 2009). Poisoning of wild predators by hunters and livestock farmers that inadvertently kills vultures is one of the leading causes of population decline (Ogada *et al.* 2016). Because the relative importance of threats to the survival of Egyptian Vultures is poorly understood, there is a critical need for research into causes of mortality and potential conservation actions that may halt and reverse population declines. Although the species has a wide range, from the Mediterranean to India, as well as parts of eastern and southern Africa, global estimates of its current population are between 12,000 and 38,000 mature individuals. Egyptian Vulture is also recognized as an endangered species in Iran. The most important factors threatening the species in Iran include loss of habitat due to changes in land cover and land use, and illegal trade and hunting. Current knowledge about this species remains extremely poor. Egyptian Vulture occurs in different regions of Iran as a resident, summer visitor, and winter visitor (Kaboli *et al.* 2015). Summer visitors are scattered in the north and west of Iran, while residents and winter visitors are scattered in coastal areas of the south.

The aims of this study were: (1) to identify factors affecting the distribution of the Egyptian Vulture during summer and winter in Iran, (2) to identify suitable habitats of the Egyptian Vulture during summer and winter in Iran, and (3) to evaluate whether the current protection network is sufficient to cover the suitable habitats of the Egyptian Vulture during summer and winter in Iran.

Materials and methods

Study area

Iran is located in the Middle East (24°–40°N, 44°–64°E), serving as a bridge between the Indian subcontinent, the Arabian Peninsula, Central Asia, and Europe. Because of high habitat diversity Iran is one of the most important countries in the Middle East for biodiversity conservation. Water scarcity, land degradation and pollution are the three main threats to biodiversity in Iran (Iranian Department of Environment 2015).

Species presence data

Egyptian Vultures inhabit different regions of the country as (1) residents that breed and remain in a specific area all year round, (2) summer visitors that arrive in spring, remain throughout the

breeding season, and leave the country in autumn and (3) winter visitors that arrive in autumn, remain throughout the winter, and leave the country in spring. Data on the historical distribution of the summer and winter visitors in Iran were obtained from atlas distribution maps (Kaboli *et al.* 2015). These maps show the distribution of each species in a point-based format and are the product of data recorded in the last 50 years. New information reported by Iranian and foreign ornithologists was added later. Species distribution data were obtained from observations and signs (e.g. nests, camera traps).

To create maps for this study, the map of Iran was converted into a 1 × 1 km grid and all reliable information on the species' presence was transferred to the cells. The records of summer and winter visitors were screened in ArcGIS 10.3 (ESRI 2014). Presence locations of residents, summer visitors and winter visitors covered 6, 136, and 14 cells respectively. Significant spatial autocorrelation of residuals was identified using Moran's correlograms ('I' predicted HSI for the species record; De Marco *et al.* 2008) and by Spatial Analysis in Macroecology (SAM; Rangel *et al.* 2006). To determine the significance of Moran's I, a randomisation test with 9,999 Monte Carlo permutations was used, corrected for multiple testing. In cases of significant spatial autocorrelation, we constrained testing and training data: the first distance lags showing positive spatial autocorrelation were used to establish a distance threshold, and data pairs closer to each other than the threshold were placed in the same data partition (Parolo *et al.* 2008).

Environmental predictors

The following six eco-geographical variables were selected for the two distribution models according to the species' time of presence in the area: vegetation, bioclimatic, topographic, and food availability; land cover, and human footprint index (Table 1). Land cover predictors were derived from 30-m Landsat Enhanced Thematic Mapper Plus (ETM+) imagery. Land cover data were obtained from the Iranian Forests, Range and Watershed Management Organization (IFRWO) and Iranian Department of Environment in 2014 (Table 1). Using neighbourhood analysis, the ratio of different land cover types in a 1-km radius from the centre of each cell was calculated. Normalized Difference Vegetation Index (NDVI, calculated as the average of seasonal values in the 2000–2016 period) was compiled based on Moderate Resolution Imaging Spectrometer at 1-km resolution (MODIS, available on: <https://lpdaac.usgs.gov>). Nineteen bioclimatic predictors were collected from the WorldClim 1.4 database (Hijmans *et al.* 2005) (Table 1) at 1-km resolution and aggregated for October–March for winter and April–August for summer. The four most important topographic factors (mean and standard deviation of elevation, and slope of all raster cells included in a 1-km radius) were calculated based on the Shuttle Radar Topography Mission (SRTM) elevation model (<http://srtm.csi.cgiar.org>), to describe physiographic properties. To evaluate human impact on the Egyptian Vulture, we calculated the human footprint index using data on settlements, land transformation, accessibility and infrastructure (Sanderson *et al.* 2002). Access to food sources is among the primary parameters determining the species' distribution (Arrondo *et al.* 2018). This parameter was modelled based on livestock density in rural areas, using data provided by the Iranian Forests, Range and Watershed Management Organization (IFRWO), Iranian Department of Environment, and Statistical Centre of Iran. Correlated predictor pairs ($r > 0.7$) were excluded through correlation tests for all pairwise combinations of environmental variables for winter and summer visitors (Table 1).

Species distribution model

Using MaxEnt, an SDM for the Egyptian Vulture was developed (Phillips *et al.* 2006). MaxEnt is a machine learning statistical tool that matches or exceeds other modelling techniques in terms of performance (e.g. Phillips and Dudik 2008, Phillips *et al.* 2009). Previous research indicates MaxEnt's similar performance to regression-based techniques that analyse species' environmental

Table 1. Environmental predictors and their relative contributions to MaxEnt model.

| Summer visitors | Relative contributions (%) | Winter visitors | Relative contributions (%) |
|-------------------------------------|----------------------------|-------------------------------------|----------------------------|
| Climatic predictors | | | |
| Max temperature of warmest month | 1.11 | Min temperature of coldest month | 1.29 |
| Mean temperature of driest quarter | 2.27 | Mean temperature of wettest quarter | 1.11 |
| Mean temperature of warmest quarter | 1.04 | Mean temperature of coldest quarter | 0.88 |
| Precipitation of driest month | 1.06 | Precipitation of wettest month | 1.42 |
| Precipitation of driest quarter | 2.70 | Precipitation of wettest quarter | 1.00 |
| Precipitation of warmest quarter | 2.15 | Precipitation of coldest quarter | 0.92 |
| Vegetation predictors | | | |
| NDVI in spring | 17.00 | NDVI in autumn | 14.02 |
| NDVI in summer | 18.01 | NDVI in winter | 14.56 |
| Topographic predictors | | | |
| Mean of the elevation | 1.04 | Mean of the elevation | 12.70 |
| Standard deviation of the elevation | 16.02 | Standard deviation of the elevation | 2.00 |
| Mean of the slop | 0.08 | Mean of the slop | 1.04 |
| Standard deviation of the slop | 1.10 | Standard deviation of the slop | 0.22 |
| Land cover predictors | | | |
| Distance from stream | 1.14 | Distance from stream | 2.00 |
| Distance from river | 1.05 | Distance from river | 1.28 |
| Distance from lake | 0.02 | Distance from lake | 2.91 |
| Distance from bare area | 0.01 | Distance from bare area | 18.92 |
| Distance from rocky area | 2.02 | Distance from rocky area | 2.61 |
| Distance from protected area | 16.09 | Distance from protected area | 17.34 |
| Anthropogenic impact predictor | | | |
| Human footprint index | 14.08 | Human footprint index | 2.11 |
| Feeding predictor | | | |
| Livestock density | 2.01 | Livestock density | 1.02 |

The highest values are in bold.

correlates (e.g. Renner and Warton 2013, Merow and Silander 2014). MaxEnt can answer a host of ecological questions when models are appropriately calibrated and evaluated (Merow *et al.* 2013). We selected MaxEnt to develop predictive models for two reasons: (1) MaxEnt has performed robustly for small sample sizes and has outperformed other methods in such cases (Elith *et al.* 2006, Franklin 2010, Shcheglovitova and Anderson 2013); (2) owing to their lower sensitivity to over-prediction compared to standard GLM models, MaxEnt models are more useful to predict and extrapolate for conservation applications (Jackson *et al.* 2015, Zlonis *et al.* 2017).

In total, 156 presence points within Iran (residents = 6, summer visitors = 136 and winter visitors = 14) were used to train the model with default settings (regularization multiplier $\beta = 1$; auto features; convergence threshold = 0.00001) using the 'dismo' package (Hijmans *et al.* 2017) in R (R Core Team 2017). The model was run separately for summer and winter visitors. Presence points for residents and winter visitors were used in modelling the winter visitors. A similar process was used for summer visitors using presence points for the summer visitors and residents. Suitable habitats for the residents were determined by overlapping the suitable habitat of winter and summer visitors. One of the underlying assumptions of MaxEnt models is that occurrence data are representative of the species' distribution. However, this assumption is often inaccurate. To eliminate sample selection bias, uniform background data can be replaced with a random

sample of background data drawn from the sampling distribution (Phillips and Dudík 2008, Kramer-Schadt *et al.* 2013). Following this procedure, background data on the species were obtained from field data, literature review, and information resulting from interviews with wildlife experts, game wardens, hunters and taxidermists in 382 points for summer visitors and 108 points for winter visitors. Predictor contribution for the best fitting model was gauged via the jackknife procedure and permutation importance. MaxEnt provided a continuous map of habitat suitability. Additionally, the continuous map was converted into a binary presence-absence map by applying the probability threshold corresponding to each model's maximum Kappa value (Gutierrez *et al.* 2014). Subsequently, to assess the overlap between suitable habitats and protected areas, binary maps of habitat suitability and protected areas were compared in IDRISI Selva. In our study, 10 bootstrap replicates were run and the following two indices of model performance were used to evaluate the models by randomly assigning the presence records as training and test datasets (80% and 20%, respectively): (1) the area under the ROC-plot curve (AUC; Fielding and Bell 1997) in which 0 and 1 mean the worst and best prediction respectively, with 0.5 corresponding to a random prediction; and (2) the True Skill Statistics (TSS; Allouche *et al.* 2006) in which -1 and 1 mean the worst and best prediction respectively, with 0 corresponding to a random prediction. AUC and TSS are commonly used as presence-background evaluators (e.g. Descombes *et al.* 2016).

Results

AUC and TSS values indicated the good predictive performance of MaxEnt models for summer visitors (AUC: 0.91 ± 0.001 , TSS: 0.89 ± 0.001) and winter visitors (AUC: 0.86 ± 0.001 , TSS: 0.80 ± 0.002), with the highest AUC and TSS values for summer visitors.

Summer visitors, winter visitors and residents in this study showed a range of occurrence probabilities (Figures 1-3). Suitable habitats of summer visitors are scattered in the north and west of Iran and mainly concentrated along the Zagros and Alborz mountain ranges (Figure 2) which cover a large area of Iran. Suitable habitats of winter visitors and residents are scattered in coastal areas in the south (Figures 1 and 3).

MaxEnt results showed that 2%, 48%, and 3% of Iran can be considered as suitable habitats for residents, summer visitors, and winter visitors, respectively. Furthermore, the suitable habitats for residents, summer visitors, and winter visitors had only 4%, 9%, and 10% overlap with protected areas, respectively. This means that Egyptian Vulture mainly occurs outside protected areas in Iran (Figures 1-3). The results also indicated that large areas in Iran can be considered suitable habitat for summer visitors (Figure 2).

The MaxEnt model calculated the proportional contribution of each environmental variable for the winter and summer distribution and demonstrated various patterns (Table 1). The relative contribution of each variable is demonstrated in Table 1 for winter and summer visitors. Environmental variables that contributed the most to modelling the potential distribution of the Egyptian Vulture during summer and winter were seasonal NDVI, and distance from protected area. In contrast, climatic variables made only small contributions to model development (Table 1). Meanwhile, we found that some model variables have a high contribution to summer or winter visitors only. For example, mean elevation is important to winter visitors and standard deviation of the elevation is important to summer visitors.

Response curves revealed the direction of effect for the most important variables on habitat suitability of the Egyptian Vulture during summer and winter (Figure 4). The results demonstrated that increased seasonal NDVI and increased human footprint index caused a decrease in habitat suitability for Egyptian Vultures in both summer and winter. Also, increased distance from protected areas led to improvements in habitat suitability for winter visitors, while the same factor had the opposite effect for summer visitors. It was also concluded that distribution area of summer visitors was limited to altitudes between 500 and 2800 m, but habitat suitability for winter visitors increased with elevation.




Figure 1. Habitat suitability of winter visitors of Egyptian Vulture in Iran and gaps between suitable habitats and protected areas.

Discussion

According to the MaxEnt models, environmental variables influencing habitat suitability of the Egyptian Vulture partially change throughout the year. For example, for winter visitors mean elevation was particularly important in explaining habitat suitability while standard deviation of elevation had a more important role for summer visitors, replacing altitude.




Figure 2. Habitat suitability of summer visitors of Egyptian Vulture in Iran and gaps between suitable habitats and protected areas.

Our results revealed that only a small portion of suitable habitats during summer and winter is located in protected areas. In contrast, Mateo-Tomás *et al.* (2010) reported that a high percentage (65.7%) of the Egyptian Vultures' spring and summer territories are located within protected areas in the Cantabrian Mountains of Spain. It seems that protected areas may contain suitable habitats for the Egyptian Vulture (Carrete *et al.* 2007, 2009, Zuberogoitia *et al.* 2008, Mateo-Tomás *et al.* 2010). There are two possible reasons for the inadequate




Figure 3. Habitat suitability of residents of Egyptian Vulture in Iran and gaps between suitable habitats and protected areas.

protection of the species in Iran. Firstly, selection of protected areas is mainly based on the distribution of large mammals, and birds play only a small role. Secondly, birds are very mobile and have broad ecological niches, therefore a significant part of their suitable habitats is located outside protected area.

Our results also showed that increased distance from protected areas led to improvements in habitat suitability for winter visitors, while the same factor had the opposite effect for summer visitors. Zuberogoitia *et al.* (2014) explain that tourist activity in protected areas creates disturbance which leads to lower habitat suitability. In Iran, better security in protected areas (as a result of lower levels of human activity and disturbance and laws banning hunting) explains why summer visitors prefer to spend their breeding season inside protected areas.

As can be seen in our results, summer- and winter-visitor Egyptian Vultures are mainly found in areas with low vegetation cover, similar to findings by Margalida *et al.* (2007), stating that the species is partial to areas with sparse vegetation due to food being more easily obtained in such areas. This result is in agreement with previous studies which have supported the relationship between habitat suitability of the Egyptian Vulture and low vegetation cover (Carrete *et al.* 2007, Margalida *et al.* 2007). Despite the fact that this species' distribution is highly variable in terms of elevation at a global scale (IUCN 2017), Zuberogoitia *et al.* (2014) and Sen *et al.* (2017) said that elevation was one of the most important habitat variables for breeding populations of the Egyptian Vulture. Oppel *et al.* (2017) identified standard deviation of elevation as one of the most decisive environmental parameters influencing territory occupancy and breeding success of Egyptian Vultures in the Balkan Peninsula. We found that mean elevation makes a high contribution only for winter visitors. Egyptian Vultures during summer and winter react differently to mean elevation. Winter visitors show a positive trend with an increase in elevation; while summer visitors show a preference for middle elevations (500–2800 m). Mateo-Tomás and Olea (2009) account for the general negative trend between elevation and habitat suitability in terms of the extreme climatic conditions at high altitudes.




Figure 4. Response curves of Maxent model for Egyptian Vulture.

In step with the growing human disturbance, the spatial and temporal distribution, foraging behaviour and breeding success of wildlife has been changing (Fernández-Juricic and Tellería, 2000, Gill *et al.* 2001, Liberatori and Penteriani 2001, Bautista *et al.* 2004). Our findings linked human activity with habitat suitability for summer and winter visitors in Iran, in agreement with previous studies on the Egyptian Vulture (Zuberogoitia *et al.* 2008) as well as other species such as Bearded Vulture *Gypaetus barbatus* and Spanish Imperial Eagle *Aquila adalberti* (Arroyo and Razin 2006, González *et al.* 2006). However, some authors suggest that the Egyptian Vulture is tolerant of human activities (Ceballos and Donazar 1989). Our study showed a negative correlation between human disturbance (e.g. human settlements and land transformation) and habitat suitability for the Egyptian Vulture, i.e. areas far from infrastructure were the most suitable for the species. This observation can be explained by better security in such areas. Hunting and trafficking of birds of prey to other Middle Eastern countries are among the major threats to birds of prey in Iran. Despite the relative security of protected areas, since the majority of suitable Egyptian Vulture habitat is located outside protected areas; further human development can restrict its suitable habitats and drive the species to extinction in the near future.

Current biodiversity policy in the world is focused on halting the loss of biodiversity (IUCN 2017). Therefore, Egyptian Vulture during summer and winter must be protected to avoid regional loss of biodiversity and extinction of the species in Iran. There is no mechanism to coordinate the scientific community and environmental departments in Iran. Accordingly, we recommend that wildlife managers use this study to develop conservation strategies directed toward maintaining or ameliorating the quality of the species' potential habitat along its migration routes in the region.

Using MaxEnt in conservation-oriented studies has become common practice (Fourcade *et al.* 2018). Nevertheless, MaxEnt models have shortcomings such as: presence-only models underperform in predicting species' distributions compared to presence-absence models since absence data can help indicate unsuitable habitats for species, a process known as "fixing the floor". Thus, the importance of absence data is in eliminating low suitability areas that could have been categorised as suitable habitats if presence-only data had been used (Brotons *et al.* 2004, Lahoz-Monfort *et al.* 2014); and MaxEnt models can take advantage of museum and atlas data, which provide presence-only information. However, there are biases in data provided by museums and atlases towards accessible areas or areas where species are expected to occur (Kadmon *et al.* 2004, Pearce and Boyce, 2006). The data we have used as input face the same challenges. Rather than being collected with strategies such as random or systematic sampling, data in museums and atlases are gathered without particular sampling schemes, and without any particular purpose in mind (Elith *et al.* 2006). Biases in sampling intensity can therefore arise (Phillips *et al.* 2009, Leitão *et al.* 2011, Yackulic *et al.* 2013, Guillera-Arroita *et al.* 2015). As opposed to presence-absence models, presence-only models such as MaxEnt are strongly affected by such sampling biases.

References

- Allouche, O., Tsoar, A. and Kadmon, R. (2006) Assessing the accuracy of species distribution models: prevalence, kappa and the true skill statistic (TSS). *J. Appl. Ecol.* 43: 1223–1232.
- Angelov, I., Hashim, I. and Oppel, S. (2013) Persistent electrocution mortality of Egyptian Vultures *Neophron percnopterus* over 28 years in East Africa. *Bird Conserv. Internatn.* 23: 1–6.
- Arrondo, E., Moleón, M., Cortés-Avizanda, A., Jiménez, J., Beja, P., Sánchez-Zapata, J. A. and Donazar, J. A. (2018) Invisible barriers: Differential sanitary regulations constrain vulture movements across country borders. *Biol. Conserv.* 219: 46–52.
- Arroyo, B. and Razin, M. (2006) Effect of human activities on bearded vulture behaviour and breeding success in the French Pyrenees. *Biol. Conserv.* 128: 276–284.
- Austin, M. P. and Van Niel, K. P. (2011) Improving species distribution models for climate change studies: variable selection and scale. *J. Biogeogr.* 38: 1–8.

- Bautista, L. M., García, J. T., Calmaestra, R. G., Palacín, C., Martín, C. A., Morales, M. B. and Viñuela, J. (2004) Effect of weekend road traffic on the use of space by raptors. *Conserv. Biol.* 18: 726–732.
- Brottons, L., Thuiller, W., Araújo, M. B. and Hirzel, A. H. (2004) Presence-absence versus presence-only modelling methods for predicting bird habitat suitability. *Ecography* 27: 437–448.
- Carrete, M., Grande, J. M., Tella, J. L., Sánchez-Zapata, J. A., Donazar, J. A., Díaz-Delgado, R. and Romo, A. (2007) Habitat, human pressure, and social behavior: Partialling out factors affecting large-scale territory extinction in an endangered vulture. *Biol. Conserv.* 136: 143–154.
- Carrete, M., Sánchez-Zapata, J. A., Benítez, J. R., Lobón, M. and Donazar, J. A. (2009) Large scale risk-assessment of wind-farms on population viability of a globally endangered long-lived raptor. *Biol. Conserv.* 142: 2954–2961.
- Ceballos, O. and Donazar, J. A. (1989) Factors influencing the breeding density and nest-site selection of the Egyptian Vulture (*Neophron percnopterus*). *J. für Ornithol.* 130: 353–359.
- Cuthbert, R., Green, R. E., Ranade, S., Saravanan, S., Pain, D. J., Prakash, V. and Cunningham, A. A. (2006) Rapid population declines of Egyptian vulture (*Neophron percnopterus*) and red-headed vulture (*Sarcogyps calvus*) in India. *Anim. Conserv.* 9: 349–354.
- De Marco, P., Diniz-Filho, J. A. F. and Bini, L. M. (2008) Spatial analysis improves species distribution modelling during De Marco range expansion. *Biol. Lett.* 4: 577–580.
- Descombes, P., Petitpierre, B., Morard, E., Berthoud, M., Guisan, A. and Vittoz, P. (2016) Monitoring and distribution modelling of invasive species along riverine habitats at very high resolution. *Biol. Invasions* 18: 3665–3679.
- Elith, J. and Leathwick, J. R. (2009) Species distribution models: ecological explanation and prediction across space and time. *Annu. Rev. Ecol. Evol. Systemat.* 40: 677–697.
- Elith, J., Graham, C. H., Anderson, R. P., Dudík, M., Ferrier, S., Guisan, A., ... and Li, J. (2006) Novel methods improve prediction of species' distributions from occurrence data. *Ecography* 129–151.
- ESRI (2014) ArcGIS Desktop. Version 10.3. World Shaded Relief. Redlands, CA: ESRI.
- Farashi, A. and Shariati, M. (2017) Biodiversity hotspots and conservation gaps in Iran. *J. Nat. Conserv.* 39: 37–57.
- Fernández-Juricic, E. and Tellería, J. L. (2000) Effects of human disturbance on spatial and temporal feeding patterns of Blackbird *Turdus merula* in urban parks in Madrid, Spain. *Bird Study* 47: 13–21.
- Ferrer, M. and Penteriani, V. (2008) Non-independence of demographic parameters: positive density-dependent fecundity in eagles. *J. Appl. Ecol.* 45: 1453–1459.
- Fielding, A. H. and Bell, J. F. (1997) A review of methods for the assessment of prediction errors in conservation presence/absence models. *Environ. Conserv.* 24: 38–49.
- Flessner, B., Henry, M. C. and Green, J. (2017) Species distribution modeling of American beech (*Fagus grandifolia*) Distribution in Southwest Ohio. *Internatn. J. Appl. Geospatial Res.* 8 (3): 16–36.
- Fourcade, Y., Besnard, A. G. and Secondi, J. (2018) Paintings predict the distribution of species, or the challenge of selecting environmental predictors and evaluation statistics. *Glob. Ecol. Biogeogr.* 27: 245–256.
- Franklin, J. (2010) *Mapping species distributions: spatial inference and prediction*. Cambridge, UK: Cambridge University Press.
- Gill, J. A., Norris, K. and Sutherland, W. J. (2001) Why behavioural responses may not reflect the population consequences of human disturbance. *Biol. Conserv.* 97: 265–268.
- González, L. M., Arroyo, B. E., Margalida, A., Sánchez, R. and Oria, J. (2006) Effect of human activities on the behaviour of breeding Spanish imperial eagles (*Aquila adalberti*): management implications for the conservation of a threatened species. *Anim. Conserv.* 9: 85–93.
- Guillera-Arroita, G., Lahoz-Monfort, J. J., Elith, J., Gordon, A., Kujala, H., Lentini, P. E., ... and Wintle, B. A. (2015) Is my species distribution model fit for purpose? Matching data and models to applications. *Glob. Ecol. Biogeogr.* 24: 276–292.
- Guisan, A. and Thuiller, W. (2005) Predicting species distribution: offering more than

- simple habitat models. *Ecol. Lett.* 8: 993–1009.
- Gutiérrez, E. E., Boria, R. A. and Anderson, R. P. (2014) Can biotic interactions cause allopatry? Niche models, competition, and distributions of South American mouse opossums. *Ecography* 37: 741–753.
- Hernández, M. and Margalida, A. (2009) Poison-related mortality effects in the endangered Egyptian vulture (*Neophron percnopterus*) population in Spain. *Eur. J. Wildl. Res.* 55: 415–423.
- Hijmans, R. J., Cameron, S. E., Parra, J. L., Jones, P. G. and Jarvis, A. (2005) Very high resolution interpolated climate surfaces for global land areas. *Internatn. J. Climatol.* 25: 1965–1978.
- Hijmans, R. J., Phillips, S., Leathwick, J., Elith, J. and Hijmans, M. R. J. (2017) Package 'dismo'. *Circles* 9: 1.
- Iranian Department of Environment (2015) *The Fifth National Report to the Convention on Biological Diversity*. Tehran, Iran: Department of Environment of Iran.
- IUCN (2017) *The IUCN Red List of threatened species*. <http://www.iucnRedList.org/> Accessed 17.03.03.
- Jackson, M. M., Gergel, S. E. and Martin, K. (2015) Citizen science and field survey observations provide comparable results for mapping Vancouver Island White-tailed Ptarmigan (*Lagopus leucura saxatilis*) distributions. *Biol. Conserv.* 181: 162–172.
- Kaboli, M., Aliabadian, M., Tohidifar, M., Hashemi, A. and Roselaar, C. S. (2015) *Atlas of birds of Iran*. Tehran, Iran: Department of Environment of Iran.
- Kadmon, R., Farber, O. and Danin, A. (2004) Effect of roadside bias on the accuracy of predictive maps produced by bioclimatic models. *Ecol. Applic.* 14: 401–413.
- Kramer-Schadt, S., Niedballa, J., Pilgrim, J. D., Schröder, B., Lindenborn, J., Reinfelder, V., ... and Cheyne, S. M. (2013) The importance of correcting for sampling bias in MaxEnt species distribution models. *Divers. Distrib.* 19: 1366–1379.
- Lahoz-Monfort, J. J., Guillera-Aroita, G. and Wintle, B. A. (2014) Imperfect detection impacts the performance of species distribution models. *Glob. Ecol. Biogeogr.* 23: 504–515.
- Leitão, P. J., Moreira, F. and Osborne, P. E. (2011) Effects of geographical data sampling bias on habitat models of species distributions: a case study with steppe birds in southern Portugal. *Internatn. J. Geogr. Inf. Sci.* 25: 439–454.
- Liberatori, F. & Penteriani, V. (2001) A long-term analysis of the declining population of the Egyptian vulture in the Italian peninsula: distribution, habitat preference, productivity and conservation implications. *Biol. Conserv.* 101: 381–389.
- Loiselle, B. A., Howell, C. A., Graham, C. H., Goerck, J. M., Brooks, T., Smith, K. G. and Williams, P. H. (2003) Avoiding pitfalls of using species distribution models in conservation planning. *Conserv. Biol.* 17: 1591–1600.
- Margalida, A., García, D. and Cortés-Avizanda, A. (2007) Factors influencing the breeding density of bearded vultures, Egyptian vultures and Eurasian griffon vultures in Catalonia (NE Spain): management implications. *Anim. Biodivers. Conserv.* 30: 189–200.
- Mateo-Tomás, P. and Olea, P. P. (2009) Combining scales in habitat models to improve conservation planning in an endangered vulture. *Acta Oecologica*, 35(4), 489–498.
- Mateo-Tomás, P., Olea, P. P. and Fombellida, I. (2010) Status of the Endangered Egyptian vulture *Neophron percnopterus* in the Cantabrian Mountains, Spain, and assessment of threats. *Oryx* 44: 434–440.
- Maxwell, S. L., Fuller, R. A., Brooks, T. M., and Watson, J. E. (2016) Biodiversity: The ravages of guns, nets and bulldozers. *Nature* 536: 143–145.
- Merow, C. and Silander, J. A. (2014) A comparison of Maxlike and Maxent for modeling species distributions. *Meth. Ecol. Evol.* 5: 215–225.
- Merow, C., Smith, M. J. and Silander, J. A. (2013) A practical guide to MaxEnt for modeling species' distributions: what it does, and why inputs and settings matter. *Ecography* 36: 1058–1069.
- Morán-Ordóñez, A., Lahoz-Monfort, J. J., Elith, J. and Wintle, B. A. (2017) Evaluating 318 continental-scale species distribution models over a 60-year prediction horizon: what factors influence the reliability of predictions? *Glob. Ecol. Biogeogr.* 26: 371–384.

- Naveda-Rodríguez, A. (2013) *Biogeography and conservation status of diurnal raptors in Venezuela*. Doctoral dissertation, MS thesis, Universidad Internacional Menéndez Pelayo, Madrid, Spain.
- Naveda-Rodríguez, A. (2015) Conservation status of diurnal raptors in Venezuela. *J. Raptor Res.* 49: 441–449.
- Nikolaus, G. (1984) Large numbers of birds killed by electric power line. *Scopus* 8: 42.
- Nikolaus, G. (2006) Where have the African vultures gone. *Vulture News* 55: 65–67.
- Ogada, D., Shaw, P., Beyers, R. L., Buij, R., Murn, C., Thiollay, J. M., ... and Krüger, S. C. (2016) Another continental vulture crisis: Africa's vultures collapsing toward extinction. *Conserv. Lett.* 9: 89–97.
- Oppel, S., Dobrev, V., Arkumarev, V., Saravia, V., Bounas, A., Manolopoulos, A., ... and Nikolov, S. C. (2017) Landscape factors affecting territory occupancy and breeding success of Egyptian Vultures on the Balkan Peninsula. *J. Ornithol.* 158: 443–457.
- Parmesan, C. (2006) Ecological and evolutionary responses to recent climate change. *Annu. Rev. Ecol. Evol. Syst.* 37: 637–669.
- Parolo, G., Rossi, G. and Ferrarini, A. (2008) Toward improved species niche modelling: *Arnica montana* in the Alps as a case study. *J. Appl. Ecol.* 45: 1410–1418.
- Pearce, J. L. and Boyce, M. S. (2006) Modelling distribution and abundance with presence-only data. *J. Appl. Ecol.* 43: 405–412.
- Penteriani, V., Ferrer, M. and Delgado, M. D. M. (2011) Floater strategies and dynamics in birds, and their importance in conservation biology: towards an understanding of nonbreeders in avian populations. *Anim. Conserv.* 14: 233–241.
- Phillips, S. J. and Dudík, M. (2008) Modeling of species distributions with Maxent: new extensions and a comprehensive evaluation. *Ecography* 31: 161–175.
- Phillips, S. J., Anderson, R. P. and Schapire, R. E. (2006) Maximum entropy modelling of species geographic distributions. *Ecol. Modell.* 190: 231–259.
- Phillips, S. J., Dudík, M., Elith, J., Graham, C. H., Lehmann, A., Leathwick, J. and Ferrier, S. (2009) Sample selection bias and presence-only distribution models: implications for background and pseudo-absence data. *Ecol. Applic.* 19: 181–197.
- R Core Team (2017) *R: a language and environment for statistical computing*. Vienna, Austria: R Foundation for Statistical Computing.
- Ramesh, V., Gopalakrishna, T., Barve, S. and Melnick, D. J. (2017) Citizen science driven species distribution models estimate drastically smaller range sizes and higher threat levels for Western Ghats endemic birds. *Biol. Conserv.* 210: 205–221.
- Rangel, T. F. L., Diniz-Filho, J. A. F., and Bini, L. M. (2006) Towards an integrated computational tool for spatial analysis in macroecology and biogeography. *Global Ecol. Biogeog.* 15: 321–327.
- Rempel, R. S. and Hornseth, M. L. (2017) Binational climate change vulnerability assessment of migratory birds in the Great Lakes Basins: Tools and impediments. *PLoS One* 12(2): e0172668.
- Renner, I. W. and Warton, D. I. (2013) Equivalence of MAXENT and Poisson point process models for species distribution modeling in ecology. *Biometrics* 69: 274–281.
- Riordan, E. C. and Rundel, P. W. (2014) Land use compounds habitat losses under projected climate change in a threatened California ecosystem. *PLoS One* 9(1): e86487.
- Rodrigues, A. S., Andelman, S. J., Bakarr, M. I., Boitani, L., Brooks, T. M., Cowling, R. M., ... and Long, J. S. (2004) Effectiveness of the global protected area network in representing species diversity. *Nature* 428(6983): 640–643.
- Sala, O. E., Chapin, F. S., Armesto, J. J., Berlow, E., Bloomfield, J., Dirzo, R., ... and Leemans, R. (2000) Global biodiversity scenarios for the year 2100. *Science* 287(5459): 1770–1774.
- Sanderson, E. W., Jaiteh, M., Levy, M. A., Redford, K. H., Wannebo, A. V. and Woolmer, G. (2002) The human footprint and the last of the wild. *BioScience* 52: 891–904.
- Sen, B., Tavares, J. and Bilgin, C. (2017) Nest site selection patterns of a local Egyptian Vulture *Neophron percnopterus* population in Turkey. *Bird Conserv. Internatn.* 27: 568–581.
- Serrano-Rodríguez, A., Escalona-Segura, G., Elias, E. E. I., Rodríguez, A. S., Uriostegui,

- J. M. and Aguilar, A. C. M. D. O. (2018) Potential distribution and climatic niche of seven species of *Campylorhynchus* (Aves, Troglodytidae): conservation implications for *C. yucatanicus*. *Wilson J. Ornithol.* 130: 3–22.
- Shcheglovitova, M. and Anderson, R. P. (2013) Estimating optimal complexity for ecological niche models: a jackknife approach for species with small sample sizes. *Ecol. Model.* 269: 9–17.
- Stralberg, D., Jongsomjit, D., Howell, C. A., Snyder, M. A., Alexander, J. D., Wiens, J. A. and Root, T. L. (2009) Re-shuffling of species with climate disruption: a no-analog future for California birds? *PLoS One* 4(9): e6825.
- Thiollay, J. M. (2006a) The decline of raptors in West Africa: long-term assessment and the role of protected areas. *Ibis* 148: 240–254.
- Thiollay, J. M. (2006b) Severe decline of large birds in the Northern Sahel of West Africa: a long-term assessment. *Bird Conserv. Internatn.* 16: 353–365.
- Thuiller, W., Lavorel, S., Araújo, M. B., Sykes, M. T. and Prentice, I. C. (2005) Climate change threats to plant diversity in Europe. *Proc. Natl. Acad. Sci. USA* 102: 8245–8250.
- Velevski, M., Nikolov, S. C., Hallmann, B., Dobrev, V., Sidiropoulos, L., Saravia, V., ... and Stara, K. (2015) Population decline and range contraction of the Egyptian Vulture *Neophron percnopterus* in the Balkan Peninsula. *Bird Conserv. Internatn.* 25: 440–450.
- Venter, O., Magrath, A., Outram, N., Klein, C. J., Possingham, H. P., Di Marco, M., and Watson, J. E. (2018) Bias in protected-area location and its effects on long-term aspirations of biodiversity conventions. *Conserv. Biol.* 32: 127–134.
- Virkkala, R., Heikkinen, R. K., Fronzek, S. and Leikola, N. (2013) Climate change, northern birds of conservation concern and matching the hotspots of habitat suitability with the reserve network. *PLoS One* 8(5): e63376.
- Yackulic, C. B., Chandler, R., Zipkin, E. F., Royle, J. A., Nichols, J. D., Campbell Grant, E. H., and Veran, S. (2013) Presence-only modelling using MAXENT: when can we trust the inferences? *Methods Ecol. Evol.* 4: 236–243.
- Zlonis, E., Panci, H., Bednar, J., Hamady, M. and Niemi, G. (2017) Habitats and landscapes associated with bird species in a lowland conifer-dominated ecosystem. *Avian Conserv. Ecol.* 12(1): 7. <https://doi.org/10.5751/ACE-00954-120107>.
- Zuberogoitia, I., Zabala, J., Martínez, J. A., Martínez, J. E., and Azkona, A. (2008) Effect of human activities on Egyptian vulture breeding success. *Anim. Conserv.* 11: 313–320.
- Zuberogoitia, I., Zabala, J., Martínez, J. E., González-Oreja, J. A. and López-López, P. (2014) Effective conservation measures to mitigate the impact of human disturbances on the endangered Egyptian vulture. *Anim. Conserv.* 17: 410–418.

AZITA FARASHI*, MOHAMMAD ALIZADEH-NOUGHANI

Department of Environmental Sciences, Faculty of Natural Resource and Environment,
Ferdowsi University of Mashhad, Iran.

*Author for correspondence; e-mail: farashi@um.ac.ir

Received 27 January 2018; revision accepted 31 July 2018;
Published online 29 November 2018