

Research Article

Cite this article: Hasanfard A and Nabati J (2025). Environmental factors influencing germination characteristics of sterile oat (*Avena sterilis*) populations from northeast and northwest Iran. *Weed Sci.* **73**(e47), 1–8. doi: [10.1017/wsc.2025.10020](https://doi.org/10.1017/wsc.2025.10020)

Received: 24 February 2025

Revised: 24 April 2025

Accepted: 25 April 2025

Associate Editor:

William Vencill, University of Georgia

Keywords:

Environmental stress; germination; local adaptation

Corresponding author:

Alireza Hasanfard;

Emails: alireza.hasanfard@yahoo.com;

alireza.hasanfard@mail.um.ac.ir

Environmental factors influencing germination characteristics of sterile oat (*Avena sterilis*) populations from northeast and northwest Iran

Alireza Hasanfard¹  and Jafar Nabati²

¹Research Scholar, Weed Science Division, Department of Agrotechnology, Faculty of Agriculture, Ferdowsi University of Mashhad, Mashhad, Iran and ²Assistant Professor in Crop Physiology, Department of Agrotechnology, Faculty of Agriculture, Ferdowsi University of Mashhad, Mashhad, Iran

Abstract

Sterile oat [*Avena sterilis* L. ssp. *ludoviciana* (Durieu) Gillet & Magne] is rapidly proliferating in cereal fields across northeastern and northwestern Iran, underscoring the necessity of studying its ecology in these two distinct climates. A study was conducted to assess the impact of environmental factors on the germination of two native populations of *A. sterilis*. The germination responses of populations from Mashhad (northeastern Iran) and Tabriz (northwestern Iran) were evaluated under various treatments, including temperature, osmotic potential, NaCl concentration, and light/dark cycles. As the temperature increased and osmotic potential decreased—indicating heightened drought stress—the germination percentages of both populations declined. The Mashhad population exhibited the highest germination percentages, reaching 99% at 10 C and 100% at 15 C, both under 0 MPa osmotic potential. Conversely, the Tabriz population demonstrated its peak germination percentages at 15 C and 20 C, also under 0 MPa osmotic potential, with rates of 97% and 96%, respectively. The highest germination rates were observed in seeds from the Mashhad and Tabriz populations at 15 C, with osmotic potentials of 0 MPa and −0.3 MPa, yielding rates of 0.52 and 0.48 seeds d^{−1}, respectively. The NaCl concentration required for 50% inhibition of seed germination was 4.76 dS m^{−1} for the Mashhad population and 3.90 dS m^{−1} for the Tabriz population. In both populations, the highest germination percentage was observed under a light/dark cycle of 10 h of light and 14 h of darkness. The differences in germination responses between the Mashhad and Tabriz populations can be attributed to local environmental adaptations. Variations in temperature, osmotic potential, and other climatic factors influence seed dormancy and germination traits, enabling populations to thrive in their specific habitats. These local adaptations contribute to differences in germination performance under various environmental conditions, ultimately affecting their potential spread across different regions.

Introduction

Sterile oat [*Avena sterilis* L. ssp. *ludoviciana* (Durieu) Gillet & Magne], a member of the Poaceae family, is a fast-growing annual grass commonly found in disturbed environments such as agricultural fields, roadsides, and grasslands (Loskutov et al. 2023). Like other weeds in the Poaceae family, *A. sterilis* is a strong competitor for water, nutrients, and light due to its dense leaf cover; robust tillering ability; tall growth habit; allelopathic properties; and ecological, biological, and botanical similarities to cereals (Hasanfard et al. 2022). This species can tolerate high temperatures and arid conditions, making it a formidable competitor in hot, dry environments. In winter wheat (*Triticum aestivum* L.) fields across Iran, *A. sterilis* is considered a problematic weed that competes with wheat for essential resources, thereby reducing crop yields—particularly in regions with limited irrigation (Hasanfard et al. 2022; Nosratti et al. 2020). Armin and Asghripour (2011) demonstrated that varying densities of *A. sterilis* significantly reduced wheat yields in Iran. For instance, densities of 25, 50, and 75 plants m^{−2} resulted in yield reductions of 10%, 19%, and 20%, respectively. The species' ability to withstand both drought and high temperatures further complicates weed management in these agricultural systems. Moreover, its capacity for rapid germination and growth enables *A. sterilis* to outcompete wheat, especially under resource-limited conditions, exacerbating the challenges faced by farmers.

Understanding the germination behavior of weed seeds in response to temperature and drought is essential for developing effective weed management strategies in both agricultural and natural ecosystems (Anwar et al. 2021; Toscano et al. 2017). Temperature and moisture are critical environmental factors that influence seed germination (Dürr et al. 2015). Different weed species exhibit varying levels of sensitivity to these conditions; some may require specific temperature ranges to initiate germination, while others may be more resilient to drought stress.

© The Author(s), 2025. Published by Cambridge University Press on behalf of Weed Science Society of America. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution and reproduction, provided the original article is properly cited.



By exploring how these factors interact, land managers can devise targeted approaches to control weed populations, thereby minimizing their impact on crop yields and native plant communities. This knowledge enables timely interventions, such as adjusting planting schedules or employing specific irrigation techniques to reduce competition from weeds. Furthermore, understanding the dynamics of seed germination can help predict weed emergence patterns, allowing for proactive management before weed populations become problematic (Ramesh 2015). Overall, integrating this knowledge into management practices not only enhances agricultural productivity but also supports biodiversity conservation in natural ecosystems.

The geographic origin of weed seeds such as *A. sterilis* significantly influences their germination characteristics, including germination percentage and rate. Seeds from different regions may adapt to local environmental conditions, such as temperature, moisture, and seasonal variations. For instance, seeds from colder climates might exhibit a longer dormancy period to prevent premature germination, while those from warmer or more stable climates may germinate more quickly and uniformly. In a study assessing nine common weed species through temperature gradient experiments, it was observed that summer annuals exhibited higher base temperatures for germination compared with winter annuals (Steinmaus et al. 2000).

Salinity is one of the major abiotic stresses that adversely affects seed germination by reducing water uptake, lowering osmotic potential, and causing ion toxicity. These effects can lead to a decline in both the rate and percentage of germination through cellular damage and metabolic disruption. Amini et al. (2024) found that increasing NaCl concentration significantly reduced the germination percentage of *Cichorium glandulosum* Boiss. et Huet seeds in both the Tabriz and Marand populations. In Tabriz, germination remained around 90% up to 2 dS m⁻¹ but decreased as salinity levels exceeded this threshold. In contrast, Marand seeds exhibited 95% germination under nonsaline conditions, with a marked decline when salinity surpassed 1 dS m⁻¹. Both populations demonstrated nearly zero germination at 10 and 12 dS m⁻¹. Factors such as soil type, photoperiod, and rainfall patterns can shape the germination traits of *A. sterilis*, leading to variations in seed viability and overall germination behavior based on their geographic origin. Therefore, examining the germination responses of *A. sterilis* populations from different regions can help identify how these populations may vary under diverse environmental conditions and predict their germination patterns. The primary objective of this study is to examine the germination responses of two *A. sterilis* populations originating from different geographic regions (Mashhad and Tabriz) under various environmental conditions, including temperature, humidity, salinity, and light requirement. By testing the hypothesis that both environmental factors and regional origin significantly influence germination behavior, this research aims to provide insights into the role of local adaptation in seed responses to abiotic stressors.

Materials and Methods

Study Design, Seed Population, and Climate

This study was conducted twice, in 2020 and 2021, at the greenhouse research laboratory of the Faculty of Agriculture at Ferdowsi University of Mashhad in Iran. It was designed as a factorial experiment utilizing a completely randomized design with six replications. The study employed two populations of *A. sterilis*

Table 1. Geographic locations and climatic characteristics of seed collection origins.

Feature	Origin of seeds	
	Mashhad	Tabriz
Location	Northeastern Iran, Khorasan Razavi Province	Northwest Iran, East Azerbaijan Province
Coordinates	36.32°N, 59.61°E	38.09°N, 46.29°E
Elevation	1,050 m (3,445 feet) above sea level	1,350 m (4,430 feet) above sea level
Topography	Foothills of the Torghabeh and Binalood Mountains	Semi-mountainous, foothills of the Sahand Mountains
Climate	Hot desert (Köppen BWh)	Cold semiarid (Köppen BSk)
Average annual temperature	14–16 C	10–12 C
Summer temperature	Hot summers	Hot, dry summers
Winter temperature	Cold winters, below freezing	Cold winters, below freezing
Annual precipitation	Around 250 mm	Around 300 mm

collected from the margins of wheat fields in northeastern Iran (Mashhad) and northwestern Iran (Tabriz) (Table 1). Mashhad, located in Razavi Khorasan, has a semiarid climate with hot summers and cold winters, receiving approximately 250 mm of rainfall annually. This region is known for cultivating wheat, barley (*Hordeum vulgare* L.), and saffron (*Crocus sativus* L.). Due to the cold and dry winters, wild oats (*Avena sterilis*), a winter weed, particularly thrives in cereal fields, such as wheat and barley, in this area. The autumn and winter rainfall in Mashhad, combined with suitable temperatures, provides ideal conditions for the growth and propagation of *A. sterilis*. This weed, which can grow well under drought and cold winter conditions, can pose strong competition to agricultural crops, negatively affecting wheat and barley yields. In contrast, Tabriz, in East Azerbaijan Province, has a continental climate with cold winters and hot summers, receiving about 300 mm of rainfall annually. While the autumn and winter rainfall in Tabriz is higher than in Mashhad, the severe cold and specific climatic conditions prevent *A. sterilis* from spreading as extensively as in Mashhad, resulting in a more limited competition with agricultural products. Given that the interaction between temperature and drought is particularly common, especially in Iran's climate (Rastgoo and Hasanfard 2021), we specifically focused on the interaction between these two factors.

Seed Preparation and Dormancy-breaking Treatment

In this study, the preparation of *A. sterilis* seeds involved the careful removal of lemma and palea, followed by the elimination of empty and damaged seeds. To break seed dormancy, a stratification treatment was employed through a moist-chilling process. Over a period of 3 d, the seeds were subjected to a constant temperature of 5 C in a dark environment, effectively mimicking their natural winter conditions. This treatment successfully facilitated germination and ensured optimal growth in the subsequent stages.

Seed Disinfection and Germination

For the experimental setup, 10 viable seeds were selected and placed in 9-cm-diameter petri dishes lined with Whatman No. 1

Filter Paper. Before the experiments, seeds were disinfected using a 5% sodium hypochlorite solution for 3 to 5 min at room temperature, followed by thorough rinsing with distilled water. Each petri dish was supplied with 10 ml of a prepared solution corresponding to the designated drought level (Rastgoo et al. 2023). To maintain precise temperature control, the germinator was pre-adjusted to the required temperature settings 1 d before the initiation of the thermal cycle. Once the temperature fluctuations were confirmed to be within ± 1 C, the petri dishes were transferred into the germinator (Grouc Engineering Design, Tehran, Iran), ensuring exposure to the desired temperature in dark conditions throughout the experiment.

Temperature and Osmotic Potential Levels

The experiment was conducted separately for each of the factors: temperature by osmotic potential, salinity, and light/dark regimes. The selected levels of temperature, osmotic potential, and salinity were determined based on regional climate characteristics and soil conditions, whereas the light/dark regimes were informed by the general daylength patterns observed in the native habitats of the studied populations. In this study, we focused on the interaction between temperature and osmotic potential, as these two factors play a fundamental role in seed germination, particularly under conditions of environmental stress. Given their strong interdependence, a more detailed analysis was necessary to understand their combined effects on the germination of *A. sterilis*. The experimental treatments consisted of six temperature levels (5, 10, 15, 20, 25, and 30 C) and five levels of osmotic potential, which were established using polyethylene glycol 6000 (PEG 6000) (Rastgoo et al. 2023). PEG is widely used to create controlled osmotic potential in experiments, as it effectively reduces water availability in a solution without interfering with the plants' nutrient uptake. The osmotic potential levels were set at 0, -0.3, -0.6, -0.9, and -1.2 MPa by dissolving varying concentrations of PEG in the solution.

Salinity Stress

Sodium chloride (NaCl) solutions with salinity concentrations of 0 (control), 1, 2, 3, 4, 5, 6, 8, 10, and 12 dS m⁻¹ were prepared to establish 10 distinct levels of salinity stress (Poljakoff-Mayber et al. 1994).

Light/Dark Regimes

The effect of various light/dark cycles, including 24/0, 12/12, 10/14, 8/16, and 0/24 h at 20/10 C, on the germination of *A. sterilis* seeds was monitored daily for 18 d following treatment. The light intensity of 156 $\mu\text{mol m}^{-2} \text{s}^{-1}$ was supplied by fluorescent lamps in a growth chamber. To assess germination in the absence of light, the petri dishes were covered with two layers of aluminum foil (Baskin and Baskin 1998).

Calculations and Statistical Analysis

The counting of germinated seeds continued for 23 d, until the number of germinating seeds became constant. Subsequently, the percentage and rate of germination were calculated using Equations 1 and 2. Additionally, the time needed to achieve 50% germination was determined by fitting a logistic model of cumulative germination percentage over time. Statistical analysis of the data was conducted using SAS v. 9.4 (SAS Institute, Cary, NC, USA) and GraphPad Prism software (version 8.0.1, La Jolla, CA, USA).

$$GP = (n/N) \times 100 \quad [1]$$

where GP represents the percentage of germination, n stands for the number of germinated seeds, and N denotes the total number of seeds.

$$R_{50} = 1/D_{50} \quad [2]$$

In this equation, R_{50} represents the germination rate, and D_{50} indicates the time needed to achieve 50% of the final germination.

Cumulative germination graphs, along with parameters for maximum germination and the time required to reach 50% germination at various temperatures and osmotic potentials, were obtained through nonlinear regression analysis using a three-parameter log-logistic equation (Equation 3).

$$Y = \frac{d}{1 + e^{b(\log x - \log e)}} \quad [3]$$

where Y represents the response rate (cumulative germination) at time x , d denotes the upper limit for Y , b indicates the slope of the curve at point e , and e is the time required to achieve 50% germination (in days).

For germination data analysis, an ANOVA was performed using SAS v. 9.4. Differences between treatments (temperature, osmotic potential, salinity, and light/dark regimes) were assessed using the standard error of the mean (SEM) values. Data from both runs were pooled, as no significant differences were observed between runs ($P > 0.05$). Cumulative germination graphs were generated using GraphPad Prism software (v. 8.0.1, La Jolla, CA, USA).

Results and Discussion

Temperature and Osmotic Potential

Cumulative Germination

The interaction among population, temperature, and osmotic potential significantly affected the germination percentage ($P < 0.01$). As temperature increased and osmotic potential decreased (indicating increased drought stress), the germination percentage of both populations decreased (Figures 1 and 2). The highest germination percentages (99% and 100%, respectively) were observed in the Mashhad population at 10 and 15 C and 0 MPa osmotic potential (Figure 1; Table 2). The Tabriz population exhibited the highest germination percentages (97% and 96%, respectively) at 15 and 20 C and 0 MPa osmotic potential (Figure 2; Table 3).

The observed decline in germination percentages as drought stress and temperatures increased highlights the significant impact of abiotic stress on seed germination, an essential phase in the life cycle of plants. Both drought stress and elevated temperatures are known to negatively affect seed viability and germination rates, with water scarcity hindering seed hydration and temperature extremes potentially disrupting enzymatic activity and cellular functions necessary for germination (Abdellaoui et al. 2019; Bewley et al. 2013).

The comparison between the Mashhad and Tabriz populations suggests that these populations have different adaptive strategies or tolerance levels to these abiotic stress factors. The Mashhad population exhibited the highest germination percentage under optimal conditions of 0 MPa osmotic potential and cooler




Figure 1. Cumulative germination of *Avena sterilis* in the Mashhad population over 23 d in response to temperature and osmotic potential. The vertical bars represent the standard error (\pm SE). The parameter estimates for the model are presented in Table 2.

temperatures. This indicates that the Mashhad population might be adapted to a more temperate or moisture-rich environment, where lower temperatures and higher water availability are common. Conversely, the Tabriz population exhibited a similar decline in germination as drought stress and temperature increased. However, the Tabriz population occasionally demonstrated the ability to tolerate slightly higher temperatures, suggesting that the Tabriz population may be better adapted to a warmer, potentially more arid environment.

This population-level variation in germination traits likely reflects local adaptation to contrasting environmental conditions. In Mashhad, cooler temperatures and more consistent moisture availability may have selected for traits that enhance germination under lower temperature and higher water conditions. In contrast, the Tabriz population may have experienced stronger selective pressures related to heat and water scarcity, possibly including episodic drought and increased salinity, which are common in

semiarid and arid regions. These environmental factors may have driven the evolution of increased thermal tolerance or more conservative germination strategies in this population. These findings imply that local environmental conditions such as temperature and water availability may drive the evolution of specific stress-tolerance mechanisms in plant populations. The Mashhad population might have evolved mechanisms that prioritize survival in cooler, wetter climates, while the Tabriz population could be better suited to withstand the higher temperatures and potentially drier conditions typical of its native environment. Thus, the observed differentiation in germination behavior may be a product of ecological divergence driven by climate variability, water availability, and, potentially, soil salinity. In various regions, distinct microclimates may exist, each influencing environmental effects (Wong et al. 2016). This suggests that the average temperature and humidity conditions in cities such as Tabriz or Mashhad may vary across specific areas




Figure 2. Cumulative germination of *Avena sterilis* in the Tabriz population over 23 d in response to temperature and osmotic potential. The vertical bars represent the standard error (\pm SE). The parameter estimates for the model are presented in Table 3.

within these regions. For instance, while Tabriz is generally cooler and more humid, some parts of the region might experience higher temperatures and increased dryness during the summer, to which the seeds of the Tabriz population may have specifically adapted. Future studies could further explore these patterns by linking environmental data from the native habitats of each population with trait-based responses in controlled experiments.

Such variations in stress tolerance are important for understanding how populations might respond to climate change, which is expected to intensify both drought conditions and temperature extremes. They also emphasize the importance of considering population-specific responses to environmental stress when designing conservation strategies or when assessing the ecological resilience of species in changing climates. The study by Saburi-Rad et al. (2012) demonstrated that the highest germination percentage in *kochia* [*Bassia scoparia* (L.) A.J. Scott] seeds occurred at temperatures between 20 and 25 C and at osmotic potentials of -0.25 , -0.50 , -0.75 , and -1 MPa.

Time Required to Reach 50% Germination

The shortest time required to achieve 50% germination was observed in the Mashhad population at 15 C and 0 MPa osmotic potential (2 d) (Figure 1; Table 2), as well as in the Tabriz population at 15 C and -0.3 MPa osmotic potential (2 d) (Figure 2; Table 3). According to Figure 1, the time required to achieve 50% germination in the Mashhad population at 5 C ranged from 9 to 13 d, which was the longest duration compared with other temperatures. In the Tabriz population at 25 C, this duration ranged from 5 to 11 d (Figure 2).

The geographic distribution of the Mashhad and Tabriz populations is closely linked to their germination characteristics, particularly their responses to water availability. The Tabriz population, which shows rapid germination at slightly negative osmotic potentials (-0.3 MPa), may be better suited to drier, more arid regions with limited water resources. This ability to tolerate low water availability could increase its geographic spread in such environments. On the other hand, the Mashhad population, which

Table 2. Nonlinear regression analysis for water potential and temperature in the *Avena sterilis* population of Mashhad^a.

Temperature	Osmotic potential	Maximum germination	<i>b</i>	<i>X</i> ₅₀	<i>R</i> ²
C	MPa	%		days	
5	0	57	0.10	12	0.99
	−0.3	30	0.21	10	0.98
	−0.6	28	0.15	10	0.99
	−0.9	26	0.16	9	0.98
	−1.2	26	0.25	13	0.99
10	0	99	0.12	3	0.97
	−0.3	85	0.07	4	0.98
	−0.6	59	0.15	3	0.99
	−0.9	35	0.10	5	0.98
	−1.2	10	0.42	11	0.95
15	0	100	0.61	2	0.99
	−0.3	88	0.11	3	0.97
	−0.6	67	0.15	3	0.90
	−0.9	61	0.06	4	0.93
	−1.2	35	0.32	6	0.99
20	0	92	0.12	3	0.95
	−0.3	80	0.14	3	0.99
	−0.6	52	0.42	4	0.95
	−0.9	31	0.06	5	0.84
	−1.2	23	0.15	10	0.99
25	0	92	0.06	3	0.92
	−0.3	66	0.07	7	0.98
	−0.6	55	0.17	5	0.99
	−0.9	21	0.08	7	0.88
	−1.2	20	0.46	13	0.98
30	0	81	0.07	4	0.97
	−0.3	65	0.10	4	0.95
	−0.6	35	0.19	7	0.99
	−0.9	18	0.25	9	0.98
	−1.2	14	0.82	13	0.99

^aAbbreviations: *b*, slope; *X*₅₀, the time required to reach 50% germination; *R*², the coefficient of determination.

germinates efficiently at 0 MPa, may thrive in regions with intermittent or seasonal droughts. The differing drought-tolerance strategies of these populations highlight how local environmental conditions influence seed germination, survival, and, ultimately, the distribution of these populations. The ability to withstand dry conditions could allow these populations to colonize a wider range of habitats, driving their geographic expansion.

Germination Rate

The interaction among population, osmotic potential, and temperature significantly affected the germination rate (*P* < 0.05). The highest germination rate was observed in the seeds of Mashhad and Tabriz populations at 15 C and at osmotic potentials of 0 MPa and −0.3 MPa, respectively (0.52 and 0.48 seeds d^{−1}, respectively) (Table 4).

These findings suggest that the Mashhad population may be more sensitive to drought stress during germination than the Tabriz population. In a similar study, the Turkish red pine (*Pinus brutia* Ten.) provenances (Burdur/Bucak) exhibited the highest germination rate under an osmotic potential of −0.2 MPa (Koç 2021).

Salt Stress

The germination rate was significantly influenced by the interaction between population and NaCl concentration (*P* < 0.01). Increasing the NaCl concentration decreased the germination percentage of *A.*

Table 3. Nonlinear regression analysis for water potential and temperature in the *Avena sterilis* population of Tabriz^a.

Temperature	Osmotic potential	Maximum germination	<i>b</i>	<i>X</i> ₅₀	<i>R</i> ²
C	MPa	%		days	
5	0	55	0.10	4	0.98
	−0.3	36	0.08	4	0.99
	−0.6	26	0.12	10	0.95
	−0.9	20	0.30	8	0.99
	−1.2	17	1.19	8	0.99
10	0	90	0.05	4	0.95
	−0.3	58	0.10	4	0.99
	−0.6	52	0.09	4	0.97
	−0.9	36	0.14	4	0.97
	−1.2	33	0.20	9	0.99
15	0	97	0.16	3	0.97
	−0.3	70	0.15	2	0.89
	−0.6	51	0.34	4	0.97
	−0.9	44	0.14	6	0.99
	−1.2	35	0.13	9	0.95
20	0	96	0.07	5	0.99
	−0.3	85	0.11	3	0.99
	−0.6	43	0.17	3	0.96
	−0.9	37	0.12	5	0.96
	−1.2	13	1.22	6	0.99
25	0	85	0.08	8	0.99
	−0.3	81	0.11	6	0.99
	−0.6	48	0.09	5	0.98
	−0.9	20	0.42	8	0.99
	−1.2	14	1.17	11	0.98
30	0	84	0.08	6	0.99
	−0.3	78	0.07	5	0.98
	−0.6	44	0.08	5	0.97
	−0.9	18	0.26	9	0.98
	−1.2 ^b	—	—	—	—

^aAbbreviations: *b*, slope; *X*₅₀, the time required to reach 50% germination; *R*², the coefficient of determination.


^bLack of proper fit.

sterilis seeds in both the Mashhad and Tabriz populations (Figure 3). The rate of germination reduction was steeper for the Tabriz population (Hill slope = −0.36) than for the Mashhad population (Hill slope = −0.34). At all NaCl concentrations, the germination rates in the Mashhad population were consistently higher than those in the Tabriz population. The NaCl concentration required for 50% inhibition of seed germination was 4.76 dS m^{−1} for the Mashhad population and 3.90 dS m^{−1} for the Tabriz population (Figure 3).

The observed decrease in germination percentage with increasing NaCl concentration aligns with previous studies highlighting the detrimental effects of salinity on seed germination (Nikolić et al. 2023). However, the more pronounced reduction in germination rates for the Tabriz population suggests a differential tolerance to salinity between the two populations. The steeper slope observed in the Tabriz population indicates a heightened sensitivity to salt stress, meaning that even at lower NaCl concentrations, Tabriz seeds experience a more rapid decline in germination compared with Mashhad seeds. The population from Mashhad exhibited relatively higher germination rates across all salinity levels, indicating better adaptation or greater intrinsic tolerance to salt stress. The lower concentration of NaCl required to achieve 50% inhibition of germination in the Tabriz population further supports the notion that this population is more sensitive to salt. These findings suggest local adaptation to environmental conditions, where the Mashhad population may be more frequently exposed to saline environments, either naturally or

Table 4. The germination rates of two *Avena sterilis* populations from Mashhad and Tabriz.

Temperature	Osmotic potential	Germination rate			
		Mashhad population		Tabriz population	
C	MPa	seed d ⁻¹ ± SE			
5	0	0.08	0.01	0.33	0.07
	−0.3	0.10	0.01	0.29	0.13
	−0.6	0.09	0.02	0.13	0.04
	−0.9	0.13	0.02	0.13	0.02
	−1.2	0.08	0.00	0.12	0.00
10	0	0.39	0.06	0.25	0.05
	−0.3	0.28	0.06	0.25	0.02
	−0.6	0.34	0.07	0.35	0.11
	−0.9	0.21	0.03	0.25	0.03
	−1.2	0.16	0.08	0.13	0.03
15	0	0.52	0.03	0.42	0.06
	−0.3	0.36	0.05	0.48	0.05
	−0.6	0.40	0.07	0.28	0.03
	−0.9	0.35	0.11	0.20	0.04
	−1.2	0.16	0.01	0.12	0.02
20	0	0.32	0.01	0.22	0.04
	−0.3	0.34	0.03	0.33	0.03
	−0.6	0.28	0.01	0.33	0.05
	−0.9	0.42	0.13	0.25	0.06
	−1.2	0.11	0.02	0.19	0.01
25	0	0.39	0.05	0.15	0.03
	−0.3	0.15	0.02	0.19	0.04
	−0.6	0.22	0.04	0.22	0.04
	−0.9	0.21	0.10	0.12	0.00
	−1.2	0.08	0.00	0.10	0.01
30	0	0.26	0.01	0.17	0.03
	−0.3	0.30	0.08	0.23	0.06
	−0.6	0.17	0.05	0.21	0.01
	−0.9	0.12	0.02	0.12	0.02
	−1.2	0.08	0.00	0.10	0.00

**Figure 3.** Effect of sodium chloride (NaCl) concentration on the germination percentage of *Avena sterilis* populations from Mashhad and Tabriz. X_{50} indicates the concentration of NaCl required to inhibit 50% of germination. The vertical bars represent the standard error (\pm SE). The equations (four-parameter log-logistic) for the populations of Mashhad and Tabriz are:

$$Y(\text{Mashhad}) = 7.14 \frac{89.92}{1 + 10^{[(4.76 - X)/(-0.34)]}}$$

$$Y(\text{Tabriz}) = 2.96 \frac{94.69}{1 + 10^{[(3.90 - X)/(-0.36)]}}$$

through agricultural practices, resulting in enhanced salt tolerance. In contrast, the Tabriz population may be adapted to less-saline conditions, which accounts for its increased sensitivity to salt. This

underscores the importance of understanding population-level variations in salinity tolerance for improved crop management and breeding strategies in areas susceptible to salinity (Amini et al. 2024).

Light/Dark Regimes

The germination rate was significantly affected by the interaction between population and light/dark regime ($P < 0.01$). In both populations, the highest germination percentage was observed under a light/dark regime of 10/14 h (Figure 4). The lowest germination percentages were recorded in the Mashhad (13%) and Tabriz (10%) populations under continuous darkness (0/24-h light/dark) (Figure 4). Germination rates under alternating light/dark regimes ranged from 64% to 93% in the Mashhad population and from 35% to 91% in the Tabriz population (Figure 4).

These results indicate that a period of darkness followed by light exposure, which simulates natural environmental conditions, is optimal for seed germination. Light plays a crucial role in regulating seed dormancy and germination through phytochromes, which are sensitive to light (Amini et al. 2017; Yang et al. 2020). The lowest germination percentages observed in both populations under continuous darkness (0/24-h light/dark) underscore the essential role of light in the germination process. Seeds kept in prolonged darkness may fail to undergo critical physiological changes, such as the breakdown of stored reserves and enzyme activation, which often depend on light. This suggests that both populations are adapted to environments where light significantly influences seed germination, with the Mashhad population potentially demonstrating greater resilience or adaptability to light/dark cycles. The study conducted by Chauhan et al. (2018) demonstrates that windmill grass (*Chloris truncata* R. Br.) seeds placed on the soil surface exhibit a higher germination percentage compared with those buried in the soil. This phenomenon can be attributed to the increased availability of light and oxygen during the germination process, which are more accessible at the soil surface.

Although there were differences in the germination ecology of *A. sterilis* populations between Mashhad and Tabriz, both populations exhibited a strong response to various temperatures, humidity levels, light conditions, and salinity. The variations in how these populations respond to environmental factors may be attributed to local adaptations to the distinct climatic and soil conditions in each region. Factors such as temperature fluctuations, precipitation, and soil composition significantly influence seed dormancy and germination, driving adaptations that enable each population to survive and establish itself in its specific environment. Understanding these germination patterns is essential for developing targeted weed management strategies. For Mashhad, where conditions favor broader germination, early-season herbicide applications, coupled with precise irrigation management to avoid excess moisture, can prevent premature germination. Additionally, implementing crop rotation as a long-term strategy will help reduce *A. sterilis* proliferation. In Tabriz, where the germination window may be narrower, a more targeted approach utilizing postemergence herbicides, along with mechanical controls such as mowing or plowing, would be more effective in managing established plants. Given the favorable conditions for *A. sterilis* germination across Iran, its unchecked spread remains a significant concern. Therefore, the integration of these tailored management strategies—incorporating predictive emergence models and optimal herbicide timing—will be crucial to prevent




Figure 4. Effect of light/dark regimes on the germination percentage of *Avena sterilis* populations from Mashhad and Tabriz. The vertical bars represent the standard error (\pm SE).

the spread of *A. sterilis* and minimize its impact on cropping systems and pastures.

Funding statement. This research received no specific grant from any funding agency or the commercial or not-for-profit sectors.

Competing interests. The authors have no conflicts of interest to declare.

References

- Abdellaoui R, Boughalleb F, Zayoud D, Neffati M, Bakhshandeh E (2019) Quantification of *Retama raetam* seed germination response to temperature and water potential using hydrothermal time concept. *Environ Exp Bot* 157:211–216
- Amini R, Gholami F, Ghanepour S (2017) Effects of environmental factors and burial depth on seed germination and emergence of two populations of *Caucalis platycarpus*. *Weed Res* 57:247–256
- Amini R, Hasanfard A, Ahmadian N, Yuzband FZ (2024) Effect of environmental, seed burial depth, and straw mulch on germination and seedling emergence in *Cichorium glandulosum*. *Weed Sci* 72:164–171
- Anwar MP, Islam AM, Yeasmin S, Rashid MH, Juraimi AS, Ahmed S, Shrestha A (2021) Weeds and their responses to management efforts in a changing climate. *Agronomy* 11:1921
- Armin M, Asghripour M (2011) Effect of plant density on wild oat competition with competitive and non-competitive wheat cultivars. *Agric Sci China* 10:1554–1561
- Baskin CC, Baskin JM (1998) *Seeds: Ecology, Biogeography, and Evolution of Dormancy and Germination*. San Diego: Academic Press. 1600 p
- Bewley JD, Bradford K, Hilhorst H, Nonogaki H (2013) *Seeds: Physiology of Development, Germination and Dormancy*. 3rd ed. New York: Springer. 392 p
- Chauhan BS, Manalil S, Florentine S, Jha P (2018) Germination ecology of *Chloris truncata* and its implication for weed management. *PLoS ONE* 13: e0199949
- Dürr C, Dickie JB, Yang XY, Pritchard HW (2015) Ranges of critical temperature and water potential values for the germination of species worldwide: contribution to a seed trait database. *Agric For Meteorol* 200:222–232
- Hasanfard A, Rastgoo M, Darbandi EI, Nezami A, Chauhan BS (2022) Freezing stress affects the efficacy of clodinafop-propargyl and 2, 4-D plus MCPA on wild oat (*Avena ludoviciana* Durieu) and turnipweed [*Rapistrum rugosum* (L.) All.] in wheat (*Triticum aestivum* L.). *PLoS ONE* 17:e0274945
- Koç İ (2021) Examining seed germination rate and seedlings gas exchange performances of some Turkish red pine provenances under water stress. *Düzce Üniversitesi Bilim ve Teknoloji Dergisi* 9:48–60
- Loskutov IG, Gnutikov AA, Blinova, EV, Rodionov AV (2023) The application of Vavilov's approaches to the phylogeny and evolution of cultivated species of the genus *Avena* L. *Vavilov J Genet Breeding* 27:921
- Nikolić N, Ghirardelli A, Schiavon M, Masin R (2023) Effects of the salinity-temperature interaction on seed germination and early seedling development: a comparative study of crop and weed species. *BMC Plant Biol* 23:446
- Nosratti I, Sabeti P, Chaghmirzaee G, Heidari H (2020) Weed problems, challenges, and opportunities in Iran. *Crop Prot* 134:104371
- Poljakoff-Mayber A, Somers GF, Werker E, Gallagher JL (1994) Seeds of *Kosteletzkya virginica* (Malvaceae): their structure, germination, and salt tolerance. II. Germination and salt tolerance. *Am J Bot* 81:54–59
- Ramesh K (2015) Weed problems, ecology, and management options in conservation agriculture: issues and perspectives. *Adv Agron* 131:251–303
- Rastgoo M, Hasanfard A (2021) Desertification in agricultural lands: approaches to mitigation. In Zhu Y, Luo Q, Liu Y, eds. *Deserts and Desertification*. London: IntechOpen. <https://doi.org/10.5772/intechopen.98795>
- Rastgoo M, Izadi Darbandi E, Hasanfard A, Mijani S (2023) Seed germination modeling of sterile oat biotypes susceptible and resistant to ACCase-inhibiting herbicides in response to temperature and drought stress. *J Crop Prot* 12:183–195
- Saburi-Rad S, Kafi M, Nezami A, Bannayan Aval M (2012) Study on seed germination behavior of *Kochia scoparia* L. Schard in response to temperature and water potential. *Iran J Range Desert Res* 18:578–592
- Steinmaus SJ, Prather TS, Holt JS (2000) Estimation of base temperatures for nine weed species. *J Exp Bot* 51:275–286
- Toscano S, Romano D, Tribulato A, Patanè C (2017) Effects of drought stress on seed germination of ornamental sunflowers. *Acta Physiol Plant* 39:1–12
- Wong PPY, Lai PC, Low CT, Chen S, Hart M (2016) The impact of environmental and human factors on urban heat and microclimate variability. *Build Environ* 95:199–208
- Yang L, Liu S, Lin R (2020) The role of light in regulating seed dormancy and germination. *J Integr Plant Biol* 62:1310–1326