


This article was downloaded by: [Ferdowsi University of Mashhad]

On: 18 January 2010

Access details: Access Details: [subscription number 912974449]

Publisher Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Phosphorus, Sulfur, and Silicon and the Related Elements

Publication details, including instructions for authors and subscription information:

<http://www.informaworld.com/smpp/title~content=t713618290>

The Green Method for Regiospecific Ring Opening of Epoxide with Dimercaptoethane

Seyed Mohammad Seyedi ^a; Hamid Sadeghian ^a; Masoomeh Rezai ^a

^a Department of Chemistry, Faculty of Sciences, Ferdowsi University of Mashhad, Mashhad, I. R., Iran

To cite this Article Seyedi, Seyed Mohammad, Sadeghian, Hamid and Rezai, Masoomeh(2007) 'The Green Method for Regiospecific Ring Opening of Epoxide with Dimercaptoethane', Phosphorus, Sulfur, and Silicon and the Related Elements, 182: 8, 1709 – 1716

To link to this Article: DOI: 10.1080/10426500701313888

URL: <http://dx.doi.org/10.1080/10426500701313888>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.informaworld.com/terms-and-conditions-of-access.pdf>

This article may be used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

The Green Method for Regiospecific Ring Opening of Epoxide with Dimercaptoethane

Seyed Mohammad Seyedi

Hamid Sadeghian

Masoomeh Rezai

Department of Chemistry, Faculty of Sciences, Ferdowsi University of Mashhad, Mashhad, I. R. Iran

A simple, efficient, and environmentally friendly one-step regiospecific ring opening of strained epoxides with dimercaptoethane as a nucleophile in nearly saturated aqueous potassium carbonate is reported. The synthesized β, β' -dihydroxydithioether compounds were confirmed by elemental analysis, ^1H NMR, ^{13}C NMR, and MS spectral studies.

INTRODUCTION

The ring opening of epoxides¹ with various nucleophils is an important synthetic transformation in organic chemistry. The preparation of β -hydroxy thioethers is generally realized by the action of thiolates² or thiols in the presence of metal salts such as Lanthanum³, Cobalt,⁴ Titanium⁵, complex of Gallium-Lithium *bis*-naphthoxide⁶ and Lewis acids ZnCl_2 ,⁷ and InCl_3 .⁸ Ring opening reaction by thiols was also carried out in the presence of alumina⁹ and polyethylene glycol.¹⁰ This reaction has also been reported in hexafluoroisopropanol without using catalyst.¹¹ The ring opening of oxiranes with dimercaptoethane has been recently performed in the presence of benzyltrimethylammonium hydroxide (Triton B) as catalyst.¹² Highly regioselective ring opening of epoxides by thiophenols in ionic liquids has also been reported.¹³ Herein, we report the ring opening reaction of two mole equivalents of epoxides with dimercaptoethane in saturated aqueous solution of potassium carbonate to prepare corresponding β, β' -dihydroxy dithioethers. Having two secondary hydroxy groups, make these compounds susceptible reagents to synthesize new thiocrown ethers possessing various sidearms.¹⁴

Received November 18, 2006; accepted February 1, 2007.


Address correspondence to Seyed Seyedi, Department of Chemistry, School of Sciences, Ferdowsi University of Mashhad, Mashhad, 41775, Iran. E-mail: smseyedi@yahoo.com

Many efforts are developed to perform chemical reactions in aqueous media to safe environmental conditions.¹⁵ Water is cheap, readily available, and nontoxic, resulting in an economic process that has clear advantages as an environmentally friendly solvent alternative in organic synthesis.¹⁶

RESULTS AND DISCUSSION



















The ring opening of the starting epoxides (**1a-i**) was regionspecific by nucleophilic attack on the terminal carbon atoms affording a secondary diols. The water solubility of epoxides are decreased in the presence of high concentrations of potassium carbonate lead to decrease epoxide ring opening percentage by the protic solvent.¹⁷ Dimercaptoethane is deprotonated by carbonate anion (rather than water deprotonation) lead to formation of water soluble dimercaptid anion, which can react with epoxides in the boundary surface of aqueous and organic phase. This is a suitable condition of the S_N2 mechanism resulted nucleophilic attack on low substituted site.¹⁷ Obviously, the ring opening of epoxides (**1a-i**) afforded corresponding dihydroxy dithioethers as a mixture of isomeric diastereomers.¹² It was notable that in low concentrations of potassium carbonate, the formation of mono-substituted ethandiols—instead of formation of corresponding β,β' -dihydroxydithioethers—was increased. This was obvious when the results were checked by TLC and sodium metaperiodate-benzidine test of 1,2-diols.¹⁸

The β,β' -dihydroxydithioethers (**3a-i**) were prepared by the action of two mole equivalents of epoxides (**1a-i**) with deprotonated dimercaptoethane (**2**), which was formed by proton abstracting of carbonate anion under reflux condition and vigorous stirring (Scheme 1). All



SCHEME 1

TABLE I Thiolysis by Dimercaptoethane of Epoxides (1a-i) in Aqueous Potassium Carbonate to Give Beta, Beta'-dihydroxy Dithioether

Epoxides	Beta, Beta'-dihydroxy Dithioether	Time (hour)	Yield (%)
		5	94
		4	97
		3	91
		3	96
		3	96
		3	96
		3	99
		3	96
		3	89

dihydroxy compounds were obtained in excellent (Table I) and indicated high regioselectivity by showing one spot on TLC, which did not need further purification. The dihydroxy compounds (**3a–g**) were isolated by decanting and (**3h–i**) by filtering follow by washing with water.

CONCLUSION

The method used here is a simple, efficient, and environmentally friendly procedure with excellent yields, high regioselectivity, and no need for any organic solvents either for reaction medium or extracting the products. Therefore, the work follows the basic aims which are important to green chemistry.¹⁹

EXPERIMENTAL

Melting points were recorded on an Electrothermal type 9100 melting point apparatus. ¹H NMR (500 MHz) and ¹³C NMR (125 MHz) were obtained by using a Bruker Avance DRX-500 Fourier transformer spectrometer. Chemical shifts are reported in ppm (δ) downfield from tetramethylsilane (TMS). Electron impact (EI) mass spectra were recorded on a Varian Match 7A spectrometer. Elemental analysis was obtained on a Thermo Finnigan Flash EA microanalyzer. All chemicals were purchased from Merck and Fluka Co. and used without further purification.

General Procedure for Ring Opening of Epoxides **1a–i** with Dimercaptoethane (**2**)

To a solution of potassium carbonate (50 g, 350 mmol) in water (65 ml) was added dimercaptoethane (**2**) (13.5 ml, 160 mmol) followed by epoxide (300 mmol). The mixture was refluxed in oil bath while stirring vigorously. The completion of reaction was checked by TLC (silica gel 60 F₂₅₄, benzene-ethylacetate 50:50).

After completion, the mixture was cooled, and the precipitated products (**3h–i**) were filtered, washed with water (3 \times 50 ml), and dried in oven at 50–55° for 4h. The liquid products (**3a–g**) in aqueous phase were decanted, washed with water (3 \times 50 ml), and then heated in oven at 80–90° for 2h to eliminate water. The remainder of the carbonate solution was reused in subsequent runs.

1-(2-[(2-Hydroxypropyl)sulfanyl]ethylsulfanyl)-2-propanol **3a**

Colorless oil (Found: C, 45.91; H, 8.57; S, 30.27. C₈H₁₈O₂S₂ requires C, 45.68; H, 8.62; S, 30.49%); δ_{H} 1.25 (6H, d, *J* 6.3, 2 \times CH₃), 2.51 (2H,

dd, J 8 and 20, $2 \times \text{CH}_2\text{S}$), 2.73 (2H, dd, J 4.1 and 12.5, $2 \times \text{CH}_2\text{S}$), 2.78 (4H, s, $\text{SCH}_2\text{CH}_2\text{S}$), 2.92 (2H, br, $2 \times \text{OH}$), 3.87 (2H, m, $2 \times \text{CH}$); δ_{C} 21.96 ($2 \times \text{q}$), 32.41 ($2 \times \text{t}$), 41.40 ($2 \times \text{t}$), 65.92 ($2 \times \text{d}$); m/z EI 210 (M^+), 165 (12%) $\text{C}_6\text{H}_{13}\text{OS}_2$, 151 (58) $\text{C}_5\text{H}_{11}\text{OS}_2$, 119 (65) $\text{C}_5\text{H}_{11}\text{OS}$, 75 (100) $\text{C}_3\text{H}_7\text{S}$.

1-(2-[(2-Hydroxybutyl)sulfanyl]ethylsulfanyl)-2-butanol 3b

Colorless oil (Found: C, 50.20; H, 9.34; S, 26.82. $\text{C}_{10}\text{H}_{22}\text{O}_2\text{S}_2$ requires C, 50.38; H, 9.30; S, 26.90%); δ_{H} 0.97 (6H, t, J 7.5, $2 \times \text{CH}_3$), 1.54 (4H, m, $2 \times \text{CH}_3\text{CH}_2$), 2.52 (2H, dd, J 8 and 20, $2 \times \text{CH}_2\text{S}$), 2.75 (2H, dd, J 4.1 and 12.5, $2 \times \text{CH}_2\text{S}$), 2.78 (4H, s, $\text{SCH}_2\text{CH}_2\text{S}$), 2.93 (2H, br, $2 \times \text{OH}$), 3.61 (2H, m, $2 \times \text{CH}$); δ_{C} 9.82 ($2 \times \text{q}$), 28.87 ($2 \times \text{t}$), 32.24 ($2 \times \text{t}$), 39.46 ($2 \times \text{t}$), 71.02 ($2 \times \text{d}$); m/z EI 238 (M^+), 179 (18%) $\text{C}_7\text{H}_{15}\text{OS}_2$, 133 (94) $\text{C}_6\text{H}_{13}\text{OS}$, 105 (95) $\text{C}_4\text{H}_9\text{OS}$, 75 (100) $\text{C}_3\text{H}_7\text{S}$.

2-(2-[(2-Hydroxy-2-phenylethyl)sulfanyl]ethylsulfanyl)-1-phenyl-1-ethanol 3c

Colorless oil (Found: C, 64.68; H, 6.57; S, 19.09. $\text{C}_{18}\text{H}_{22}\text{O}_2\text{S}_2$ requires C, 64.63; H, 6.63; S, 19.17%); δ_{H} 2.55-2.93 (4H, m, $4 \times \text{CH}_2\text{S}$), 2.76 (4H, s, $\text{SCH}_2\text{CH}_2\text{S}$), 3.45 (2H, br, $2 \times \text{OH}$), 4.74 (2H, m, $2 \times \text{CH}$), 7.37 (10 H, m, $2 \times \text{Ph}$); δ_{C} 32.63 ($2 \times \text{t}$), 42.78 ($2 \times \text{t}$), 77.65 ($2 \times \text{d}$), 128.74 ($2 \times \text{d}$), 130.02 (d), 130.88 ($2 \times \text{d}$), 139.84 (d); m/z EI 334 (M^+), 317 (11%) $\text{C}_{18}\text{H}_{21}\text{OS}_2$, 227 (23) $\text{C}_{11}\text{H}_{15}\text{OS}_2$, 213 (28) $\text{C}_{10}\text{H}_{13}\text{OS}_2$, 196 (72) $\text{C}_{10}\text{H}_{12}\text{S}_2$, 136 (92) $\text{C}_4\text{H}_8\text{OS}_2$, 121 (96) $\text{C}_8\text{H}_9\text{O}$, 107 (100) $\text{C}_7\text{H}_7\text{O}$.

1-(2-[(2-Hydroxy-3-isopropoxypropyl)sulfanyl]ethylsulfanyl)-3-isopropoxy-2-propanol 3d

Colorless oil (Found: C, 51.43; H, 9.21; S, 19.39. $\text{C}_{14}\text{H}_{30}\text{O}_4\text{S}_2$ requires C, 51.50; H, 9.26; S, 19.64%); δ_{H} 1.21 (12H, d, J 6.1, $2 \times \text{CH}_3\text{CHCH}_3$), 2.69 (2H, dd, J 7 and 19.5, $2 \times \text{CH}_2\text{S}$), 2.77 (2H, dd, J 5.2 and 17.4, $2 \times \text{CH}_2\text{S}$), 2.84 (4H, s, $\text{SCH}_2\text{CH}_2\text{S}$), 2.92 (2H, br, $2 \times \text{OH}$), 3.47 (2H, dd, J 6.3 and 15.7, $2 \times \text{CH}_2\text{O}$), 3.55 (2H, dd, J 4.1 and 13.5, $2 \times \text{CH}_2\text{O}$), 3.65 (2H, m, $2 \times \text{CH}_3\text{CHCH}_3$), 3.90 (2H, m, $2 \times \text{CH}$); δ_{C} 22.45 ($2 \times \text{q}$), 32.99 ($2 \times \text{t}$), 36.14 ($2 \times \text{t}$), 67.05 ($2 \times \text{d}$), 71.02 ($2 \times \text{t}$), 72.75 ($2 \times \text{t}$); m/z EI 326 (M^+), 297 (11%) $\text{C}_{12}\text{H}_{25}\text{O}_4\text{S}_2$, 209 (48) $\text{C}_8\text{H}_{17}\text{O}_2\text{S}_2$, 177 (57) $\text{C}_8\text{H}_{17}\text{O}_2\text{S}$, 103 (88) $\text{C}_5\text{H}_{11}\text{O}_2$, 59 (100) $\text{C}_3\text{H}_7\text{O}$.

1-(Tert-butoxy)-3-[(2-[3-(tert-butoxy)-2-hydroxypropyl]sulfanylethyl)sulfanyl]-2-propanol 3e

Colorless oil (Found: C, 54.43; H, 9.71; S, 17.98. $\text{C}_{16}\text{H}_{34}\text{O}_4\text{S}_2$ requires C, 54.20; H, 9.67; S, 18.09%); δ_{H} 1.18 (18H, s, $2 \times \text{C}_4\text{H}_9$), 2.68 (2H, dd, J 7 and 19.5, $2 \times \text{CH}_2\text{S}$), 2.76 (2H, dd, J 5.2 and 17.4, $2 \times \text{CH}_2\text{S}$), 2.83 (4H, s, $\text{SCH}_2\text{CH}_2\text{S}$), 2.87 (2H, br, $2 \times \text{OH}$), 3.46 (2H, dd, J 6.3 and 15.7,

2 × CH₂O), 3.54 (2H, dd, *J* 4.1 and 13.5, 2 × CH₂O), 3.86 (2H, m, 2 × CH); δ_C 27.45 (2 × q), 33.09 (2 × t), 36.24 (2 × t), 67.76 (2 × d), 71.62 (2 × t), 73.15 (2 × t); *m/z* EI 354 (M⁺), 223 (53%) C₉H₁₉O₂S₂, 166 (54) C₅H₁₀O₂S₂, 137 (60) C₄H₉OS₂, 57 (100) C₄H₉.

1-Butoxy-3-(2-[(3-butoxy-2-hydroxypropyl)sulfanyl]ethylsulfanyl)-2-propanol 3f

Colorless oil (Found: C, 54.16; H, 9.73; S, 18.22. C₁₆H₃₄O₄S₂ requires C, 54.20; H, 9.67; S, 18.09%); δ_H 0.89 (6H, t, *J* 8, 2 × CH₃), 1.18–1.69 (8H, m, 2 × CH₂CH₂), 2.64 (2H, dd, *J* 1.1, 7 and 19.5, 2 × CH₂S), 2.70 (2H, dd, *J* 5.2 and 17.4, 2 × CH₂S), 2.76 (4H, s, SCH₂CH₂S), 2.90 (2H, br, 2 × OH), 3.42 (2H, dd, *J* 6.3 and 15.7, 2 × CH₂O), 3.49 (2H, dd, *J* 4.1 and 13.5, 2 × CH₂O), 3.46 (4H, t, *J* 8, 2 × CH₂CH₂O), 3.87 (2H, m, 2 × CH); δ_C 14.44 (2 × q), 19.30 (2 × t), 31.2 (2 × t), 33.15 (2 × t), 36.08 (2 × t), 67.70 (2 × d), 71.53 (2 × t), 72.89 (2 × t); *m/z* EI 354 (M⁺), 223 (43%) C₉H₁₉O₂S₂, 191 (63) C₉H₁₉O₂S, 57 (100) C₄H₉.

1-(Allyloxy)-3-[(2-[(3-allyloxy)-2-hydroxypropyl]sulfanylethyl)sulfanyl]-2-propanol 3g

Colorless oil (Found: C, 52.16; H, 8.18; S, 19.78. C₁₄H₂₆O₄S₂ requires C, 52.14; H, 8.13; S, 19.89%); δ_H 2.68 (2H, dd, *J* 7.1 and 13.8, 2 × CH₂S), 2.76 (2H, dd, *J* 4.2 and 17.8, 2 × CH₂S), 2.81 (4H, s, SCH₂CH₂S), 3.50 (2H, dd, *J* 6.2 and 15.9, 2 × CH₂O), 3.55 (2H, dd, *J* 4.1 and 13.7, 2 × CH₂O), 3.95 (2H, m, 2 × CH), 4.05 (4H, d, *J* 5.6, 2 × CH₂=CHCH₂O), 5.06 (2H, br, 2 × OH), 5.22 (2H, dd, *J* 10 and 1.2, 2 × CH₂=CH), 5.30 (2H, dd, *J* 17 and 1.2, 2 × CH₂=CH), 5.86–5.94 (2H, m, 2 × CH₂=CH); δ_C 33.06 (2 × t), 36.08 (2 × t), 69.33 (2 × d), 70.11 (2 × t), 72.77 (2 × t), 117.99 (2 × t), 134.68 (2 × d); *m/z* EI 322 (M⁺), 247 (13%) C₁₀H₁₅O₃S₂, 207 (30) C₈H₁₅O₂S₂, 175 (54) C₈H₁₅O₂S, 147 (57) C₆H₁₁O₂S, 71 (62) C₄H₇O, 41 (100) C₃H₅.

1-(2-[(2-Hydroxy-3-phenoxypropyl)sulfanyl]ethylsulfanyl)-3-phenoxy-2-propanol 3h

White solid (Found: C, 61.13; H, 6.72; S, 16.07. C₂₀H₂₆O₄S₂ requires C, 60.88; H, 6.64; S, 16.25%); m.p. 79°C (from carbon tetrachloride); δ_H 2.39 (2H, br, 2 × OH), 2.71 (2H, dd, *J* 7.2 and 15.5, 2 × CH₂S), 2.79 (2H, dd, *J* 4.2 and 17.8, 2 × CH₂S), 2.85 (4H, s, SCH₂CH₂S), 4.04 (2H, dd, *J* 6.2 and 15.8, 2 × CH₂O), 4.12 (2H, m, 2 × CH), 4.15 (2H, dd, *J* 4.1 and 13.7, 2 × CH₂O), 6.84–7.38 (10 H, m, 2 × Ph); δ_C 33.10 (2 × t), 35.80 (2 × t), 69.91 (2 × t), 70.92 (2 × d), 114.90 (2 × d), 121.09 (d), 120.74 (2 × d), 159.23 (d); *m/z* EI 394 (M⁺), 287 (85%) C₁₃H₁₉O₃S₂, 243 (86) C₁₁H₁₅O₂S₂, 211 (90) C₁₁H₁₅O₂S, 75 (100) C₆H₅.

2-(2-[(2-Hydroxycyclohexyl)sulfanyl]ethylsulfanyl)-1-cyclohexanol 3i

White solid (Found: C, 58.13; H, 9.06; S, 22.07. $C_{14}H_{26}O_2S_2$ requires C, 57.89; H, 9.02; S, 22.08%); m.p. 72°C (from carbon tetrachloride); δ_H 1.25–1.54 (8H, m, $2 \times CH_2CH_2$), 1.73 (4H, m, $2 \times CH_2CHS$), 2.08 (4H, m, $2 \times CH_2CHOH$), 2.33 (2H, br, $2 \times OH$), 2.27–2.44 (2H, m, $2 \times CHS$), 2.80 and 2.83 (4H, two singlets, $2 \times SCH_2CH_2S$), 3.17–3.41 (2H, m, $2 \times CH$); δ_C 24.22 ($2 \times t$), 26.03 ($2 \times t$), 30.72 and 31.11 (two peaks, $2 \times t$), 32.87 and 32.97 (two peaks, $2 \times t$), 33.83 ($2 \times t$), 53.33 and 53.42 (two peaks, $2 \times d$), 72.41 and 72.79 (two peaks, $2 \times d$); m/z EI 290 (M^+), 273 (10%) $C_{14}H_{25}OS_2$, 159 (94) $C_8H_{15}OS$, 99 (95) $C_6H_{11}O$, 83 (100) C_6H_{11} .

REFERENCES

- [1] (a) S. Garzynski, *Synthesis*, 1984, 629; (b) C. Bonini and G. Righi, *Synthesis*, 1994, 225; (c) W. M. Dai, J. Wu, and A. Wu, *Tetrahedron Lett.*, 1998, **39**, 4091; (d) I. C. González and C. J. Forsyth, *Tetrahedron Lett.*, 2000, **41**, 3805; (e) O. A. Ozhogina, *Tetrahedron Lett.*, 2002, **43**, 553; (f) K. Surendra, N. S. Krishnaveni, Y. V. D. Nageswar, and K. R. Rao, *J. Org. Chem.*, 2003, **68**, 4994; (g) A. Solladié-Cavallo, P. Lupattelli, and C. Bonini, *J. Org. Chem.*, 2005, **70**, 1605; (h) R. M. Jong, J. J. W. Tiesinga, A. Villa, L. Tang, D. B. Janssen, and B. W. Dijkstra, *J. Am. Chem. Soc.*, 2005, **127**, 13338.
- [2] (a) E. J. Corey, D. A. Clark, G. Goto, A. Marfat, C. Mioskowski, B. Samuelsson, and S. Hammarstorm, *J. Am. Chem. Soc.*, 1980, **102**, 1436; (b) M. F. Sebban, P. Vottero, A. Alagui, and G. Dupuy, *Tetrahedron Lett.*, 2000, **41**, 1019; (c) S. Patai, Ed., *The Chemistry of The Thiol Group*, (John Wiley, New York, 1974).
- [3] A. E. Vougiokas and H. B. Kagan, *Tetrahedron Lett.*, 1987, **28**, 6065.
- [4] J. Iqbal, A. Pandey, A. Shukla, R. R. Srivastava, and S. Tripathi, *Tetrahedron*, 1990, **46**, 6423.
- [5] (a) S. Y. Ko and K. B. Sharpless, *J. Org. Chem.*, 1986, **51**, 5413; (b) Y. Belokon, M. Flego, N. Ikonnikov, M. Moscalenko, and M. North, *Tetrahedron Asymm.*, 1996, **7**, 851; (c) Y. Jiang, L. Gong, X. Feng, and W. Hu, *Tetrahedron*, 1997, **53**, 14327; (d) J. Wu, X. L. Hou, L. J. Xia and M. H. Tang, *Tetrahedron Asymm.*, 1998, **9**, 3431.
- [6] (a) R. M. Hanson and K. B. Sharpless, *J. Org. Chem.*, 1986, **51**, 1922; (b) K. Narasaka, N. Iwasawa, M. Inoue, T. Yamada, and M. Nakashima, *J. Am. Chem. Soc.*, 1989, **111**, 5340. (c) K. Mikami, M. Terada, and T. Nakai, *J. Am. Chem. Soc.*, 1990, **112**, 3949; (d) H. Sasai, T. Arai, Y. Satow, K. N. Houk, and M. Shibasaki, *J. Am. Chem. Soc.*, 1995, **117**, 6194; (e) T. Arai, Y. M. A. Yamada, N. Yamamoto, H. Sasai, and M. Shibasaki, *Chem. Eur. J.*, 1996, **2**, 1368; (f) T. Aria, H. Sasai, K. Aoe, K. Okamura, and M. Shibasaki, *Angew. Chem. Int. Ed. Engl.*, 1996, **35**, 104.
- [7] (a) D. Amantini, F. Fringuelli, F. Pizzo, S. Tortoioli, and L. Vaccaro, *Synlett*, 2003, **15**, 2292; (b) F. Fringuelli, F. Pizzo, S. Tortoioli, and L. Vaccaro, *J. Org. Chem.*, 2003, **68**, 8248.
- [8] F. Fringuelli, F. Pizzo, S. Tortoioli, and L. Vaccaro, *Adv. Synth. Catal.*, 2002, **344**, 379.
- [9] (a) G. H. Posmer and O. Z. Rogers, *J. Am. Chem. Soc.*, 1977, **99**, 8208; (b) A. S. Rao, S. K. Paknihar and J. G. Kirtane, *Tetrahedron*, 1983, **39**, 2323.
- [10] A. K. Maiti and P. Behattacharyya, *Tetrahedron*, 1994, **50**, 10483.

- [11] V. Kesavan, D. Bennet-Delpar, and P. J. Bégué, *Tetrahedron Lett.*, 2000, **41**, 2895.
- [12] M. R. Younes, M. M. Chaabouni, and A. Baklouti, *Tetrahedron Lett.*, 2001, **42**, 3167.
- [13] J. Chen, H. Wu, C. Jin, X. Zhang, Y. Xie, and W. Su, *Green Chem.*, 2006, **8**, 330.
- [14] (a) V. Guyon, A. Guy, J. Foos, M. Lemaire, and M. Draye, *Tetrahedron*, **51**, 4065 (1995); (b) G. W. Gokel, W. M. Leevy, and E. M. Weber, *Chem. Rev.*, **104**, 2723 (2004).
- [15] P. T. Anastas, Ed., *Green Chemistry: Challenging Perspective*, Oxford Sc. Pub., Oxford, 1999.
- [16] V. Pironti and S. Colonna, *Green. Chem.*, 2005, **7**, 43.
- [17] F. A. Carey and R. J. Sundberg, *Advanced Organic Chemistry*, (Kluwer Academic/Plenum Pub., New York, 2000), 4th. ed., Vol. 1, Chap. 5, pp. 263–295.
- [18] E. Stahl, Ed., *Thin-Layer Chromatography* (Springer-Verlag Berlin, Heidelberg, New York, 1969), 2nd ed., Chap. Z, p. 885.
- [19] J. H. Clark, *Green Chem.*, 2006, **8**, 17.