

Proceedings

of the British Society of Animal Science
and the Agricultural Research Forum

2010

Advances in Animal Biosciences

This book is part of a series which is a companion to the journal ANIMAL



- 290 Fatty acid composition of milk from Holstein cows fed Fish oil, Canola oil, or their combination in early lactation
T Sadat Vafa, A A Naserian, A R Heravi Moussavi, R Valizadeh, M Danesh Mesgaran, M A Khorashadizadeh
- 291 Effects of different levels of fish oil and canola oil on productive performance of early lactating Holstein dairy cows
T Sadat Vafa, A A Naserian, A R Heravi Moussavi, R Valizadeh, M Danesh Mesgaran, M A Khorashadizadeh
- 292 Evaluation of nutritional values of caraway-seed pulp fed to Holstein dairy cattle
M M Moheghi, A.M Tahmasbi, A.A Naserian, A Halim, M Kazemi
- 293 The effect of grazing different pasture herbage masses on rumen pH in lactating dairy cows
E Lewis, F Coughlan, M O'Donovan, C Wims
- 294 Effect of rumen acid load on *in vitro* ruminal total bacteria and *fibrobacter succinogenes* populations determined by real-time PCR
S Danesh Mesgaran, A Heravi Mousavi, J Arshami, A Vakili, S Ghovvati, H Jahani-azizabadi

DAIRY

- 295 Effect of different sources of pectin feedstuffs on milk yield and composition of early lactating Holstein cows
M Kordi, A A Naserian, R Valizadeh, A M Tahmasbi
- 296 Investigation of milking equipment cleaning procedures in three geographical areas where milk chemical residues were identified in bulk milk
D Gleeson, B O'Brien
- 297 Comparative grazing behaviour of Holstein-Friesian and Jersey dairy cows and their F₁ cross in pasture based production systems
R Prendiville, K M Pierce, F Buckley
- 298 Comparative somatic cell score and milking characteristics of Holstein-Friesian, Jersey and Jersey×Holstein-Friesian cows at pasture
R Prendiville, K M Pierce, F Buckley
- 299 Animal performance and economic implications of Holstein-Friesian, Jersey and Jersey×Holstein-Friesian cows under seasonal pasture based systems
R Prendiville, L Shalloo, K M Pierce, F Buckley
- 300 Estimation of the effect of dilution due to milk yield on milk somatic cell count across cow parities
S J More, K O'Sullivan, B O'Brien, D P Berry, P T Kelly

GENETICS

- 301 Genetic aspects of lambing difficulty in Scottish Blackface Sheep
J Ilkska, J Conington, N Lambe
- 302 Genetic associations between Johne's disease and susceptibility to *Mycobacterium bovis* and *Mycobacterium avium* subsp *avium* in Irish Holstein Friesian dairy cows
M L Bermingham, S J More, M Good, A R Cromie, P Mullowney, I M Higgins, D P Berry
- 303 Genetic variation in serological response to *Mycobacterium avium* subspecies paratuberculosis and its association with performance in Irish Holstein-Friesian cows
D P Berry, M Good, A R Cromie, S J More
- 304 Transcriptional activities of bovine lactoferrin (LTF) gene promoter haplotypes
B Bahar, F O'Halloran, L Giblin, T Sweeney
- 305 The impact of selection on milk production on the lactoferrin content of milk in Irish Holstein-Friesians
S McParland, L Giblin, R F Veerkamp, D P Berry
- 306 The influence of genetic selection on the milk fatty acid profile of spring calving dairy cows
S Mc Parland, D P Berry, J Coleman, R F Veerkamp, H Soyeurt

Fatty acid composition of milk from Holstein cows fed fish oil, canola oil, or their combination in early lactation

T Sadat Vafa, A Ali Naserian, A R Heravi Moussavi, R Valizadeh, M Danesh Mesgaran, M Ali Khorashadizadeh
 Excellence Centre for Animal Science, Faculty of Agriculture, Ferdowsi University of Mashhad, Mashhad, Khorasan
 Razavi, Islamic Republic of Iran
 Email: vafa_toktam@yahoo.com

Introduction Supplementing the diet of cows with long chain unsaturated lipid decreased the medium chain fatty acids (10:0, 12:0, and 14:0) and increased the 18:0 and 18:1 content of milk fat with the greatest decrease when a high linoleic acid source was fed. Fish oil which contains relatively high concentrations of two polyunsaturated fatty acids of the n-3 family: eicosapentaenoic acid (EPA, C20:5) and docosahexaenoic acid (DHA, C22:6), is an effective means of increasing milk fat CLA content, due to an inhibition of *trans* C18:1 reduction in the rumen that results in an increased supply of *trans*-11 C18:1 available for endogenous conversion in the mammary gland. The combination of marine lipids with plant oils is an established strategy for increasing milk fat CLA content (Palmquist and Griinari, 2006). Lipids from canola seeds contain about 6% C16:0 and as high as 58% *cis*-C18:1, which have potentially interesting characteristics to change fatty acids profile of milk fat. Our objective was to evaluate the effects of feeding fish oil and canola oil separately and in combination on the fatty acid profile of milk fat and on the yield and composition of milk from dairy.

Material and methods Eight multiparous early lactation Holstein cows (42±12 DIM, 40±6 kg daily milk yield) were fed a total mixed ration supplemented with either 0% oil (Control), 2% canola oil (CO), 2% fish oil (FO), or 1% canola oil + 1% fish oil (COFO), according to a double 4 × 4 Latin square design. Oils were added at a level of about 2% of dietary DM, resulting in a dietary ether extract content of 4.7%. Milk samples were collected at the regular milking time over a 48-h period at the last two days of each period and stored at -20°C until analysis for fatty acid composition using gas liquid chromatography (GC Younglin Acme 6000 column bpx70 100*0.25*250 micron SGE company). Using hexane, milk lipid extracted, then following extraction fatty acid methyl esters were synthesised using methanolic sodium. Methyl esters of fatty acids were analyzed by gas chromatography equipped with a flamed ionized detector and silica capillary column [30 m × 0.32 (internal diameter) with 0.25-µm film thickness] with hydrogen gas as the carrier gas. Data were analyzed as a replicated 4×4 Latin square using generalized linear model (PROC GLM, SAS Inst, Inc., Cary, NC).

Results The proportions of *trans* 18:1, *cis*-11 18:1, *trans*-10,*cis*-12 18:2, *cis*-9,*trans*-11 18:2 (CLA), 18:3, 20:5(EPA) and 22:6 (DHA) were affected significantly by oil supplemented diets (Table 1, P<0.05).

Table 1 Milk fatty acid content in cows fed control, fish oil (FO), fish oil with canola oil (FOCO), or canola oil (CO) diet

Fatty Acids (g/100g fatty acids)	Treatments ¹				SE	p Value
	Control	FO	FOCO	CO		
18:1 <i>trans</i>	0.38 ^a	1.57 ^b	0.57 ^a	0.58 ^a	0.05	0.001
18:1 <i>cis</i> -9	22.2	23.53	22.75	25.53	1.06	0.22
18:1 <i>cis</i> -11	0.78 ^a	0.84 ^a	1.81 ^b	1.12 ^b	0.08	0.006
18:2 <i>trans</i> -9, <i>trans</i> -12	0.25	0.27	0.29	0.26	0.04	0.93
18:2 <i>cis</i> -9, <i>cis</i> -12	2.56	2.74	2.98	2.84	0.06	0.11
18:2 <i>trans</i> -10, <i>cis</i> -12	0.05 ^a	0.3 ^b	0.15 ^a	0.06 ^a	0.02	0.03
18:2 <i>cis</i> -9, <i>trans</i> -11 CLA	0.47 ^a	1.05 ^b	1.26 ^b	0.88 ^{ab}	0.09	0.03
18:3	0.44 ^a	0.69 ^b	0.60 ^b	0.48 ^a	0.01	0.009
20:5 EPA	0.05 ^a	0.26 ^b	0.17 ^b	0.04 ^a	0.001	0.03
22:6 DHA	0.04 ^a	0.28 ^b	0.14 ^b	0.06 ^a	0.01	0.04

Conclusion Milk fatty acid profile was significantly affected by diet containing fish oil and canola oil. AbuGhazaleh *et al* (2003) reported that the concentrations of *cis*-9, *trans*-11 CLA in milk fat increased when cows were fed the high oleic and high linoleic sunflower seed with fish oil. The results of the current experiment reveal that combining fish oil with canola oil positively improves fatty acid composition of milk fat.

References

- AbuGhazaleh A. A, Schingoethe D. J, Hippen, A. R and Kalscheur K. F. 2003. Conjugated Linoleic Acid and Vaccenic Acid in rumen, plasma, and milk of cows fed fish oil and fats differing in saturation of 18 Carbon Fatty Acids. *Journal of Dairy Science*. 86, 3648–3660.
- Bayourthe C, Enjalbert F and Moncoulon R. 2000. Effects of different forms of canola oil fatty acids plus canola meal on milk composition and physical properties of butter. *Journal of Dairy Science*. 83, 690–696.
- Palmquist D. L and Griinari J. M. 2006. Milk fatty acid composition in response to reciprocal combinations of sunflower and fish oil in the diet. *Animal Feed Science and Technology*. 131, 358-369.