

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368


Editors: W. T. A. Harrison, H. Stoeckli-Evans,
E. R. T. Tiekink and M. Weil

N,N'-Dicyclohexyl-*N''*-(2,6-difluorobenzoyl)-*N,N'*-dimethylphosphoric triamide

Mehrdad Pourayoubi, Atekeh Tarahhomi, Arnold L. Rheingold and James A. Golen

Acta Cryst. (2011). E67, o3027

This open-access article is distributed under the terms of the Creative Commons Attribution Licence <http://creativecommons.org/licenses/by/2.0/uk/legalcode>, which permits unrestricted use, distribution, and reproduction in any medium, provided the original authors and source are cited.



Acta Crystallographica Section E: Structure Reports Online is the IUCr's highly popular open-access structural journal. It provides a simple and easily accessible publication mechanism for the growing number of inorganic, metal-organic and organic crystal structure determinations. The electronic submission, validation, refereeing and publication facilities of the journal ensure very rapid and high-quality publication, whilst key indicators and validation reports provide measures of structural reliability. The journal publishes over 4000 structures per year. The average publication time is less than one month.

Crystallography Journals **Online** is available from journals.iucr.org

N,N'-Dicyclohexyl-*N''*-(2,6-difluorobenzoyl)-*N,N'*-dimethylphosphoric triamide

Mehrdad Pourayoubi,^{a*} Atekeh Tarahhomi,^a Arnold L. Rheingold^b and James A. Golen^b

^aDepartment of Chemistry, Ferdowsi University of Mashhad, Mashhad 91779, Iran, and ^bDepartment of Chemistry, University of California, San Diego, 9500 Gilman Drive, La Jolla, CA 92093, USA

Correspondence e-mail: mehrdad_pourayoubi@yahoo.com


Received 20 September 2011; accepted 17 October 2011

Key indicators: single-crystal X-ray study; $T = 100$ K; mean $\sigma(\text{C}-\text{C}) = 0.002$ Å; R factor = 0.037; wR factor = 0.097; data-to-parameter ratio = 18.9.

In the title molecule, $\text{C}_{21}\text{H}_{32}\text{F}_2\text{N}_3\text{O}_2\text{P}$, the $\text{P}=\text{O}$ and $\text{N}-\text{H}$ groups are *syn* with respect to each other, and the P atom is bonded in a distorted tetrahedral environment. The phosphoryl group adopts an *anti* orientation with respect to the carbonyl group. The angles at the tertiary N atoms (with bond-angle sums of 358.4 and 357.0°) confirm their sp^2 character. In the crystal, inversion dimers linked by pairs of $\text{N}-\text{H}\cdots\text{O}$ hydrogen bonds generate $R_2^2(8)$ loops.

Related literature

For hydrogen-bond patterns in compounds containing a $\text{C}(\text{O})\text{NHP}(\text{O})$ skeleton, see: Toghraee *et al.* (2011); Pourayoubi *et al.* (2011). For background to phosphoric triamide compounds containing a $\text{C}(\text{O})\text{NHP}(\text{O})$ skeleton, and related bond lengths, angles and torsion angles, see: Pourayoubi *et al.* (2010); Amir Khanov *et al.* (2010); Tarahhomi *et al.* (2011). For a description of hydrogen-bond motifs, see: Bernstein *et al.* (1995).



Experimental

Crystal data

$\text{C}_{21}\text{H}_{32}\text{F}_2\text{N}_3\text{O}_2\text{P}$
 $M_r = 427.47$
 Triclinic, $P\bar{1}$
 $a = 10.2322$ (6) Å
 $b = 10.6188$ (5) Å
 $c = 11.2658$ (6) Å
 $\alpha = 69.419$ (1)°
 $\beta = 79.269$ (1)°
 $\gamma = 81.536$ (1)°
 $V = 1121.45$ (10) Å³
 $Z = 2$
 Mo $K\alpha$ radiation
 $\mu = 0.16$ mm⁻¹
 $T = 100$ K
 $0.35 \times 0.30 \times 0.25$ mm

Data collection

Bruker SMART CCD area-detector diffractometer
 Absorption correction: multi-scan (SADABS; Bruker, 2005)
 $T_{\min} = 0.946$, $T_{\max} = 0.961$
 17147 measured reflections
 5053 independent reflections
 4421 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.029$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.037$
 $wR(F^2) = 0.097$
 $S = 1.05$
 5053 reflections
 267 parameters
 1 restraint
 H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\text{max}} = 0.32$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.34$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{N1}-\text{H1N}\cdots\text{O2}^i$	0.86 (1)	1.90 (1)	2.7330 (13)	165 (1)

Symmetry code: (i) $-x + 1, -y + 1, -z$.

Data collection: SMART (Bruker, 2005); cell refinement: SAINT (Bruker, 2005); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: SHELXTL (Sheldrick, 2008); software used to prepare material for publication: SHELXTL and enCIFer (Allen *et al.*, 2004).

Support of this investigation by Ferdowsi University of Mashhad is gratefully acknowledged.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BH2384).

References

- Allen, F. H., Johnson, O., Shields, G. P., Smith, B. R. & Towler, M. (2004). *J. Appl. Cryst.* **37**, 335–338.
- Amir Khanov, O. V., Moroz, O. V., Znoviyak, K. O., Trush, E. A. & Sliva, T. Y. (2010). *Acta Cryst.* **E66**, o1102.
- Bernstein, J., Davis, R. E., Shimoni, L. & Chang, N.-L. (1995). *Angew. Chem. Int. Ed. Engl.* **34**, 1555–1573.
- Bruker (2005). *SADABS, SMART and SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Pourayoubi, M., Tarahhomi, A., Rheingold, A. L. & Golen, J. A. (2010). *Acta Cryst.* **E66**, o3159.
- Pourayoubi, M., Tarahhomi, A., Saneei, A., Rheingold, A. L. & Golen, J. A. (2011). *Acta Cryst.* **C67**, o265–o272.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Tarahhomi, A., Pourayoubi, M., Rheingold, A. L. & Golen, J. A. (2011). *Struct. Chem.* **22**, 201–210.
- Toghraee, M., Pourayoubi, M. & Divjakovic, V. (2011). *Polyhedron*, **30**, 1680–1690.

supplementary materials

Acta Cryst. (2011). E67, o3027 [doi:10.1107/S1600536811043029]

N,N'-Dicyclohexyl-*N''*-(2,6-difluorobenzoyl)-*N,N'*-dimethylphosphoric triamide

M. Pourayoubi, A. Tarahhomi, A. L. Rheingold and J. A. Golen

Comment

The patterns of hydrogen bonds and their strengths on phosphoric triamides containing a C(O)NHP(O) skeleton have been discussed (Toghraee *et al.*, 2011; Pourayoubi *et al.*, 2011). The structure determination of the title compound, P(O)[2,6-F₂—C₆H₃C(O)NH][N(CH₃)C₆H₁₁]₂ (Fig. 1), was performed as part of a project in our laboratory on the synthesis of new derivatives of benzoyl phosphoric triamides with formula XC₆H_{*n*}C(O)NHP(O)Y₂ (for example, *n* = 3, X = 2,6-F₂ and Y = NHC(CH₃)₃; Pourayoubi *et al.*, 2010).

In the title phosphoric triamide, the phosphoryl group adopts the *anti* orientation with respect to the carbonyl group; whereas it is in a *syn* position relative to the N—H unit. The tetrahedral environment at the P atom is distorted, as previously noted for other phosphoric triamides (Amirkhanov *et al.*, 2010). The P=O, C=O and P—N bond lengths are within the expected ranges (Tarahhomi *et al.*, 2011).

The hydrogen atom of the C(=O)NHP(=O) moiety is involved in an intermolecular —P=O···H—N— hydrogen bond. Pairs of this type of hydrogen bonds form hydrogen-bonded dimers in the crystal (as *R*₂²(8) rings; see: Bernstein *et al.*, 1995).

Experimental

2,6—F₂C₆H₃C(O)NHP(O)Cl₂ was prepared according to the literature method reported by Pourayoubi *et al.* (2010).

The title compound was synthesized from the reaction of 2,6—F₂C₆H₃C(O)NHP(O)Cl₂ (1.09 mmol) with *N*-methylcyclohexylamine (4.36 mmol) in dry chloroform (30 ml). The amine was added dropwise to a solution of 2,6—F₂C₆H₃C(O)NHP(O)Cl₂ at 273 K, with continuous stirring. After 4 h, the solvent was evaporated and the obtained solid was washed with distilled water and recrystallized from a mixture of chloroform and DMF (4:1 *v/v*) at room temperature. IR (KBr, *v*, cm⁻¹): 3063 (NH), 2912, 2752, 1698, 1608, 1470, 1262, 1192, 1155, 997, 864, 815.

Refinement

All non-H atoms were refined anisotropically by full-matrix least-squares on *F*². Hydrogen H1N was found in a difference map and N1—H1N distance was set at 0.87 (1) Å and allowed to refine with *U*_{iso} = 1.2*U*_{eq}(N1). All other H atoms were placed in calculated positions with C—H distances for CH₂ of 0.99 Å, CH₃ of 0.98 Å, methine CH of 1.00 Å, and C(Ar)H 0.95 Å, and with *U*_{iso} of 1.2 or 1.5 times that of the parent C atom.

Figures




Fig. 1. An *ORTEP*-style plot for the title compound. Displacement ellipsoids are given at 50% probability level and H atoms are drawn as small spheres of arbitrary radii.

N,N'-Dicyclohexyl-*N''*-(2,6-difluorobenzoyl)-*N,N'*-dimethylphosphoric triamide

Crystal data

$C_{21}H_{32}F_2N_3O_2P$

$M_r = 427.47$

Triclinic, $P\bar{1}$

Hall symbol: $-P\ 1$

$a = 10.2322$ (6) Å

$b = 10.6188$ (5) Å

$c = 11.2658$ (6) Å

$\alpha = 69.419$ (1)°

$\beta = 79.269$ (1)°

$\gamma = 81.536$ (1)°

$V = 1121.45$ (10) Å³

$Z = 2$

$F(000) = 456$

$D_x = 1.266$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 9954 reflections

$\theta = 2.4$ – 28.0 °

$\mu = 0.16$ mm⁻¹

$T = 100$ K

Block, colourless

$0.35 \times 0.30 \times 0.25$ mm

Data collection

Bruker SMART CCD area-detector diffractometer

Radiation source: fine-focus sealed tube graphite

φ and ω scans

Absorption correction: multi-scan (*SADABS*; Bruker, 2005)

$T_{\min} = 0.946$, $T_{\max} = 0.961$

17147 measured reflections

5053 independent reflections

4421 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.029$

$\theta_{\max} = 28.0$ °, $\theta_{\min} = 2.0$ °

$h = -13 \rightarrow 13$

$k = -13 \rightarrow 13$

$l = -14 \rightarrow 14$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.037$

Primary atom site location: structure-invariant direct methods

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

$wR(F^2) = 0.097$

$S = 1.05$

5053 reflections

267 parameters

1 restraint

0 constraints

H atoms treated by a mixture of independent and constrained refinement

$$w = 1/[\sigma^2(F_o^2) + (0.0429P)^2 + 0.4607P]$$

$$\text{where } P = (F_o^2 + 2F_c^2)/3$$

$$(\Delta/\sigma)_{\max} < 0.001$$

$$\Delta\rho_{\max} = 0.32 \text{ e } \text{\AA}^{-3}$$

$$\Delta\rho_{\min} = -0.34 \text{ e } \text{\AA}^{-3}$$

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
P1	0.42370 (3)	0.41644 (3)	0.20824 (3)	0.01482 (9)
F1	0.91928 (9)	0.33633 (12)	0.27902 (10)	0.0491 (3)
F2	0.68054 (9)	0.73518 (9)	0.04970 (9)	0.0363 (2)
O1	0.64907 (10)	0.41510 (13)	0.35266 (9)	0.0347 (3)
O2	0.35023 (8)	0.47928 (9)	0.09668 (8)	0.01753 (19)
N1	0.57558 (10)	0.47579 (11)	0.15972 (10)	0.0170 (2)
H1N	0.5985 (14)	0.5054 (14)	0.0781 (9)	0.020*
N2	0.34278 (10)	0.45423 (10)	0.33252 (9)	0.0165 (2)
N3	0.45145 (11)	0.25173 (11)	0.25782 (11)	0.0244 (2)
C1	0.91678 (14)	0.4643 (2)	0.19652 (16)	0.0392 (4)
C2	1.03613 (16)	0.5176 (3)	0.14059 (19)	0.0497 (5)
H2	1.1182	0.4669	0.1619	0.060*
C3	1.03445 (16)	0.6460 (3)	0.0529 (2)	0.0546 (6)
H3	1.1166	0.6842	0.0137	0.066*
C4	0.91532 (17)	0.7212 (2)	0.02025 (17)	0.0463 (5)
H4	0.9148	0.8099	-0.0405	0.056*
C5	0.79751 (14)	0.66235 (18)	0.07925 (15)	0.0333 (4)
C6	0.79311 (13)	0.53390 (17)	0.16838 (13)	0.0283 (3)
C7	0.66617 (13)	0.46961 (15)	0.23652 (13)	0.0236 (3)
C8	0.27329 (12)	0.59130 (12)	0.31148 (11)	0.0169 (2)
H8	0.2504	0.6255	0.2224	0.020*
C9	0.36202 (14)	0.69035 (14)	0.32103 (14)	0.0258 (3)
H9A	0.3890	0.6578	0.4075	0.031*
H9B	0.4439	0.6951	0.2575	0.031*
C10	0.28793 (18)	0.83088 (16)	0.29625 (17)	0.0388 (4)
H10A	0.3451	0.8918	0.3081	0.047*
H10B	0.2701	0.8673	0.2064	0.047*
C11	0.15626 (17)	0.82817 (16)	0.38598 (17)	0.0385 (4)
H11A	0.1743	0.8016	0.4753	0.046*
H11B	0.1085	0.9197	0.3635	0.046*
C12	0.06889 (15)	0.72885 (15)	0.37619 (16)	0.0328 (3)
H12A	0.0434	0.7605	0.2892	0.039*
H12B	-0.0139	0.7250	0.4385	0.039*
C13	0.14254 (13)	0.58770 (14)	0.40322 (13)	0.0222 (3)
H13A	0.1616	0.5529	0.4927	0.027*
H13B	0.0852	0.5257	0.3933	0.027*

supplementary materials

C14	0.36037 (14)	0.37310 (14)	0.46557 (12)	0.0248 (3)
H14A	0.3893	0.4297	0.5066	0.037*
H14B	0.4281	0.2975	0.4658	0.037*
H14C	0.2754	0.3381	0.5129	0.037*
C15	0.57908 (14)	0.17673 (14)	0.22547 (14)	0.0277 (3)
H15	0.6473	0.2439	0.1874	0.033*
C16	0.57337 (16)	0.11177 (15)	0.12540 (14)	0.0335 (3)
H16A	0.5479	0.1824	0.0465	0.040*
H16B	0.5042	0.0468	0.1585	0.040*
C17	0.70838 (18)	0.03820 (16)	0.09313 (15)	0.0400 (4)
H17A	0.6998	-0.0088	0.0334	0.048*
H17B	0.7750	0.1049	0.0497	0.048*
C18	0.7570 (2)	-0.06431 (18)	0.21368 (16)	0.0476 (5)
H18A	0.6957	-0.1369	0.2517	0.057*
H18B	0.8468	-0.1059	0.1906	0.057*
C19	0.76312 (19)	0.00220 (19)	0.31179 (17)	0.0465 (5)
H19A	0.8303	0.0692	0.2766	0.056*
H19B	0.7914	-0.0671	0.3901	0.056*
C20	0.62703 (17)	0.07244 (16)	0.34615 (15)	0.0362 (4)
H20A	0.6341	0.1178	0.4076	0.043*
H20B	0.5613	0.0045	0.3877	0.043*
C21	0.33158 (15)	0.17706 (14)	0.30012 (15)	0.0304 (3)
H21A	0.3101	0.1560	0.2283	0.046*
H21B	0.2564	0.2325	0.3300	0.046*
H21C	0.3483	0.0930	0.3702	0.046*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
P1	0.01077 (15)	0.01879 (16)	0.01385 (15)	0.00019 (11)	-0.00010 (11)	-0.00559 (12)
F1	0.0221 (5)	0.0863 (8)	0.0401 (6)	0.0158 (5)	-0.0113 (4)	-0.0274 (6)
F2	0.0281 (5)	0.0433 (5)	0.0395 (5)	-0.0109 (4)	0.0008 (4)	-0.0161 (4)
O1	0.0187 (5)	0.0684 (8)	0.0170 (5)	0.0035 (5)	-0.0043 (4)	-0.0164 (5)
O2	0.0119 (4)	0.0264 (5)	0.0154 (4)	-0.0013 (3)	-0.0006 (3)	-0.0092 (4)
N1	0.0116 (5)	0.0268 (5)	0.0126 (5)	-0.0012 (4)	-0.0006 (4)	-0.0074 (4)
N2	0.0140 (5)	0.0206 (5)	0.0124 (5)	0.0026 (4)	-0.0013 (4)	-0.0044 (4)
N3	0.0186 (6)	0.0198 (5)	0.0281 (6)	0.0026 (4)	0.0060 (5)	-0.0061 (5)
C1	0.0152 (7)	0.0836 (13)	0.0351 (8)	0.0013 (7)	-0.0044 (6)	-0.0418 (9)
C2	0.0152 (7)	0.1089 (17)	0.0493 (11)	-0.0083 (9)	0.0001 (7)	-0.0576 (12)
C3	0.0188 (8)	0.1161 (19)	0.0591 (12)	-0.0293 (10)	0.0157 (8)	-0.0679 (13)
C4	0.0354 (9)	0.0766 (13)	0.0434 (10)	-0.0298 (9)	0.0141 (7)	-0.0405 (10)
C5	0.0183 (7)	0.0595 (10)	0.0340 (8)	-0.0111 (7)	0.0033 (6)	-0.0307 (8)
C6	0.0125 (6)	0.0588 (10)	0.0244 (7)	-0.0046 (6)	-0.0005 (5)	-0.0274 (7)
C7	0.0127 (6)	0.0419 (8)	0.0202 (6)	0.0028 (5)	-0.0026 (5)	-0.0170 (6)
C8	0.0153 (6)	0.0198 (6)	0.0155 (6)	0.0015 (5)	-0.0024 (4)	-0.0070 (5)
C9	0.0236 (7)	0.0311 (7)	0.0262 (7)	-0.0076 (6)	0.0032 (5)	-0.0154 (6)
C10	0.0485 (10)	0.0270 (8)	0.0419 (9)	-0.0089 (7)	0.0094 (8)	-0.0186 (7)
C11	0.0439 (10)	0.0299 (8)	0.0419 (9)	0.0036 (7)	0.0046 (7)	-0.0202 (7)

C12	0.0257 (8)	0.0348 (8)	0.0367 (8)	0.0103 (6)	-0.0032 (6)	-0.0163 (7)
C13	0.0143 (6)	0.0279 (7)	0.0246 (7)	0.0012 (5)	0.0004 (5)	-0.0120 (5)
C14	0.0206 (7)	0.0320 (7)	0.0141 (6)	0.0053 (5)	-0.0007 (5)	-0.0023 (5)
C15	0.0244 (7)	0.0221 (7)	0.0261 (7)	0.0086 (5)	0.0052 (6)	-0.0041 (5)
C16	0.0380 (9)	0.0265 (7)	0.0252 (7)	0.0099 (6)	0.0039 (6)	-0.0049 (6)
C17	0.0461 (10)	0.0314 (8)	0.0272 (8)	0.0161 (7)	0.0073 (7)	-0.0051 (6)
C18	0.0545 (12)	0.0374 (9)	0.0338 (9)	0.0271 (8)	0.0017 (8)	-0.0075 (7)
C19	0.0461 (11)	0.0447 (10)	0.0351 (9)	0.0269 (8)	-0.0058 (8)	-0.0087 (7)
C20	0.0383 (9)	0.0343 (8)	0.0266 (8)	0.0154 (7)	-0.0009 (7)	-0.0080 (6)
C21	0.0281 (8)	0.0204 (7)	0.0364 (8)	-0.0035 (5)	0.0085 (6)	-0.0078 (6)

Geometric parameters (\AA , $^\circ$)

P1—O2	1.4848 (9)	C11—C12	1.523 (2)
P1—N2	1.6348 (11)	C11—H11A	0.9900
P1—N3	1.6363 (11)	C11—H11B	0.9900
P1—N1	1.6854 (10)	C12—C13	1.5295 (19)
F1—C1	1.350 (2)	C12—H12A	0.9900
F2—C5	1.3552 (19)	C12—H12B	0.9900
O1—C7	1.2205 (16)	C13—H13A	0.9900
N1—C7	1.3612 (15)	C13—H13B	0.9900
N1—H1N	0.859 (9)	C14—H14A	0.9800
N2—C14	1.4725 (15)	C14—H14B	0.9800
N2—C8	1.4819 (15)	C14—H14C	0.9800
N3—C21	1.4707 (17)	C15—C16	1.528 (2)
N3—C15	1.4810 (17)	C15—C20	1.5314 (19)
C1—C2	1.369 (2)	C15—H15	1.0000
C1—C6	1.399 (2)	C16—C17	1.532 (2)
C2—C3	1.374 (3)	C16—H16A	0.9900
C2—H2	0.9500	C16—H16B	0.9900
C3—C4	1.390 (3)	C17—C18	1.526 (2)
C3—H3	0.9500	C17—H17A	0.9900
C4—C5	1.384 (2)	C17—H17B	0.9900
C4—H4	0.9500	C18—C19	1.521 (3)
C5—C6	1.382 (2)	C18—H18A	0.9900
C6—C7	1.5046 (18)	C18—H18B	0.9900
C8—C13	1.5271 (17)	C19—C20	1.530 (2)
C8—C9	1.5294 (17)	C19—H19A	0.9900
C8—H8	1.0000	C19—H19B	0.9900
C9—C10	1.527 (2)	C20—H20A	0.9900
C9—H9A	0.9900	C20—H20B	0.9900
C9—H9B	0.9900	C21—H21A	0.9800
C10—C11	1.524 (2)	C21—H21B	0.9800
C10—H10A	0.9900	C21—H21C	0.9800
C10—H10B	0.9900		
O2—P1—N2	110.40 (5)	C11—C12—C13	110.95 (12)
O2—P1—N3	116.97 (6)	C11—C12—H12A	109.5
N2—P1—N3	105.60 (6)	C13—C12—H12A	109.5
O2—P1—N1	105.57 (5)	C11—C12—H12B	109.5

supplementary materials

N2—P1—N1	112.66 (5)	C13—C12—H12B	109.5
N3—P1—N1	105.73 (6)	H12A—C12—H12B	108.0
C7—N1—P1	126.27 (9)	C8—C13—C12	110.68 (11)
C7—N1—H1N	118.7 (10)	C8—C13—H13A	109.5
P1—N1—H1N	114.7 (10)	C12—C13—H13A	109.5
C14—N2—C8	116.58 (10)	C8—C13—H13B	109.5
C14—N2—P1	123.33 (9)	C12—C13—H13B	109.5
C8—N2—P1	118.49 (8)	H13A—C13—H13B	108.1
C21—N3—C15	116.90 (11)	N2—C14—H14A	109.5
C21—N3—P1	115.51 (9)	N2—C14—H14B	109.5
C15—N3—P1	124.57 (9)	H14A—C14—H14B	109.5
F1—C1—C2	118.10 (16)	N2—C14—H14C	109.5
F1—C1—C6	118.56 (14)	H14A—C14—H14C	109.5
C2—C1—C6	123.28 (19)	H14B—C14—H14C	109.5
C1—C2—C3	118.40 (17)	N3—C15—C16	113.22 (12)
C1—C2—H2	120.8	N3—C15—C20	111.07 (11)
C3—C2—H2	120.8	C16—C15—C20	111.21 (12)
C2—C3—C4	121.52 (15)	N3—C15—H15	107.0
C2—C3—H3	119.2	C16—C15—H15	107.0
C4—C3—H3	119.2	C20—C15—H15	107.0
C5—C4—C3	117.7 (2)	C15—C16—C17	111.21 (14)
C5—C4—H4	121.1	C15—C16—H16A	109.4
C3—C4—H4	121.1	C17—C16—H16A	109.4
F2—C5—C6	118.33 (12)	C15—C16—H16B	109.4
F2—C5—C4	118.37 (17)	C17—C16—H16B	109.4
C6—C5—C4	123.29 (16)	H16A—C16—H16B	108.0
C5—C6—C1	115.79 (14)	C18—C17—C16	111.24 (13)
C5—C6—C7	124.16 (13)	C18—C17—H17A	109.4
C1—C6—C7	120.02 (15)	C16—C17—H17A	109.4
O1—C7—N1	123.71 (12)	C18—C17—H17B	109.4
O1—C7—C6	120.91 (11)	C16—C17—H17B	109.4
N1—C7—C6	115.38 (11)	H17A—C17—H17B	108.0
N2—C8—C13	111.30 (10)	C19—C18—C17	111.16 (14)
N2—C8—C9	112.22 (10)	C19—C18—H18A	109.4
C13—C8—C9	110.75 (10)	C17—C18—H18A	109.4
N2—C8—H8	107.4	C19—C18—H18B	109.4
C13—C8—H8	107.4	C17—C18—H18B	109.4
C9—C8—H8	107.4	H18A—C18—H18B	108.0
C10—C9—C8	110.65 (12)	C18—C19—C20	110.98 (16)
C10—C9—H9A	109.5	C18—C19—H19A	109.4
C8—C9—H9A	109.5	C20—C19—H19A	109.4
C10—C9—H9B	109.5	C18—C19—H19B	109.4
C8—C9—H9B	109.5	C20—C19—H19B	109.4
H9A—C9—H9B	108.1	H19A—C19—H19B	108.0
C11—C10—C9	111.80 (13)	C19—C20—C15	110.31 (12)
C11—C10—H10A	109.3	C19—C20—H20A	109.6
C9—C10—H10A	109.3	C15—C20—H20A	109.6
C11—C10—H10B	109.3	C19—C20—H20B	109.6
C9—C10—H10B	109.3	C15—C20—H20B	109.6

H10A—C10—H10B	107.9	H20A—C20—H20B	108.1
C12—C11—C10	110.74 (13)	N3—C21—H21A	109.5
C12—C11—H11A	109.5	N3—C21—H21B	109.5
C10—C11—H11A	109.5	H21A—C21—H21B	109.5
C12—C11—H11B	109.5	N3—C21—H21C	109.5
C10—C11—H11B	109.5	H21A—C21—H21C	109.5
H11A—C11—H11B	108.1	H21B—C21—H21C	109.5
O2—P1—N1—C7	164.83 (11)	P1—N1—C7—C6	-178.43 (10)
N2—P1—N1—C7	44.27 (13)	C5—C6—C7—O1	-131.18 (15)
N3—P1—N1—C7	-70.59 (12)	C1—C6—C7—O1	47.0 (2)
O2—P1—N2—C14	157.51 (10)	C5—C6—C7—N1	49.58 (19)
N3—P1—N2—C14	30.20 (11)	C1—C6—C7—N1	-132.22 (14)
N1—P1—N2—C14	-84.74 (11)	C14—N2—C8—C13	-50.03 (14)
O2—P1—N2—C8	-37.41 (10)	P1—N2—C8—C13	143.89 (9)
N3—P1—N2—C8	-164.72 (9)	C14—N2—C8—C9	74.73 (13)
N1—P1—N2—C8	80.35 (9)	P1—N2—C8—C9	-91.36 (11)
O2—P1—N3—C21	-60.31 (12)	N2—C8—C9—C10	178.90 (11)
N2—P1—N3—C21	62.93 (11)	C13—C8—C9—C10	-56.04 (15)
N1—P1—N3—C21	-177.45 (10)	C8—C9—C10—C11	55.54 (17)
O2—P1—N3—C15	99.32 (12)	C9—C10—C11—C12	-55.57 (19)
N2—P1—N3—C15	-137.44 (11)	C10—C11—C12—C13	56.09 (18)
N1—P1—N3—C15	-17.82 (13)	N2—C8—C13—C12	-177.37 (10)
F1—C1—C2—C3	177.59 (14)	C9—C8—C13—C12	57.05 (14)
C6—C1—C2—C3	0.3 (2)	C11—C12—C13—C8	-57.15 (16)
C1—C2—C3—C4	-0.3 (2)	C21—N3—C15—C16	54.69 (16)
C2—C3—C4—C5	0.0 (2)	P1—N3—C15—C16	-104.68 (13)
C3—C4—C5—F2	179.09 (13)	C21—N3—C15—C20	-71.27 (17)
C3—C4—C5—C6	0.2 (2)	P1—N3—C15—C20	129.36 (12)
F2—C5—C6—C1	-179.04 (12)	N3—C15—C16—C17	178.75 (11)
C4—C5—C6—C1	-0.2 (2)	C20—C15—C16—C17	-55.36 (16)
F2—C5—C6—C7	-0.8 (2)	C15—C16—C17—C18	54.61 (19)
C4—C5—C6—C7	178.10 (13)	C16—C17—C18—C19	-55.4 (2)
F1—C1—C6—C5	-177.37 (12)	C17—C18—C19—C20	56.9 (2)
C2—C1—C6—C5	-0.1 (2)	C18—C19—C20—C15	-57.2 (2)
F1—C1—C6—C7	4.3 (2)	N3—C15—C20—C19	-176.48 (14)
C2—C1—C6—C7	-178.47 (13)	C16—C15—C20—C19	56.45 (18)
P1—N1—C7—O1	2.4 (2)		

Hydrogen-bond geometry (Å, °)

<i>D</i> —H··· <i>A</i>	<i>D</i> —H	H··· <i>A</i>	<i>D</i> ··· <i>A</i>	<i>D</i> —H··· <i>A</i>
N1—H1N···O2 ⁱ	0.86 (1)	1.90 (1)	2.7330 (13)	165.(1)

Symmetry codes: (i) $-x+1, -y+1, -z$.

Fig. 1

