

ISSN:2229-712X

Available online at www.elixirjournal.org

Statistics

Elixir Statistics 52A (2012) 11691-11695

Elixir Online
Journal

On Bayesian estimator from exponential distribution based on records with presence of outliers

Parviz Nasiri¹ and Mehdi Jabbari Nooghabi²¹Department of Statistics, University of Payame Noor, 19395-4697 Tehran, I. R of Iran.²Department of Statistics, Ferdowsi University of Mashhad, Mashhad, I. R of Iran.

ARTICLE INFO

Article history:

Received: 24 September 2012;

Received in revised form:

19 November 2012;

Accepted: 29 November 2012;

Keywords

Exponential model,
Bayes estimators,
Outliers,
Records value.

ABSTRACT

In this paper, Bayes estimator is derived for the parameter of the exponential model with presence of outliers based on records value. This estimator compared with estimator when we not use of records value.

© 2012 Elixir All rights reserved.

Introduction

The exponential distribution plays an important role in life testing problems. A great deal with research has been done on estimating the parameters of exponential distribution using both classical and Bayesian techniques and a very good summary of this work can be found in Johanson, Kotz, and Balakrishnan (1994). There are also some papers on estimation and prediction for exponential distribution parameters based of record and censored samples. See for example Balasubramanian and Balakrishnan (1992), Ohandrasekar, Leo Alexander and Balakrishnan (2002), Jaheen (2004), Ahmadi, Doostparast and Parsian (2005) and references therein. In 2001, Dixit and Nasiri have estimated the parameter of exponential distribution with presence of outliers, and in 2009 Asgharzadeh has estimated the parameter of exponential model based on records value.

In section two, we estimate Bayes estimator of parameter of the exponential distribution with presence of outlier, in section three, we estimate the parameter (θ) of exponential distribution based on records value with presence of outliers, in section four, we estimate , Bayes estimation of (θ) based on records value with presence of outliers. These estimators compare in section five.

Bayes estimation of θ with presence of outliers

According to Dixit, Moor and Barnett (1996), we assume that a set of random variables (X_1, X_2, \dots, X_n) represent of the distribution of an infected sampled plant from a plot of plants inoculated with a virus. Some of the observations are derived from the airborne dispersal of the spores and are distributed according to the exponential distribution. The other observations out of n random variables (say k) are present because aphids which are known to be carriers of BYMDV have passed the virus into the plant when the aphids feed on the sap. Theses k (known) aphids are considered to be exponential

distribution. Thus, we assume that the random variables (X_1, X_2, \dots, X_n) are such that k of them are distributed with probability density function (pdf) $g(x, \lambda, \theta)$ as

$$g(x|\lambda, \theta) = \frac{\lambda}{\theta} \exp\left(-\frac{\lambda}{\theta}x\right), \quad x > 0, \quad \lambda > 0, \quad \theta > 0 \quad (1)$$

and the remaining $(n-k)$ random variables are distributed with the following pdf.

$$f(x|\theta) = \frac{1}{\theta} \exp\left(-\frac{x}{\theta}\right), \quad x > 0, \quad \theta > 0 \quad (2)$$

Then the joint pdf of (X_1, X_2, X_n) is

$$f(x|\lambda, \theta) = \frac{k!(n-k)!}{n!} \prod_{i=1}^n f(x_i|\theta) \sum_{j=1}^k \prod_{j=1}^k \frac{g(x_{A_j})}{f(x_{A_j})} \quad (3)$$

$$= \frac{k!(n-k)!}{n!} \frac{1}{\theta^n} \exp\left(-\frac{\sum_{i=1}^n x_i}{\theta}\right) \sum_{j=1}^k \prod_{j=1}^k \frac{\lambda}{\theta} \exp\left(-\frac{\lambda x_{A_j}}{\theta}\right) \frac{1}{\theta} \exp\left(-\frac{x_{A_j}}{\theta}\right) \quad (4)$$

$$= \frac{k!(n-k)!}{n!} \frac{\lambda^k \exp\left(-\frac{n\bar{x}}{\theta}\right)}{\theta^n} \sum_{j=1}^k \exp\left\{-\left(\frac{\lambda}{\theta} - \frac{1}{\theta}\right) \sum_{j=1}^k x_{A_j}\right\} \quad (5)$$

where

$$\sum_{j=1}^k = \sum_{A_1=A_1}^{n-k+1} \sum_{A_2=A_1+1}^{n-k+2} \dots \sum_{A_k=A_{k-1}+1} \quad (6)$$

From (3) the marginal distribution of X is given by

$$f(x) = \frac{k\lambda}{n\theta} \exp\left(-\frac{\lambda x}{\theta}\right) + \frac{(n-k)}{n} \frac{1}{\theta} \exp\left(-\frac{x}{\theta}\right), \quad x > 0 \quad (6)$$

Tele:

E-mail addresses: jabbarinm@yahoo.com

For more details, see Dixit and Nasiri (2001).

Let $\prod(\theta|\alpha, \beta)$ is the prior pdf of θ as

$$\prod(\theta|\alpha, \beta) = \frac{\beta^\alpha}{\Gamma(\alpha)} \left(\frac{1}{\theta}\right)^{\alpha-1} \exp\left(-\frac{\beta}{\theta}\right), \quad \theta > 0.$$

Using (3) and (5), $f(x, \theta|\alpha, \beta, \lambda)$ is given by

$$f(\theta|\alpha, \beta, \lambda) = \frac{k!(n-k)! \beta^\alpha}{n! \Gamma(\alpha)} \frac{\lambda \exp\left(-\frac{n\bar{x} + \beta}{\theta}\right)}{\theta^{n+\alpha-1}} \sum_{j=1}^k \exp\left\{-\left(\frac{\lambda-1}{\theta}\right) \sum_{j=1}^k x_{A_j}\right\} \quad (7)$$

Then the posterior distribution of θ is given by

$$\prod(\theta|x) = \frac{\frac{\exp\left(-\frac{n\bar{x} + \beta}{\theta}\right)}{\theta^{n+\alpha-1}} \sum_{j=1}^k \exp\left\{-\frac{(\lambda-1) \sum_{j=1}^k x_{A_j}}{\theta}\right\}}{\int_0^\infty \frac{\exp\left(-\frac{n\bar{x} + \beta}{\theta}\right)}{\theta^{n+\alpha-1}} \sum_{j=1}^k \exp\left\{-\frac{(\lambda-1) \sum_{j=1}^k x_{A_j}}{\theta}\right\} d\theta}$$

Therefore, by using the notation of I as the integral

$$E(\theta^r|x) = \int_0^\infty \theta^r \prod(\theta|x) d\theta = \frac{1}{I_0} \int_0^\infty \frac{\theta^r \exp\left(-\frac{n\bar{x} + \beta}{\theta}\right)}{\theta^{n+\alpha-1}} \sum_{j=1}^k \exp\left\{-\frac{(\lambda-1) \sum_{j=1}^k x_{A_j}}{\theta}\right\} d\theta \quad (8)$$

$$= \frac{I_r}{I_0} \quad (9)$$

Where

$$I_r = \int_0^\infty \frac{\theta^r \exp\left(-\frac{n\bar{x} + \beta}{\theta}\right)}{\theta^{n+\alpha-1}} \sum_{j=1}^k \exp\left\{-\frac{(\lambda-1) \sum_{j=1}^k x_{A_j}}{\theta}\right\} d\theta \quad (10)$$

For $r = 1$, the bayes estimator of θ is given by

$$\theta^* = E(\theta|x) = \frac{I_1}{I_0}$$

We can use the Lindely approximation to estimate θ

Estimation of θ based on records value with presence of outliers

In this section, we shall be concerned with estimation of the parameter of the exponential distribution based on upper record values with the presence of outliers. Suppose that we observe n upper record values $X_{u(1)} = x_1, X_{u(2)} = x_2, \dots, X_{u(n)} = x_n$ from the exponential model with presence of outliers given by (4). Then the joint pdf of

$$X_{u(1)} = x_1, X_{u(2)} = x_2, \dots, X_{u(n)} = x_n \quad (11)$$

is given as

$$f(x|\theta) = f(x_n, \theta) \prod_{i=1}^{n-1} h(x_i, \theta), \quad x_1 < x_2 < \dots < x_n$$

where

$$h(x_i, \theta) = \frac{f(x_i, \theta)}{1 - F(x_i, \theta)}$$

For more details, refer to Asgharzadeh (2009). So

$$f(x|\theta) = \frac{\left[\frac{k\lambda}{n\theta} \exp\left(-\frac{\lambda x_n}{\theta}\right) + \frac{n-k}{n\theta} \exp\left(-\frac{x_n}{\theta}\right) \right] \prod_{i=1}^{n-1} \left[\frac{k\lambda}{n\theta} \exp\left(-\frac{\lambda x_i}{\theta}\right) + \frac{n-k}{n\theta} \exp\left(-\frac{x_i}{\theta}\right) \right]}{\prod_{i=1}^n \left[\frac{k\lambda}{n\theta} \exp\left(-\frac{\lambda x_i}{\theta}\right) + \frac{n-k}{n\theta} \exp\left(-\frac{x_i}{\theta}\right) \right]} \quad (12)$$

Let $l(\theta) = \frac{d}{d\theta} \ln(f(x|\theta))$, then

$$l(\theta) = \frac{d}{d\theta} \left[\sum_{i=1}^n \ln \left(\frac{k\lambda}{n\theta} \exp\left(-\frac{\lambda x_i}{\theta}\right) + \frac{n-k}{n\theta} \exp\left(-\frac{x_i}{\theta}\right) \right) - \sum_{i=1}^{n-1} \ln \left(\frac{k}{n} \exp\left(-\frac{\lambda x_i}{\theta}\right) + \frac{n-k}{n} \exp\left(-\frac{x_i}{\theta}\right) \right) \right]$$

$$= \frac{1}{\theta^2} \sum_{i=1}^n \frac{k\lambda(\lambda x_i - \theta) \exp\left(-\frac{\lambda x_i}{\theta}\right) + (n-k)(x_i - \theta) \exp\left(-\frac{x_i}{\theta}\right)}{k\lambda \exp\left(-\frac{\lambda x_i}{\theta}\right) + (n-k) \exp\left(-\frac{x_i}{\theta}\right)}$$

$$- \frac{1}{\theta^2} \sum_{i=1}^n \frac{k\lambda x_i \exp\left(-\frac{\lambda x_i}{\theta}\right) + (n-k)x_i \exp\left(-\frac{x_i}{\theta}\right)}{k \exp\left(-\frac{\lambda x_i}{\theta}\right) + (n-k) \exp\left(-\frac{x_i}{\theta}\right)} \quad (13)$$

To estimate θ , consider $\frac{dl(\theta)}{d\theta} = 0$. So

$$g(\theta) = \sum_{i=1}^n \frac{k\lambda(\lambda x_i - \theta) \exp\left(-\frac{\lambda x_i}{\theta}\right) + (n-k)(x_i - \theta) \exp\left(-\frac{x_i}{\theta}\right)}{k\lambda \exp\left(-\frac{\lambda x_i}{\theta}\right) + (n-k) \exp\left(-\frac{x_i}{\theta}\right)} - \sum_{i=1}^n \frac{k\lambda x_i \exp\left(-\frac{\lambda x_i}{\theta}\right) + (n-k)x_i \exp\left(-\frac{x_i}{\theta}\right)}{k \exp\left(-\frac{\lambda x_i}{\theta}\right) + (n-k) \exp\left(-\frac{x_i}{\theta}\right)} \quad (14)$$

Then by using numerical methods, for example, Newton-Raphson

$$\theta_{i+1} = \theta_i - \frac{g'(\theta)}{g(\theta)}$$

where $g'(\theta)$ is derivation of θ

Bayes estimation θ based on records value with the presence of outliers

To obtain Bayes estimation of θ , consider $f(x|\theta)$ as

$$f(x|\theta) = \frac{\left[\frac{k\lambda}{n\theta} \exp\left(-\frac{\lambda x_n}{\theta}\right) + \frac{n-k}{n\theta} \exp\left(-\frac{x_n}{\theta}\right) \right] \prod_{i=1}^{n-1} \left[\frac{k\lambda}{n\theta} \exp\left(-\frac{\lambda x_i}{\theta}\right) + \frac{n-k}{n\theta} \exp\left(-\frac{x_i}{\theta}\right) \right]}{\prod_{i=1}^n \left[\frac{k\lambda}{n\theta} \exp\left(-\frac{\lambda x_i}{\theta}\right) + \frac{n-k}{n\theta} \exp\left(-\frac{x_i}{\theta}\right) \right]} = \frac{\prod_{i=1}^n \left[\frac{k\lambda}{n\theta} \exp\left(-\frac{\lambda x_i}{\theta}\right) + \frac{n-k}{n\theta} \exp\left(-\frac{x_i}{\theta}\right) \right]}{\prod_{i=1}^{n-1} \left[\frac{k}{n} \exp\left(-\frac{\lambda x_i}{\theta}\right) + \frac{n-k}{n} \exp\left(-\frac{x_i}{\theta}\right) \right]}$$

$$\begin{aligned}
 &= \frac{1}{n\theta^n} \frac{\prod_{i=1}^n \left[k\lambda \exp\left(-\frac{\lambda x_i}{\theta}\right) + (n-k)\exp\left(-\frac{x_i}{\theta}\right) \right]}{\prod_{i=1}^{n-1} \left[k \exp\left(-\frac{\lambda x_i}{\theta}\right) + (n-k)\exp\left(-\frac{x_i}{\theta}\right) \right]} \\
 &= \frac{1}{n\theta^n} \frac{\prod_{i=1}^n A_i}{\prod_{i=1}^{n-1} B_i}
 \end{aligned}
 \tag{15}$$

where

$$\begin{aligned}
 A_i &= k\lambda \exp\left(-\frac{\lambda x_i}{\theta}\right) + (n-k)\exp\left(-\frac{x_i}{\theta}\right) \\
 B_i &= k \exp\left(-\frac{\lambda x_i}{\theta}\right) + (n-k)\exp\left(-\frac{x_i}{\theta}\right)
 \end{aligned}$$

For $k = 0$ it is given by Asgharzadeh (2009).

By using (5) and (15), the joint pdf of \underline{x} and θ is given by

$$\prod(\theta, \underline{x}) = \frac{n\beta^\alpha}{\Gamma(\alpha)} \frac{\exp\left(-\frac{\beta}{\theta}\right)}{\theta^{n+\alpha-1}} \frac{\prod_{i=1}^n A_i}{\prod_{i=1}^{n-1} B_i}
 \tag{16}$$

Then

$$\begin{aligned}
 \prod(\theta, \underline{x}) &= \frac{\exp\left(-\frac{\beta}{\theta}\right)}{\theta^{n+\alpha-1}} \frac{\prod_{i=1}^n A_i}{\prod_{i=1}^{n-1} B_i} \\
 \prod(\theta, \underline{x}) &= \frac{\int_0^\infty \frac{\exp\left(-\frac{\beta}{\theta}\right)}{\theta^{n+\alpha-1}} \frac{\prod_{i=1}^n A_i}{\prod_{i=1}^{n-1} B_i} d\theta}
 \end{aligned}
 \tag{17}$$

Hence

$$E(\theta^r | \underline{x}) = \frac{I_r}{I_0}
 \tag{18}$$

where

$$I_r = \int_0^\infty \theta^r \frac{\exp\left(-\frac{\beta}{\theta}\right)}{\theta^{n+\alpha-1}} \frac{\prod_{i=1}^n A_i}{\prod_{i=1}^{n-1} B_i}$$

Numerical Analysis

To get an idea about the Bias and mean square error (MSE) of three type of estimators, we have generated a sample of size $n=5, 6, \dots, 50$, $k=1, 2, 3$, $\alpha=0.5, 2, 5$, $\beta=0.3, 3$, $\theta=0.4, 2, 5$, $\lambda=2, 3$. The results are derived by using R software based on 1000 independent replication (Figure 1-4). In each graph, red color is for Bayes estimator of θ with outliers, green color is for estimation of θ based on records value with the presence of outliers and blue color is to show the behavior of Bayes estimator of θ based on records value in the presence of outliers.

From the graphs, it is seen that the estimator of θ based on records value in the presence of outliers is unbiased and consistent. It has also a better behavior among the other types of estimator for all values of n and the parameters α, β , and λ .




Figure 1. Comparison Bias and MSE of the estimators for $k=1$, $(\alpha, \beta, \lambda, \theta) = (0.5, 0.3, 2, 0.4), (0.5, 3, 3, 2), (0.5, 0.3, 3, 5)$ and $(2, 0.3, 3, 0.4)$, respectively.




Figure 2. Comparison Bias and MSE of the estimators for $k=1$, $(\alpha, \beta, \lambda, \theta) = (2, 3, 2, 2), (5, 3, 3, 0.4), (5, 3, 3, 2)$ and $(5, 3, 2, 5)$, respectively.




Figure 3. Comparison Bias and MSE of the estimators for $k=2$, $(\alpha, \beta, \lambda, \theta) = (0.5, 0.3, 2, 0.4)$, $(0.5, 3, 3, 2)$, $(0.5, 0.3, 3, 5)$ and $(2, 0.3, 3, 0.4)$, respectively




Figure 4. Comparison Bias and MSE of the estimators for $k=2$, $(\alpha, \beta, \lambda, \theta) = (2, 3, 2, 2)$, $(5, 3, 3, 0.4)$, $(5, 3, 3, 2)$ and $(5, 3, 2, 5)$, respectively.




Figure 5. Comparison Bias and MSE of the estimators for $k=3$, $(\alpha, \beta, \lambda, \theta) = (0.5, 0.3, 2, 0.4)$, $(0.5, 3, 3, 2)$, $(0.5, 0.3, 3, 5)$ and $(2, 0.3, 3, 0.4)$, respectively.




Figure 6. Comparison Bias and MSE of the estimators for $k=3$, $(\alpha, \beta, \lambda, \theta) = (2, 3, 2, 2)$, $(5, 3, 3, 0.4)$, $(5, 3, 3, 2)$ and $(5, 3, 2, 5)$, respectively.

**References**

1. Ahmadi, J., Doostparast, M. and Parsian, A. (2005). Estimation and prediction in a two parameter exponential distribution based on k-record value under Linex loss function. *Commun. Statist. Theor. Meth.* 34, 795-805.
2. Asgharzadeh, A. (2009). On Bayesian estimation from exponential distribution based on records. *Journal of the Korean Statistical Society* 38, 125-130.
3. Balasubramanian, K. and Balakrishnan, N. (1992). Estimation for one and two parameter exponential distribution under multiple type II censoring. *Statistische Hefte* 33, 203-216.
4. Chandraeskar, B., Leo Alexander, T. and Balakrishnan, N. (2002). Equivariant estimation for parameters of exponential distribution based type II progressively censored samples. *Commun. Statist. Theor. Meth.* 31(10), 1675-1686.
5. Dixit, U. J., Moore, K. L. and Barnett, V. (1996). On the estimation of the scale parameter of the exponential distribution in the presence of outliers generated from uniform distribution. *Metron* LIV(3-4), 201-211.
6. Dixit, U. J. and Nasiri, P. F. (2001). Estimation of parameters of the exponential distribution in the presence of outliers generated from uniform distribution. *Metron* LIX(3-4), 187--198.
7. Jaheen, Z. F. (2004). Empirical Bayes analysis of record statistics based on Linex and quadratic loss functions. *Computers and Mathematics with Applications* 47, 947-954.
8. Johnson, N. L., Kotz, S. and Balakrishnan, N. (1994). *Continuous univariate distributions: Vol. 1* (2nd ed.). Now York: John Wiley and Sons.