

Increasing aridity reduces soil microbial diversity and abundance in global drylands

Fernando T. Maestre^{a,1}, Manuel Delgado-Baquerizo^b, Thomas C. Jeffries^b, David J. Eldridge^c, Victoria Ochoa^a, Beatriz Gozalo^a, José Luis Quero^d, Miguel García-Gómez^e, Antonio Gallardo^f, Werner Ulrich^g, Matthew A. Bowker^h, Tulio Arredondoⁱ, Claudia Barraza-Zepeda^j, Donaldo Bran^k, Adriana Florentino^l, Juan Gaitán^m, Julio R. Gutiérrez^{i,n,o}, Elisabeth Huber-Sannwaldⁱ, **Mohammad Jankju^p**, Rebecca L. Mau^q, Maria Miriti^r, **Kamal Naseri^p**, Abelardo Ospina^l, Ilan Stavi^s, Deli Wang^t, Natasha N. Woods^r, Xia Yuan^t, Eli Zaady^u, and Brajesh K. Singh^{b,v}

^aDepartamento de Biología y Geología, Física y Química Inorgánica, Escuela Superior de Ciencias Experimentales y Tecnología, Universidad Rey Juan Carlos, 28933 Móstoles, Spain; ^bHawkesbury Institute for the Environment, Western Sydney University, Penrith, 2751, NSW, Australia; ^cSchool of Biological, Earth and Environmental Sciences, University of New South Wales, Sydney, NSW 2052, Australia; ^dDepartamento de Ingeniería Forestal, Escuela Técnica Superior de Ingeniería Agronómica y de Montes, Universidad de Córdoba, Campus de Rabanales, C.P. 14071, Córdoba, Spain; ^eDepartamento de Ingeniería y Morfología del Terreno, Escuela Técnica Superior de Ingenieros de Caminos, Canales y Puertos, Universidad Politécnica de Madrid, 28040 Madrid, Spain; ^fDepartamento de Sistemas Físicos, Químicos y Naturales, Universidad Pablo de Olavide, 41013 Sevilla, Spain; ^gChair of Ecology and Biogeography, Nicolaus Copernicus University in Toruń, Lwowska 1, 87-100 Toruń, Poland; ^hSchool of Forestry, Northern Arizona University, Flagstaff, AZ 86011; ⁱDivision de Ciencias Ambientales, Instituto Potosino de Investigación Científica y Tecnológica, San Luis Potosí, SLP, México; ^jDepartamento de Biología, Facultad de Ciencias, Universidad de La Serena, Benavente 980, Casilla 554, La Serena, Chile; ^kNacional de Tecnología Agropecuaria, Estación Experimental San Carlos de Bariloche, San Carlos de Bariloche, Río Negro, Argentina; ^lInstituto de Edafología, Facultad de Agronomía, Universidad Central de Venezuela, Campus UCV-Maracay, ZP 2101, Estado Aragua, Venezuela; ^mInstituto de Suelos, Centro de Investigación en Recursos Naturales, Instituto Nacional de Tecnología Agropecuaria, Nicolás Repetto y de los Reseros Sin Número, 1686 Hurlingham, Buenos Aires, Argentina; ⁿCentro de Estudios Avanzados en Zonas Áridas (CEAZA), La Serena, Chile; ^oInstituto de Ecología y Biodiversidad, Facultad de Ciencias, Universidad de Chile, Las Palmeras 3425, Casilla 653, Ñuñoa, Santiago, Chile; ^pDepartment of Range and Watershed Management, Ferdowsi University of Mashhad, Mashhad, 91775-1363, Iran; ^qCenter for Ecosystem Science and Society, Northern Arizona University, Flagstaff, AZ 86011; ^rDepartment of Evolution, Ecology and Organismal Biology, Ohio State University, Columbus, OH 43210; ^sDead Sea and Arava Science Center, Yotvata 88840, Israel; ^tInstitute of Grassland Science, Key Laboratory for Vegetation Ecology, Northeast Normal University, Changchun, Jilin 130024, China; ^uDepartment of Natural Resources, Agriculture Research Organization, Ministry of Agriculture, Gilat Research Center, Mobile Post Negev 85280, Israel; and ^vGlobal Centre for Land-Based Innovation, Western Sydney University, Penrith, 2751, NSW, Australia

Edited by William H. Schlesinger, Cary Institute of Ecosystem Studies, Millbrook, NY, and approved November 10, 2015 (received for review August 21, 2015)

Soil bacteria and fungi play key roles in the functioning of terrestrial ecosystems, yet our understanding of their responses to climate change lags significantly behind that of other organisms. This gap in our understanding is particularly true for drylands, which occupy ~41% of Earth's surface, because no global, systematic assessments of the joint diversity of soil bacteria and fungi have been conducted in these environments to date. Here we present results from a study conducted across 80 dryland sites from all continents, except Antarctica, to assess how changes in aridity affect the composition, abundance, and diversity of soil bacteria and fungi. The diversity and abundance of soil bacteria and fungi was reduced as aridity increased. These results were largely driven by the negative impacts of aridity on soil organic carbon content, which positively affected the abundance and diversity of both bacteria and fungi. Aridity promoted shifts in the composition of soil bacteria, with increases in the relative abundance of *Chloroflexi* and α -*Proteobacteria* and decreases in *Acidobacteria* and *Verrucomicrobia*. Contrary to what has been reported by previous continental and global-scale studies, soil pH was not a major driver of bacterial diversity, and fungal communities were dominated by *Ascomycota*. Our results fill a critical gap in our understanding of soil microbial communities in terrestrial ecosystems. They suggest that changes in aridity, such as those predicted by climate-change models, may reduce microbial abundance and diversity, a response that will likely impact the provision of key ecosystem services by global drylands.

bacteria | fungi | climate change | arid | semiarid

Climate change is a major driver of biodiversity loss from local to global scales, in both terrestrial and aquatic ecosystems (1, 2). Given the dependence of crucial ecosystem processes and services on biodiversity (3–5), climate-change-driven biodiversity losses will dramatically alter the functioning of natural ecosystems (4, 6). Key ecosystem processes—such as nutrient cycling, carbon (C) sequestration, and organic matter decomposition—depend on soil bacteria and fungi (7–9). However, we have limited knowledge of the role of climatic factors as drivers of their abundance and diversity at regional and global scales (10–12). This gap in our understanding is particularly true for

drylands, areas with an aridity index (precipitation/potential evapotranspiration ratio) below 0.65 (13), which are among the most sensitive ecosystems to climate change (14). Drylands are expected to expand in global area by 11–23% by 2100 (15), experiencing increased aridity and reduced soil moisture (16). Land degradation and desertification already affect ~250 million people in the developing world (17). Altered climate and the growth of human populations will almost inevitably exacerbate these problems in drylands (14, 17). Because the provisioning of ecosystem services essential for human development (e.g., soil fertility, food, and biomass production) heavily relies on the abundance, composition, and diversity of soil fungi and bacteria (18, 19), it is

Significance

Climate change is increasing the degree of aridity in drylands, which occupy 41% of Earth's surface and support 38% of its population. Soil bacteria and fungi are largely responsible for key ecosystem services, including soil fertility and climate regulation, yet their responses to changes in aridity are poorly understood. Using a field survey conducted in drylands worldwide and DNA-sequencing approaches, we found that increases in aridity reduce the diversity and abundance of soil bacteria and fungi. This study represents an important advancement in our understanding of soil microbial communities and their likely responses to ongoing climate change.

Author contributions: F.T.M., M.D.-B., and B.K.S. designed research; F.T.M., D.J.E., V.O., B.G., J.L.Q., M.G.-G., A.G., M.A.B., T.A., C.B.-Z., D.B., A.F., J.G., J.R.G., E.H.-S., M.J., R.L.M., M.M., K.N., A.O., I.S., D.W., N.N.W., X.Y., and E.Z. performed research; T.C.J., A.G., and W.U. contributed new reagents/analytic tools; F.T.M., M.D.-B., and W.U. analyzed data; and F.T.M., M.D.-B., D.J.E., M.A.B., and B.K.S. wrote the paper.

The authors declare no conflict of interest.

This article is a PNAS Direct Submission.

Data deposition: The primary data have been deposited in figshare, [dx.doi.org/10.6084/m9.figshare.1487693](https://doi.org/10.6084/m9.figshare.1487693). The raw sequence data have been deposited in the GenBank SRA database (BioProject accession no. [PRJNA301533](https://www.ncbi.nlm.nih.gov/bioproject/PRJNA301533)).

¹To whom correspondence should be addressed. Email: fernando.maestre@urjc.es.

This article contains supporting information online at www.pnas.org/lookup/suppl/doi:10.1073/pnas.1516684112/-DCSupplemental.

crucial to understand how changes in aridity affect soil microbial communities. Drylands, however, are poorly represented in global soil bacteria and fungi databases (10–12, 20), and no field study has simultaneously examined how the abundance, composition, and diversity of these organisms vary along aridity gradients in drylands worldwide.

Here, we present a global field study conducted across 80 dryland sites from all continents, except Antarctica (Fig. S1), to assess how changes in aridity, as defined by the aridity index, affect the total abundance and diversity of soil bacteria and fungi and the relative abundance of major bacterial and fungal taxa. The studied ecosystems encompass a wide variety of the climatic, edaphic, and vegetation conditions found in drylands worldwide (*Materials and Methods*). We predict that increases in aridity should reduce the abundance and diversity of soil bacteria and fungi due to the negative relationships typically found between aridity and the availability of resources such as water and C (21), which largely drive soil microbial abundance and activity in drylands (22–24). To test this hypothesis, we characterized bacterial and fungal communities in the soil surface (top 7.5 cm) along natural aridity gradients by using Illumina Miseq profiling of ribosomal genes and internal transcribed spacer (ITS) markers, quantified bacterial and fungal abundances with quantitative PCR (qPCR), and gathered information on multiple biotic and abiotic factors known to influence soil microbes (Fig. S2).

Results and Discussion

Bacterial communities were dominated by *Actinobacteria*, *Proteobacteria*, *Acidobacteria*, and *Planctomycetes* (Fig. S3A), which are common bacterial phyla in soils worldwide (8, 11). The most abundant soil fungi were those from the phylum *Ascomycota*, followed by *Basidiomycota*, *Chytridiomycota*, and *Zygomycetous*

fungi (Fig. S3B). Our findings contrast with a recent global survey of soil fungi (12) reporting a greater dominance of *Basidiomycota* than we observed (56% vs. 22%). These discrepancies probably relate to the low proportion of drylands surveyed in that study (<1% of 350 sites) and underscore our limited understanding of fungal communities inhabiting dryland soils. Minor differences in the dominant bacterial and fungal phyla were observed when assessing differences among major vegetation types, because variation in their relative abundance was <5% when comparing grasslands and woodlands (Fig. S3A and B).

We evaluated the direct relationship between the diversity and abundance of soil bacteria and fungi and aridity using ordinary least-squares (OLS) regression models (*Materials and Methods*). In addition—and to account for possible large-scale spatial non-independence of the sites surveyed (25)—we included the dominant eigenvector of the Euclidean distance matrix of sites as an additional predictor into these models (spatial models; refs. 26 and 27). Increases in aridity were linearly associated with reductions in fungal and bacterial diversity and abundance (Fig. 1). The diversity of several bacterial and fungal taxa also followed this pattern (Figs. S4 and S5), although nonlinear (e.g., *Acidobacteria* and *Proteobacteria*; Fig. S4B and E) and nonsignificant (e.g., *Chloroflexi*; Fig. S4D; and *Zygomycetous* fungi; Fig. S5E) relationships were also found. Aridity also affected the dominance of major bacterial phyla. The relative abundance of *Acidobacteria* declined linearly as aridity increased (Fig. 2A), whereas that of *Chloroflexi* followed the opposite pattern (Fig. 2C). Other bacterial phyla and classes were nonlinearly related to aridity, with peaks in their relative abundance at either low (e.g., *Verrucomicrobia*) or high (e.g., α -*Proteobacteria*) aridity levels (Fig. 2B and D–F). These results can be explained by the different life strategies of bacterial taxa. *Chloroflexi* are known to have multiple adaptations to environmental




Fig. 1. Relationships between aridity and the diversity and abundance of soil bacteria and fungi. The solid lines represent the fitted OLS model. The proportion of variance explained (R^2) of regressions including the dominant eigenvector of the Euclidean distance matrix of sites (for spatial models, see *SI Materials and Methods*), and the differences in the second-order Akaike Information Criterion [ΔAICc] between these models and those shown in the figure, are as follows: $R^2 = 0.197$, $\Delta\text{AICc} = -1.501$ (A); $R^2 = 0.169$, $\Delta\text{AICc} = 1.399$ (B); $R^2 = 0.193$, $\Delta\text{AICc} = -8.879$ (C); and $R^2 = 0.298$, $\Delta\text{AICc} = -15.86$ (D). A negative ΔAICc value indicates that the AICc of the spatial model is lower than that of the nonspatial model.




Fig. 2. Relationships between aridity and the relative abundance (arcsine-transformed proportions) of dominant soil bacterial phyla and classes. The solid lines represent the fitted linear or quadratic OLS model. The proportion of variance explained (R^2) of regressions including the dominant eigenvector of the Euclidean distance matrix of sites, and the differences in the Δ AICc between these models and those shown in the figure are as follows: $R^2 = 0.319$, Δ AICc = -6.288 (A); $R^2 = 0.164$, Δ AICc = -0.173 (B); $R^2 = 0.112$, Δ AICc = 1.047 (C); $R^2 = 0.245$, Δ AICc = -1.376 (D); $R^2 = 0.341$, Δ AICc = -12.93 (E); and $R^2 = 0.250$, Δ AICc = -6.234 (F). The rest of the legend is as in Fig. 1.

harshness (28), whereas *Acidobacteria* and *Verrucomicrobia* typically show opportunistic responses to short-term changes in water availability, with rapid declines and increases in ribosomal synthesis during drought and after soil rewetting, respectively (24). The relative abundance of major fungal phyla did not change with aridity (R^2 of linear/logarithmic/quadratic regression < 0.07 , $P > 0.08$ in all cases). These findings are consistent with those from studies showing that the relative abundance of fungal phyla remained largely unchanged with soil desiccation during a summer drought (24). Spatial models that included aridity were superior predictors of changes in the abundance of fungi and bacteria (Fig. 1), in the diversity of *Actinobacteria* and *Acidobacteria* (Fig. S4), and in the relative abundance of *Acidobacteria*, *Verrucomicrobia*, and δ -*Proteobacteria* (Fig. 2). These results are likely to be driven by the strong relationships between aridity and latitude/longitude found in our database (Fig. 3A).

To further investigate the direct and indirect effects of aridity on soil bacteria and fungi, we generated structural equation models (SEMs) based on the known effects and relationships among aridity and other key drivers of the diversity and abundance of these microorganisms (mean diurnal temperature range, plant

cover, soil pH, and organic C content; Fig. S2). Latitude and longitude were also included in our models, given their effects on microbial abundance (Fig. 2C and D) and on the rest of biotic and abiotic variables evaluated (Fig. 3A and B). Our models explained between 34% and 49% of the variance found in microbial diversity and abundance among the sites surveyed (Fig. 3A and B). Aridity indirectly impacted the diversity and abundance of soil bacteria and fungi by strongly affecting soil pH, soil organic C content, and total plant cover (Fig. 3C and D). Organic C content had a direct positive effect on the diversity and abundance of both bacteria and fungi (Fig. 3A and B and Fig. S6) and was the strongest predictor of such community attributes in the case of bacteria (Fig. 3C). These results suggest that soil microbial communities are limited by C in dryland soils (22, 23) and align with studies from polar regions, indicating that soil C content is a major driver of the diversity of soil bacteria and fungi (29). Our findings also mimic observed global patterns in microbial biomass C, which have been found to increase in tandem with soil C contents from southern to northern latitudes (30, 31). The negative effect of mean diurnal temperature range (MDR) on bacterial and fungal abundance is likely due to (i) increases in physiological stress

which does not account for the different effects of human activities and other climate-change drivers that may change with aridity. For example, if the introduction of exotic, invasive species by humans in ecosystems occurs along with increasing aridity, some of the observed links between aridity, soil organic C content, and microbial diversity could break down (36). Increases in water use efficiency (WUE) due to elevated [CO₂] may enhance overall plant growth and soil C fixation in drylands (37), which could mitigate the reduction of microbial abundance and diversity expected as aridity increases. Whether this enhancement of WUE can compensate for the detrimental effects of increased aridity on water availability and plant growth is largely unknown. A recent study showed that increased aridity over the last four decades was responsible for a sustained decline in plant productivity, regardless of CO₂-induced increases in WUE during this period (38).

Conclusions

Here we show that increases in aridity, such as those predicted by climate-change models (15, 16), reduce the diversity and abundance of soil bacteria and fungi in drylands, the largest biome on Earth (39). These responses are mainly driven by reductions in soil organic C content associated with increases in aridity and, in the case of microbial abundance, with increases in diurnal temperature variations. Soil pH affected both the abundance of fungi and the relative abundance of major bacterial phyla and classes, but had no effect on the diversity of soil bacteria. Unlike reports of previous global surveys from terrestrial ecosystems (12), fungal communities in dryland soils were dominated by *Ascomycota*. These findings highlight the unique features of soil microbial communities in drylands. Both the community structure and relative importance of environmental factors driving variation in microbial communities in global drylands differ from previous records from other terrestrial ecosystems. Our results fill a critical gap in our understanding of microbial community structure in global drylands and provide additional insights into how soil microbial communities may respond to climate change. Ecosystem models are beginning to incorporate information on microbial abundance, composition, and diversity, which is needed to improve predictions of soil C stocks and their dynamics (31), and the links between aridity, soil organic C content, and these microbial community attributes shown here can be used to refine and validate them.

Materials and Methods

Complete documentation of the study sites, field survey, sample collection, and laboratory procedures, as well as additional details on the statistical analyses are provided in *SI Materials and Methods*.

Field data were collected from 80 dryland sites selected to represent a wide range of the environmental and biotic characteristics of global drylands (Fig. S1; figshare DOI 10.6084/m9.figshare.1487693). At each site, the cover of

perennial vegetation was measured by using the line-intercept method along four 30-m-long transects (5). Replicated soil samples (0- to 7.5-cm depth) were randomly taken under the canopy of the dominant perennial plant species and in open areas devoid of perennial vegetation (10–15 samples per site). After field collection, a fraction of the soil samples was immediately frozen at –20 °C for microbial analyses. These analyses were conducted on composite samples of each microsite (open and vegetated areas) and site. Soil DNA was extracted from 0.5 g of defrosted soil samples by using the Powersoil DNA Isolation Kit (Mo Bio Laboratories). qPCR reactions were performed in triplicate by using 96-well plates on an ABI 7300 Real-Time PCR (Applied Biosystems). The bacterial 16S-rRNA genes and fungal ITS were amplified with the Eub 338-Eub 518 and ITS 1-5.8S primer sets (40). After qPCR analyses, the extracted DNA samples were frozen and shipped to the Next Generation Genome Sequencing Facility of Western Sydney University, where they were defrosted and analyzed by using the Illumina MiSeq platform (41) and the 341F/805R (bacteria) and FIT57/ITS4 (fungi) primer sets (42, 43). Initial sequence processing and diversity analyses for both bacterial 16S rDNA and fungal ITS genes were conducted as described in *SI Materials and Methods*.

Before numerical and statistical analyses, all soil and microbial variables used in this study were averaged to obtain site-level estimates by using the mean values observed in bare ground and vegetated areas, weighted by their respective cover at each site (5). We first modeled the relationships between aridity and the abundance and diversity of bacteria and fungi using either linear or curvilinear (quadratic) regressions. The aridity [1 – aridity index (AI), where AI is precipitation/potential evapotranspiration] of each site was obtained by using data from ref. 44. Similarly, we explored the relationships between aridity and the abundance and diversity of main bacterial and fungal phyla and classes, as well as between soil organic C content/pH and these taxa using regression and partial correlation analyses as described in *SI Materials and Methods*. To determine the mechanisms underlying the observed effects of aridity on microbial abundance and diversity, we used SEM (45). This approach tests the plausibility of a causal model encompassing a set of a priori hypotheses (Fig. S2). Our a priori model included spatial structure (latitude and longitude), aridity, mean diurnal temperature range (obtained from ref. 46), soil pH, organic C, and total plant cover as predictors of the total amount (as measured with qPCR) and diversity (Shannon index) of both bacteria and fungi. SEM analyses were conducted as described in *SI Materials and Methods* by using AMOS (Version 18.0; Amos Development). All of the data used in the primary analyses are available from figshare (DOI 10.6084/m9.figshare.1487693).

ACKNOWLEDGMENTS. We thank D. Encinar for his help with the laboratory analyses and data management; Stefan Hempel for his help with the nomenclature of fungi; and Matthias C. Rillig, Mark Bradford, Nicholas J. Gotelli, Santiago Soliveres, Stavros Veresoglou, and Jeff Powell for revising previous versions of the manuscript. This research is supported by the European Research Council (ERC) under the European Community's Seventh Framework Programme FP7/2007-2013/ERC Grant Agreement 242658 (BIOCOM); by Spanish Ministry of Economy and Competitiveness BIOMOD Project CGL2013-44661-R; and by Australian Research Council Project DP13010484. F.T.M. was supported by Salvador de Madariaga program of the Spanish Ministry of Education, Culture and Sports Grant PRX14/00225; the Research Exchange Program of the Hawkesbury Institute for the Environment; and a Research Award granted by the Alexander Von Humboldt Foundation. J.R.G. was supported by Iniciativa Científica Milenio PO5-002 (MIDEPLAN) and Comisión Nacional de Investigación Científica y Tecnológica PFB-23.

- Bellard C, Bertelsmeier C, Leadley P, Thuiller W, Courchamp F (2012) Impacts of climate change on the future of biodiversity. *Ecol Lett* 15(4):365–377.
- Bálint M, et al. (2011) Cryptic biodiversity loss linked to global climate change. *Nat Clim Chang* 1(6):313–318.
- Isbell F, et al. (2011) High plant diversity is needed to maintain ecosystem services. *Nature* 477(7363):199–202.
- Cardinale BJ, et al. (2012) Biodiversity loss and its impact on humanity. *Nature* 486(7401):59–67.
- Maestre FT, et al. (2012) Plant species richness and ecosystem multifunctionality in global drylands. *Science* 335(6065):214–218.
- Peñuelas J, et al. (2013) Evidence of current impact of climate change on life: A walk from genes to the biosphere. *Glob Change Biol* 19(8):2303–2338.
- Trivedi P, Anderson IC, Singh BK (2013) Microbial modulators of soil carbon storage: Integrating genomic and metabolic knowledge for global prediction. *Trends Microbiol* 21(12):641–651.
- Fierer N, et al. (2013) Reconstructing the microbial diversity and function of pre-agricultural tallgrass prairie soils in the United States. *Science* 342(6158):621–624.
- Zhou J, et al. (2011) Microbial mediation of carbon-cycle feedbacks to climate warming. *Nat Clim Chang* 2(2):106–110.
- Serna-Chavez HM, Fierer N, Van Bodegom PM (2013) Global drivers and patterns of microbial abundance in soil. *Glob Ecol Biogeogr* 22(10):1162–1172.
- Ramirez KS, et al. (2014) Biogeographic patterns in below-ground diversity in New York City's Central Park are similar to those observed globally. *Proc Biol Sci* 281(1795):20141988.
- Tedesco L, et al. (2014) Fungal biogeography. Global diversity and geography of soil fungi. *Science* 346(6213):1256688.
- Safirel U, Adeel Z (2005) Dryland systems. *Ecosystems and Human Well-Being: Current State and Trends*, eds Hassan R, Scholes R, Neville A (Island, Washington, DC), Vol 1, pp 623–662.
- Maestre FT, Salguero-Gómez R, Quero JL (2012) It is getting hotter in here: Determining and projecting the impacts of global environmental change on drylands. *Philos Trans R Soc Lond B Biol Sci* 367(1606):3062–3075.
- Huang J, Yu H, Guan X, Wang G, Guo R (October 26, 2015) Accelerated dryland expansion under climate change. *Nat Clim Chang*, 10.1038/nclimate2837.
- Fu Q, Feng S (2014) Responses of terrestrial aridity to global warming. *J Geophys Res Atmos* 119:7863–7875.
- Reynolds JF, et al. (2007) Global desertification: Building a science for dryland development. *Science* 316(5826):847–851.
- Wall DH, et al. (eds) (2012) *Soil Ecology and Ecosystem Services* (Oxford Univ Press, Oxford).
- Pointing SB, Belpaen J (2012) Microbial colonization and controls in dryland systems. *Nat Rev Microbiol* 10(8):551–562.

20. Lauber CL, Hamady M, Knight R, Fierer N (2009) Pyrosequencing-based assessment of soil pH as a predictor of soil bacterial community structure at the continental scale. *Appl Environ Microbiol* 75(15):5111–5120.
21. Delgado-Baquerizo M, et al. (2013) Decoupling of soil nutrient cycles as a function of aridity in global drylands. *Nature* 502(7473):672–676.
22. Manzoni S, Schimel JP, Porporato A (2012) Responses of soil microbial communities to water stress: Results from a meta-analysis. *Ecology* 93(4):930–938.
23. Angel R, Pasternak Z, Soares MIM, Conrad R, Gillor O (2013) Active and total prokaryotic communities in dryland soils. *FEMS Microbiol Ecol* 86(1):130–138.
24. Barnard RL, Osborne CA, Firestone MK (2013) Responses of soil bacterial and fungal communities to extreme desiccation and rewetting. *ISME J* 7(11):2229–2241.
25. Hawkins BA (2012) Eight (and a half) deadly sins of spatial analysis. *J Biogeogr* 39(1):1–9.
26. Dormann CF, et al. (2007) Methods to account for spatial autocorrelation in the analysis of species distributional data: A review. *Ecography* 30(5):609–628.
27. Ulrich W, et al. (2014) Climate and soil attributes determine plant species turnover in global drylands. *J Biogeogr* 41(12):2307–2319.
28. Battistuzzi FU, Hedges SB (2009) A major clade of prokaryotes with ancient adaptations to life on land. *Mol Biol Evol* 26(2):335–343.
29. Siciliano SD (2014) Soil fertility is associated with fungal and bacterial richness, whereas pH is associated with community composition in polar soil microbial communities. *Soil Biol Biochem* 78:10–20.
30. Xu X, Thornton PE, Post WM (2013) A global analysis of soil microbial biomass carbon, nitrogen and phosphorus in terrestrial ecosystems. *Glob Ecol Biogeogr* 22(6):737–749.
31. Wieder WR, Bonan GB, Allison SD (2013) Global soil carbon projections are improved by modelling microbial processes. *Nat Clim Chang* 3(10):909–912.
32. Van Gestel NC, Schwilk DW, Tissue DT, Zak JC (2011) Reductions in daily soil temperature variability increase soil microbial biomass C and decrease soil N availability in the Chihuahuan Desert: Potential implications for ecosystem C and N fluxes. *Glob Change Biol* 17(12):3564–3576.
33. Fierer N, Jackson RB (2006) The diversity and biogeography of soil bacterial communities. *Proc Natl Acad Sci USA* 103(3):626–631.
34. FAO/IIASA/ISRIC/ISSCAS/JRC (2012) *Harmonized World Soil Database* (FAO, Rome), Version 1.2.
35. Rousk J, et al. (2010) Soil bacterial and fungal communities across a pH gradient in an arable soil. *ISME J* 4(10):1340–1351.
36. Zou J, Rogers WE, DeWalt SJ, Siemann E (2006) The effect of Chinese tallow tree (*Sapium sebiferum*) ecotype on soil-plant system carbon and nitrogen processes. *Oecologia* 150(2):272–281.
37. Evans RD, et al. (2014) Greater ecosystem carbon in the Mojave Desert after ten years exposure to elevated CO₂. *Nat Clim Chang* 4(5):394–397.
38. Brookshire ENJ, Weaver T (2015) Long-term decline in grassland productivity driven by increasing dryness. *Nat Commun* 6:7148.
39. Schimel DS (2010) Climate: Drylands in the Earth system. *Science* 327(5964):418–419.
40. Evans SE, Wallenstein MD (2011) Soil microbial community response to drying and rewetting stress: Does historical precipitation regime matter? *Biogeochemistry* 109(1–3):101–116.
41. Caporaso JG, et al. (2012) Ultra-high-throughput microbial community analysis on the Illumina HiSeq and MiSeq platforms. *ISME J* 6(8):1621–1624.
42. Herlemann DP, et al. (2011) Transitions in bacterial communities along the 2000 km salinity gradient of the Baltic Sea. *ISME J* 5(10):1571–1579.
43. Ihrmark K, et al. (2012) New primers to amplify the fungal ITS2 region—evaluation by 454-sequencing of artificial and natural communities. *FEMS Microbiol Ecol* 82(3):666–677.
44. Trabucco A, Zomer RJ (2009) Global Aridity Index (global-aridity) and global potential evapo-transpiration (global-PET) geospatial database. CGIAR Consortium for Spatial Information. The CGIAR-CSI GeoPortal. Available at www.cgiar-csi.org/data/global-aridity-and-pet-database.
45. Grace JB (2006) *Structural Equation Modeling and Natural Systems* (Cambridge Univ Press, Cambridge, U.K.).
46. Hijmans RJ, et al. (2005) Very high resolution interpolated climate surfaces for global areas. *Int J Clim* 25(15):1965–1978.
47. Caporaso JG, et al. (2010) QIIME allows analysis of high-throughput community sequencing data. *Nat Methods* 7(5):335–336.
48. Magoč T, Salzberg SL (2011) FLASH: Fast length adjustment of short reads to improve genome assemblies. *Bioinformatics* 27(21):2957–2963.
49. Aronesty E (2011) Command-line tools for processing biological sequencing data. *ea-utils: Fast Q Processing Utilities*. Available at code.google.com/p/ea-utils.
50. Edgar RC (2010) Search and clustering orders of magnitude faster than BLAST. *Bioinformatics* 26(19):2460–2461.
51. Edgar RC, Haas BJ, Clemente JC, Quince C, Knight R (2011) UCHIME improves sensitivity and speed of chimera detection. *Bioinformatics* 27(16):2194–2200.
52. Cole JR, et al. (2005) The Ribosomal Database Project (RDP-II): Sequences and tools for high-throughput rRNA analysis. *Nucleic Acids Res* 33(Database issue, Suppl 1):D294–D296.
53. DeSantis TZ, et al. (2006) Greengenes, a chimera-checked 16S rRNA gene database and workbench compatible with ARB. *Appl Environ Microbiol* 72(7):5069–5072.
54. McDonald D, et al. (2012) An improved Greengenes taxonomy with explicit ranks for ecological and evolutionary analyses of bacteria and archaea. *ISME J* 6(3):610–618.
55. Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ (1990) Basic local alignment search tool. *J Mol Biol* 215(3):403–410.
56. Kõljalg U, et al. (2013) Towards a unified paradigm for sequence-based identification of fungi. *Mol Ecol* 22(21):5271–5277.
57. McLaughlin DJ, Spatafora JW (eds) (2014) *The Mycota. VII Systematics and Evolution Part A* (Springer, Heidelberg), 2nd Ed.
58. Haegeman B, et al. (2013) Robust estimation of microbial diversity in theory and in practice. *ISME J* 7(6):1092–1101.
59. Sugiura N (1978) Further analysis of the data by Akaike's information criterion and the finite corrections. *Commun Stat Theory Methods* 7(1):13–26.
60. Burnham KP, Anderson DR (2002) *Model Selection and Multimodel Inference. A Practical Information-Theoretical Approach* (Springer, Heidelberg).
61. Good PI (2004) *Permutation, Parametric and Bootstrap Tests of Hypotheses* (Springer, New York), 3rd Ed.
62. Lange K (2010) *Numerical Analysis for Statisticians* (Springer, New York).
63. Boos DD, Stefanski LA (2013) *Essential Statistical Inference* (Springer, New York).
64. Anderson MJ, Legendre P (1999) An empirical comparison of permutation methods for tests of partial regression coefficients in a linear model. *J Stat Comput Simul* 62(3):271–303.
65. Hammer Ø, Harper DAT, Ryan PD (2001) PAST: Paleontological statistics software package for education and data analysis. *Palaeontol Electronica* 4:1–9.
66. Schermelleh-Engel K, et al. (2003) Evaluating the fit of structural equation models, tests of significance descriptive goodness-of-fit measures. *Methods Psychol Res Online* 8(2):23–74.
67. Delgado-Baquerizo M, Gallardo A, Wallenstein MD, Maestre FT (2013) Vascular plants mediate the effects of aridity and soil properties on ammonia-oxidizing bacteria and archaea. *FEMS Microbiol Ecol* 85(2):273–282.
68. Liu X, et al. (2006) Temporal trends and variability of daily maximum and minimum, extreme temperature events, and growing season length over the eastern and central Tibetan Plateau during 1961–2003. *J Geophys Res Atmos* 111(19):1–19.
69. Elagib NA, Addin Abdu AS (1997) Climate variability and aridity in Bahrain. *J Arid Environ* 36(3):405–419.
70. Zhou L, Dickinson RE, Tian Y, Vose RS, Dai Y (2007) Impact of vegetation removal and soil aridation on diurnal temperature range in a semiarid region: Application to the Sahel. *Proc Natl Acad Sci USA* 104(46):17937–17942.
71. Bell C, McIntyre N, Cox S, Tissue D, Zak J (2008) Soil microbial responses to temporal variations of moisture and temperature in a chihuahuan desert grassland. *Microb Ecol* 56(1):153–167.
72. Wen-Jie W, et al. (2011) Changes in soil organic carbon, nitrogen, pH and bulk density with the development of larch (*Larix gmelinii*) plantations in China. *Glob Change Biol* 17(8):2657–2676.
73. Fierer N, Lennon JT (2011) The generation and maintenance of diversity in microbial communities. *Am J Bot* 98(3):439–448.
74. Schlesinger WH (1996) *Biogeochemistry, an Analysis of Global Change* (Academic, San Diego).

Supporting Information

Maestre et al. 10.1073/pnas.1516684112

SI Materials and Methods

Field Survey and Sample Collection. Field data were collected from 80 dryland sites located in 12 countries from all continents except Antarctica (Fig. S1). Data collection took place between June 2006 and January 2011, and all of the sites were surveyed by using a standardized sampling protocol. At each site, a 30-m × 30-m plot representative of the dominant vegetation was established, and the cover of perennial vegetation was measured by using the line-intercept method along four 30-m-long transects (5). The coordinates and elevation of each plot were recorded in situ with a portable global positioning system and were standardized to the WGS84 ellipsoid for visualization and analyses by using Google Earth (<https://www.google.com/earth/>). We estimated the degree of aridity of each site by obtaining its aridity index (precipitation/potential evapotranspiration) using data interpolations provided by Worldclim (44, 46). To facilitate the interpretation of our results, we used 1-AI as our surrogate of aridity (21). This index increases with decreasing annual mean precipitation in the sites studied (Spearman $\rho = -0.829$, $n = 80$, $P < 0.001$).

Soils were sampled during the dry season in most of the sites using a stratified random procedure. At each plot, five 50-cm × 50-cm quadrats were randomly placed under the canopy of the dominant perennial vegetation element and in open areas devoid of perennial vegetation. A composite sample consisting of five 145-cm³ soil cores (0- to 7.5-cm depth) was collected from each quadrat, bulked, and homogenized in the field. When more than one dominant plant species was found, samples were also collected under the canopies of five randomly selected individuals of the codominant species. Thus, the number of soil samples varied between 10 and 15 per site. To avoid problems associated with the use of multiple laboratories when analyzing the soils from different sites, and to facilitate the comparison of results between them, dried and frozen soil samples from all of the countries were shipped to Spain for laboratory analyses.

Microbial Analyses. The extracted DNA was of high quality, with ratios of A260/A230 and A260/A280 above 1.5 and 1.8, respectively. Initial sequence processing and diversity analyses for both bacterial 16S rDNA and fungal ITS genes were conducted by using the QIIME package (47). Initially, low-quality regions ($Q < 20$) were trimmed from the 5' end of sequences, and paired ends were joined with FLASH (48) for 16S rDNA sequences and Fastq-join (49) for ITS reads. Sequences were demultiplexed, and a further round of quality control was conducted to remove sequences containing ambiguous bases (N) and reads containing bases with a quality score < 25 . Chimeric 16S rDNA sequences were detected by using the UCHIME algorithm from the USEARCH package (50, 51) implemented within VSEARCH (<https://github.com/torognes/vsearch>). The RDP training dataset (Version 9; ref. 52) was used as a reference for chimera detection, as recommended by the UCHIME documentation. De novo (abundance-based) chimera detection was used for ITS data using USEARCH (50). The remaining high-quality chimera-free sequences were used for downstream analysis. Operational taxonomic units (OTUs) were defined as clusters of 97% sequence similarity using UCLUST (50). Taxonomy was assigned using UCLUST (50) against the Greengenes database (Version 13_850) for 16S rDNA OTUs (53, 54). For fungal ITS sequences, taxonomy was assigned by using BLAST (55) against the UNITE database (Version 6.9.7; ref. 56) ($E < 10^{-5}$). This database, however, considers the *Zygomycota* as a phylum, whereas the most recent taxonomical references no longer do so (57).

We thus refer to Zygomycetous fungi when referring to the different taxonomical units formerly included in the *Zygomycota* (57). The resultant OTU abundance tables for both primer sets were filtered to remove singletons and rarefied to an even number of sequences per samples to ensure an equal sampling depth (11,925 and 17,000 for 16S rDNA and ITS, respectively). Shannon diversity was calculated on these rarefied OTU tables by using QIIME (47). We estimated diversity using this metric because it has been recommended when quantifying and comparing microbial diversity (58). The number of bacterial sequences obtained from two of the sites surveyed was too low to estimate microbial diversity accurately, so they were not used in further analyses.

Assessing Relationships Between Microbial Community Attributes, Aridity, and Soil Variables. We explored the relationships between aridity and the abundance and diversity of main bacterial and fungal phyla and classes, as well as between soil organic C content/pH and these taxa by using OLS regression analyses. We used the AICc (59) to compare linear, logarithmic, and quadratic models; when differences in the AICc ($\Delta AICc$) values of these models were < 2 , the linear model was selected (60). To account for possible large-scale spatial nonindependence of sample sites (25), we also included the dominant eigenvector of the Euclidean distance matrix of sites as an additional predictor into the OLS regressions (26, 27). Results of OLS regressions with and without this spatial predictor were compared by using AICc; $\Delta AICc$ values of spatial vs. nonspatial models > 4 suggest that the models are clearly different (60). Some of the regressions conducted did not fulfill some of the assumptions of OLS regression (normality and homogeneity of variance). Thus, we conducted additional analyses using permutation tests to obtain P values (61), because these provide superior type I error control when assumptions of normality and homogeneity of variance are violated (62–64). Nonparametric significance levels were obtained from 10,000 permutations each. In all of the regressions conducted, the differences between OLS- and permutation-obtained P values were minimal and did not change either the statistical significance of the regression or the interpretation of the results. Hence, for simplicity, we report in the main text results of OLS regression. We conducted partial correlation analyses between soil pH and the relative abundance of major bacteria phyla and classes controlling for the effects of aridity and soil organic C content. OLS regression and nonparametric permutation analyses were conducted with Sigma Plot (Version 11; Systat) and Past (Version 3.09) (65), respectively. Spatial regression models were calculated by using Statistica (Version 12; StatSoft). Partial correlation analyses were conducted by using SPSS (Version 21; IBM).

Evaluating Direct and Indirect Effects of Aridity on Microbial Community Attributes: SEM. Our a priori SEM is based on the known effects and relationships among key drivers of bacterial and fungal diversity and abundance (Fig. S2). This model included: spatial structure (latitude and longitude), aridity, mean diurnal temperature range, soil pH, organic C, and total plant cover as predictors of the total amount (as measured with qPCR) and diversity (Shannon index) of both bacteria and fungi. The mean diurnal temperature range was obtained as the average of monthly differences between maximum and minimum air temperatures, as provided by Worldclim (46). Before modeling, we examined the distributions of all of our variables and tested their normality. Bacterial and fungal abundance were log₁₀-transformed.

When these data manipulations were complete, we parameterized our model using our dataset and tested its overall goodness of fit. There is no single universally accepted test of overall goodness of fit for SEMs, so we used multiple goodness-of-fit criteria. We used the χ^2 test (the model has a good fit when χ^2 is low and probability value is high, traditionally $P > 0.05$; ref. 66) and the root mean square error of approximation (RMSEA; the model has a good fit when RMSEA is near 0 and the probability is high, traditionally $P > 0.05$; ref. 66). Additionally, and because some variables were not normal, we confirmed the fit of the model using the Bollen–Stine bootstrap test (the model has a good fit when bootstrap P is high, traditionally $P > 0.05$; ref. 66). Our a priori model attained an acceptable fit by all criteria (Fig. 3), and thus no post hoc alterations were made. After attaining a satisfactory model fit, we introduced composite variables into our model. The use of composite variables does not alter the underlying model, but collapses the effects of multiple conceptually related variables into a single composite effect, aiding interpretation of model results (45). Both latitude and longitude were necessary to represent the spatial distribution of the dryland sites and thus were included as a composite variable.

With a reasonable model fit, and composite variables constructed, we were free to interpret the path coefficients of the model and their associated P values. A path coefficient is analogous to a partial

correlation coefficient and describes the strength and sign of the relationships between two variables (45). Because some of the variables introduced did not follow a normal distribution, the probability of a path coefficient to differ from zero was tested by using bootstrap tests (45). Bootstrapping is preferred to the classical maximum-likelihood estimation in these cases because, in bootstrapping, probability assessments are not based on an assumption that the data match a particular theoretical distribution. Thus, data are randomly sampled with replacement to arrive at estimates of SEs that are empirically associated with the distribution of the data in the sample (45).

Another important capability of SEM is its ability to partition direct and indirect effects that one variable may have on another and estimate the strengths of these multiple effects. To aid final interpretation in light of this ability of SEM, we calculated the standardized total effects of latitude, longitude, aridity, mean diurnal temperature range, plant cover, soil organic C content, and soil pH (and also total microbial abundance in the case of diversity). The net influence that one variable has upon another was calculated by summing all direct and indirect pathways between the two variables. If the model fits the data well, the total effect should approximate the bivariate correlation coefficient for that pair of variables (45).




Fig. S1. Location of the 80 sites used in this study. Some of them overlap and are thus indistinguishable. Exact locations and additional site characteristics are provided in figshare (DOI 10.6084/m9.figshare.1487693).




Fig. S4. Relationships between aridity and the diversity (Shannon index, in bits) of dominant bacterial phyla in the ecosystems studied. The proportion of variance explained (R^2) of regressions including the dominant eigenvector of the Euclidean distance matrix of sites (spatial models) and the ΔAICc between these models and those shown in the figure, are as follows: $R^2 = 0.267$, $\Delta\text{AICc} = -12.301$ (A); $R^2 = 0.269$, $\Delta\text{AICc} = -4.991$ (B); $R^2 = 0.115$, $\Delta\text{AICc} = 2.072$ (C); $R^2 = 0.333$, $\Delta\text{AICc} = -2.477$ (E); and $R^2 = 0.068$, $\Delta\text{AICc} = 1.706$ (F). The relationship presented in D is nonsignificant ($P > 0.05$), hence no regression model is shown. A negative ΔAICc value indicates that the AICc of the spatial model is lower than that of the nonspatial model.




Fig. S9. Relationships between soil pH and organic C content and the relative abundance (arcsine-transformed proportions) of dominant fungal phyla in the ecosystems studied. The solid lines represent the fitted OLS model. The proportion of variance explained (R^2) of regressions including the dominant eigenvector of the Euclidean distance matrix of sites and $\Delta AICc$ between these models and those shown in the figure, are as follows: $R^2 = 0.108$, $\Delta AICc = -0.997$ (D); $R^2 = 0.066$, $\Delta AICc = 2.177$ (E); and $R^2 = 0.129$, $\Delta AICc = -4.016$ (F). The relationships presented in A, B, and C are nonsignificant ($P > 0.05$), hence no regression models are shown. The rest of the legend is as in Fig. S4.