

harvested the next morning (AM), in 2010 on June 13/14, 19/20 and 26/27 for EB, LB and EF, respectively. After 4 field days, these hays ($n=6$) were baled in a factorial arrangement. Advancing from EB to EF, the DM content of fresh alfalfa increased ($P<0.01$) and delaying the time of cutting in the day increased its DM content ($P<0.01$). With advancing maturity, leaf content and leaf/stem ratio decreased ($P<0.01$). These traits increased at PM cutting compared with AM cutting ($P<0.01$). Across maturity stages ash and CP content decreased and total carbohydrate (TCHO), NDF, ADF and cellulose content increased ($P<0.05$). These components were similar between alfalfa harvested at sunset and sunrise. Fat content was similar among the maturity stages and between PM and AM harvests ($P<0.05$). Alfalfa harvested at sunset had a higher true potentially degradable protein (PB) and lower buffer soluble protein (BSP) compared with sunrise harvests ($P<0.05$). These protein fractions were similar among three stages of maturity. There was an interaction between stage of maturity and cutting time for dry matter digestibility (DMD) after 24 h and 48 h of in situ incubation ($P<0.05$), which were both highest for alfalfa harvest at EBPM. Our findings confirm, as previously reported, that with advancing maturity ash, CP and leaf contents decrease and NDF, ADF and TCHO contents increase. Alfalfa harvest at sunset had higher DM, leaf and PB contents and lower BSP content. Results from the current study indicate that early bud stage at sunset is the optimum time to harvest alfalfa.

Key words: Alfalfa hay, sunset and sunrise, botanical traits, chemical composition and dry matter digestibility, stage of maturity

Alfalfa at three stages of maturity harvested at sunset and sunrise: effects on dairy cattle performance. Mojtaba Yari^{1†}, Reza Valizadeh¹, Abbas Ali Naserian¹, Gholam Reza Ghorbani², Parviz Rezvani Moghaddam¹, Arjan Jonker^{3*†}, and Peiqiang Yu³. ¹Department of Plant Science, Ferdowsi University of Mashhad, Mashhad; Iran; ²Department of Animal Science, Isfahan University of Technology, Isfahan, Iran; ³Department of Animal and Poultry Science, University of Saskatchewan, Saskatoon, SK, Canada.

Information is lacking on the optimum time to harvest alfalfa to obtain high-quality forage for dairy cattle. The objective of this study was to determine optimum maturity stage and time of the day to harvest alfalfa in the east of Iran. The third cut of a second-year alfalfa field (4.2 ha) was divided into three plots (1.4 ha) with each plot assigned to one of the following three stages of maturity: early bud (EB), late bud (LB) and early flowering (EF, local commercial harvesting stage). Half of each plot was cut at sunset (PM) after a sunny day and the rest was cut the following morning (AM) in

2010 on June 13, 14, and 19, 20 and 26, 27 for EB, LB and EF, respectively, and hays ($N=6$) baled after 4 d. Eighteen Holstein-Frisian dairy cows (66.3 d in milk and 43 kg milk d^{-1} pretrial) were used in a cyclic changeover design with three 21-d periods in a 3×2 factorial arrangement. Total mixed rations consisted of % DM of 20% experimental alfalfa hays plus 15% barley silage and 65% concentrates delivered twice daily. Organic matter intake of cows consuming EBAM alfalfa hay was higher than those consuming EFAM ($P<0.05$). With advancing maturity stages, net energy intake of cows tended to decrease linearly ($P<0.1$). In vivo total tract DM, OM, CP and NDF digestibility and daily DM, OM and NDF digestible intake were highest in EFPM alfalfa hay and were lowest for LFAM alfalfa hay ($P<0.05$). Cows fed alfalfa harvested at PM had more digestible CP intake per day than AM fed cows ($P<0.05$). With advancing maturity stages, digestible CP intake of cows decreased, while the efficiency of conversion of feed N into milk N of cows increased linearly ($P<0.05$). Feed efficiency, measured as kilograms milk per kilogram DM intake, was higher at EFAM compared with LBAM ($P<0.05$). Alfalfa harvested at sunset tended to increase milk fat yield ($kg\ d^{-1}$, $P<0.14$). Daily eating time per kilogram DM intake was shorter for cows fed EB alfalfa hay compared with other hays ($P<0.05$). Alfalfa harvested at early bud had higher nutrient availability, but including this forage in dairy cattle diets containing high concentrate level decreased feed efficiency. Alfalfa harvested at sunset had higher in vivo feed digestibility and increased digestible nutrient intake. Overall, alfalfa harvested at early bud at sunset has higher nutrient availability for dairy cattle.

Key words: Alfalfa hay, stage of maturity, diurnal cutting, dairy cattle performance

Growth and pigmentation in Arctic charr (*Salvelinus alpinus*) fed diets with different lipid sources and astaxanthin. H. Lin^{*†} and D. M. Anderson. Nova Scotia Agricultural college, Truro, NS, Canada.

Pigment, a significant dietary cost in salmonid production, is estimated to be over 10% of the cost of diets. Astaxanthin is the pigment of choice. Fish oil is available in finite quantities, so alternate lipid sources need investigation. Tissue deposition of astaxanthin can be related to the nature of dietary lipid as lipid source in diets influences absorption. This study compared the growth performance, rate and level of flesh pigmentation in Arctic charr (initial weight approximately 50 g) fed diets with different lipids; beef tallow (BT), poultry grease (PG) camelina oil (CO), flax oil (FLO) or fish oil (FO) with synthetic astaxanthin added at 60 mg kg^{-1} . A control diet (CON) containing fish oil was compared with diets containing 65% of the FO replaced with one